

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 July 2002 (04.07.2002)


PCT

(10) International Publication Number
WO 02/051982 A2

- (51) International Patent Classification⁷: C12N (71) Applicant (for all designated States except US): ELITRA PHARMACEUTICALS, INC. [US/US]; Suite A, 3510 Dunhill Street, San Diego, CA 92121 (US).
- (21) International Application Number: PCT/US01/50250
- (22) International Filing Date: 21 December 2001 (21.12.2001) (72) Inventors; and (75) Inventors/Applicants (for US only): HASELBECK, Robert [US/US]; 4672 El Cerrito Drive, San Diego, CA 92115 (US). WALL, Dan [US/US]; 3734 Brookshire Street, San Diego, CA 92111 (US). GROSS, Molly [US/US]; #2101, 10954 W. Ocean Air Drive, San Diego, CA 92130 (US).
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/259,434 27 December 2000 (27.12.2000) US (74) Agent: REISMAN, Joseph, M.; Knobbe, Martens, Olson & Bear, LLP, 16th Floor, 620 Newport Center Drive, Newport Beach, CA 92660 (US). 09/948,993 6 September 2001 (06.09.2001) US
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application: 61/230,335 (CIP) (81) Designated States (national): AE, AG, AL, AM, AT, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EC, EE, Filed on 6 September 2000 (06.09.2000)

[Continued on next page]

(54) Title: BACTERIAL PROMOTERS AND METHODS OF USE



(57) Abstract: Compositions and methods are disclosed herein that relate to the development of fusion promoters for regulating gene expression in bacteria. Embodiments include fusion promoters comprising one or more operators linked to a promoter that is modified to have altered activity in Gram-positive organisms. Vectors and cells containing these fusion promoters are also described. Other embodiments include, methods of using these fusion promoters to regulate nucleic acid and/or polypeptide expression, methods of using these fusion promoters to identify proliferation-required genes, and methods of using these fusion promoters to identify molecules having potential antibiotic activity.

BEST AVAILABLE COPY

WO 02/051982 A2



EE (utility model), ES, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished upon receipt of that report
- entirely in electronic form (except for this front page) and available upon request from the International Bureau

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

BACTERIAL PROMOTERS AND METHODS OF USE

RELATED APPLICATIONS

5 This application claims priority to United States Provisional Patent Application 60/259434, filed December 27, 2000, by Robert Haselbeck, and entitled "BACTERIAL PROMOTERS AND METHODS OF USE" and United States Patent Application Serial No. 09/948,993, filed September 6, 2001, by Daniel Tishkoff, and entitled "RAPID METHOD FOR REGULATING GENE EXPRESSION", which
10 claims priority to United States Provisional Patent Application Serial No. 60/230,335, filed September 6, 2000, by Daniel Tishkoff, and entitled "RAPID REPLACEMENT OF GENOMIC PROMOTERS TO GENERATE STRAINS FOR USE IN A CELL-BASED ASSAY FOR ANTIBIOTICS", the disclosures of which are incorporated herein by reference in their entirety.

15

FIELD OF THE INVENTION

 The invention described herein relates to fusion promoters which comprise one or more promoters and one or more operators for directing transcription in Gram-positive bacteria. More specifically, some embodiments relate to fusion promoters
20 comprising a promoter sequence that has been modified so as to alter its transcription activity in Gram-positive organisms linked to a xylose operator, cells having constructs that include such fusion promoters, and methods of using of these manufactures.

25

BACKGROUND OF THE INVENTION

 Genetic information is expressed in bacteria by a series of steps involving transcription of DNA into mRNA and the subsequent translation of the mRNA into polypeptides or proteins. The expression of encoded information is initiated at a region on the DNA molecule called a "promoter", onto which RNA polymerase
30 recognizes and, thereby, initiates transcription. Promoters, which usually precede

protein-coding nucleotide sequences ("structural genes"), vary in "strength" depending on the ability to recognize and bind RNA polymerase in a manner facilitating the initiation of transcription.

The regulation of transcription levels from certain promoters in bacteria can be facilitated by "operators" that are positioned adjacent to or within the promoter. Operator sequences generally function by interacting with modulator proteins, which either activate or repress the transcriptional activity of the promoter. Thus, the inherent strength of the promoter and the amount of regulation that can be imposed upon the promoter by an associated operator sequence interacting with the modulator protein contribute significantly to the control of transcription.

Current techniques in molecular biology enable the transfer to bacteria of exogenous nucleic acids, which one desires to transcribe. Successful transcription of the transferred nucleic acid in the host microorganism, however, depends on the association of the transferred nucleic acid with a suitable promoter and/or operator sequence. Frequently, the promoter and/or operator that is naturally associated with the nucleic acid to be transcribed will not support desired levels of transcription in the new host and, a new promoter and/or operator sequence must be operably linked to the nucleic acid. A variety of promoters and operators have been studied in this regard. (See *e.g.*, von Gabain, *et al.*, *Proc. Natl. Acad. Sci. (USA)*, 76:189-193 (1979); Casadaban, *et al.*, *J. Molecular Biology*, 138:179-207 (1980); West, *et al.*, *Gene*, 9:175-193 (1980); (Chang *et al.*, *Nature*, 275:615 (1978); Itakura *et al.*, *Science*, 198:1056 (1977); Goeddel *et al.*, *Nature*, 281:544 (1979); and (Goeddel *et al.*, *Nucleic Acids Res.*, 8:4057 (1980); and EPO Application Publication No. 0036776). However, many of the currently available regulatable promoters are designed for use in Gram-negative bacteria, such as *E. coli*. There remains a need for regulatable promoters that function in Gram-positive bacteria. In particular, the need exists for promoters that can be used to achieve the precise levels of nucleic acid and/or protein expression desired for applications such as cell-based gene and drug discovery in Gram-positive bacteria. For example, in the discovery of proliferation-required genes, the expression of a proliferation-required mRNA which is transcribed at a low level

may be sufficiently reduced by the levels of antisense transcript produced from a promoter having a low maximal level of transcriptional activity to inhibit cell growth. However, the expression of a proliferation-required mRNA which is transcribed at higher levels may only be sufficiently reduced by levels of antisense transcript that can

5 produced from a promoter having a high maximal transcriptional activity. As a result, the number of types of proliferation-required genes that can be discovered are determined in part by the transcriptional activity of the promoter used to express the proliferation-inhibiting antisense RNA. Accordingly, there exists a need for a promoter suite having a wide dynamic range of transcriptional activities in Gram-

10 positive bacteria. In methods in which genes required for proliferation of Gram-positive organisms are identified by transcribing an antisense nucleic acid which is complementary to at least a portion of such genes, the endogenous transcription levels of the genes may vary. Accordingly, it is valuable to have a suite of promoters which provide a wide range of antisense nucleic acid transcription levels to ensure that a

15 sufficient amount of antisense nucleic acid is obtained to allow the identification of proliferation-required genes which are normally expressed at high levels in the Gram-positive organism. In drug discovery applications which utilize Gram-positive cells that are sensitized by the expression of proliferation-inhibiting antisense RNAs, the degree to which the cells are sensitized is an important factor in determining the

20 candidate compounds that are discovered. Furthermore, the expression of some proliferation-required target genes may be so greatly reduced by the production of proliferation-inhibiting antisense RNA from a highly active promoter that, rather than becoming sensitized, the cell dies. Accordingly, there exists a need for a promoter suite containing a collection of promoters which provide a wide range of basal

25 transcription levels, maximal transcription levels, and degrees of induction which can be used to obtain a level of antisense nucleic acid which is optimal for the identification of a particular gene as being required for proliferation or optimal for identifying compounds which act on the gene product of a particular gene which is required for proliferation in cell-based assays using sensitized cells as described

herein. Thus, there remains a need for a suite of promoters having the ability to tightly and/or finely regulate transcription in Gram-positive bacteria.

BRIEF SUMMARY OF THE INVENTION

5 Some aspects of the present invention are described in the numbered paragraphs below.

1. An isolated nucleic acid comprising a fusion promoter said fusion promoter comprising at least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at least one operator
10 selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter.

2. The isolated fusion promoter of Claim 1, wherein said at least one promoter is selected from the group consisting of SEQ ID NOs.: 36-45.

15 3. The isolated nucleic acid of Claim 1, wherein said at least one operator is *xylO*.

4. The isolated nucleic acid of Claim 3, wherein said at least one promoter is T5.

20 5. The isolated nucleic acid of Claim 3, further comprising a second operator.

6. The isolated nucleic acid of Claim 5, wherein said second operator is *lacO*.

7. The isolated nucleic acid of Claim 1, wherein said fusion promoter is responsive to an inducer.

25 8. The isolated nucleic acid of Claim 7, wherein said inducer is xylose.

9. The isolated nucleic acid of Claim 1, wherein said fusion promoter is titratable.

10. The isolated nucleic acid of Claim 1, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*,
30 *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium*

teteni, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*,
Lactococcus lactis, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium*
tuberculosis, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus*
epidermidis, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus*
5 *mutans* and any species falling within the genera of any of the above species.

11. The isolated nucleic acid of Claim 1, wherein said at least one gram-positive organism is *Staphylococcus aureus*.

12. The isolated nucleic acid of Claim 1, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

10 13. An isolated nucleic acid comprising a fusion promoter said fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter being linked to at least one operator selected from the group consisting of *xyIO*, *tetO*, *trpO*, *malO* and *lacIO*, wherein said at least one operator is positioned such that binding of at least one repressor to said at
15 least one operator represses transcription from said fusion promoter.

14. The isolated nucleic acid of Claim 13, wherein said at least one operator is *xyIO*.

15. The isolated nucleic acid of Claim 14, wherein said at least one promoter is T5.

20 16. The isolated nucleic acid of Claim 14, further comprising a second operator.

17. The isolated nucleic acid of Claim 16, wherein said second operator is *lacO*.

18. The isolated nucleic acid of Claim 13, wherein said fusion promoter is
25 responsive to an inducer.

19. The isolated nucleic acid of Claim 18, wherein said inducer is xylose.

20. The isolated nucleic acid of Claim 13, wherein said fusion promoter is titratable.

21. The isolated nucleic acid of Claim 13, wherein said at least one gram-
30 positive organism is selected from the group consisting of *Bacillus anthracis*,

Clostridium botulinum, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

22. The isolated nucleic acid of Claim 13, wherein said at least one gram-positive organism is *Staphylococcus aureus*.

23. The isolated nucleic acid of Claim 13, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

24. An isolated fusion promoter comprising one of SEQ ID NO.: 26-35.

25. A vector comprising the isolated nucleic acid of Claim 1.

26. A vector comprising the isolated nucleic acid of Claim 13.

27. A vector comprising the isolated nucleic acid of Claim 24.

28. The vector of Claim 25 further comprising at least one replicon selected from the group consisting of p15a, pC194 and pCT1138.

29. The vector of Claim 25 further comprising a reporter gene operably linked to said fusion promoter.

30. The vector of Claim 29, wherein said reporter gene is *lacL-lacM*.

31. The vector of Claim 30, wherein *lacL-lacM* is derived from *Leuconostoc mesenteroides*.

32. The vector of Claim 25, wherein said at least one operator is *xyIO*.

33. The vector of Claim 32, wherein said at least one promoter is T5.

34. The vector of Claim 32, further comprising a second operator.

35. The vector of Claim 34, wherein said second operator is *lacO*.

36. The vector of Claim 25, wherein said fusion promoter is responsive to an inducer.

37. The vector of Claim 36, wherein said inducer is xylose.

38. The vector of Claim 25, wherein said fusion promoter is titratable.

39. The vector of Claim 25, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*,
5 *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

40. The vector of Claim 25, wherein said at least one gram-positive
10 organism is *Staphylococcus aureus*.

41. The vector of Claim 25, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

42. The vector of Claim 25, further comprising a random fragment of a microbial genome operably linked to said fusion promoter.

15 43. The vector of Claim 25, further comprising a nucleic acid that encodes a peptide, wherein said nucleic acid is operably linked to said fusion promoter.

44. The vector of Claim 25, further comprising a nucleic acid that is complementary to a portion of a microbial genome, wherein said nucleic acid is operably linked to said fusion promoter.

20 45. The vector of Claim 25, further comprising a nucleic acid that encodes a molecule that inhibits the proliferation of microbe, wherein said nucleic acid is operably linked to said fusion promoter.

46. A host cell comprising the nucleic acid of Claim 1.

47. A host cell comprising the nucleic acid of Claim 13

25 48. A method of expressing a biomolecule in a cell said method comprising:

(a) introducing a construct into a cell said construct comprising a fusion promoter operably linked to a nucleic acid encoding a biomolecule, said fusion promoter comprising at least one promoter that is modified to have
30 altered activity in at least one gram-positive organism said promoter being

linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter; and

- 5 (b) transcribing said nucleic acid encoding a biomolecule from said fusion promoter.

49. The method of Claim 48, wherein said at least one operator is *xylO*.

50. The method of Claim 49, wherein said at least one promoter is T5.

51. The method of Claim 49, wherein said fusion promoter further
10 comprises a second operator.

52. The method of Claim 51, wherein said second operator is *lacO*.

53. The method of Claim 48, wherein said fusion promoter is responsive to an inducer.

54. The method of Claim 53, wherein said inducer is xylose.

- 15 55. The method of Claim 48, wherein said fusion promoter is titratable.

56. The method of Claim 48, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*,
20 *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

57. The method of Claim 48, wherein said at least one gram-positive
25 organism is *Staphylococcus aureus*.

58. The method of Claim 48, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

59. The method of Claim 48, wherein said nucleic acid is a random fragment of a microbial genome.

60. The method of Claim 48, wherein said biomolecule is a peptide or nucleic acid.

61. The method of Claim 48, wherein said biomolecule is a nucleic acid that is complementary to a portion of a microbial genome.

5 62. The method of Claim 48, wherein said biomolecule is a nucleic acid that encodes a molecule that inhibits the proliferation of a microbe.

63. A biomolecule produced by the method of Claim 48.

64. A method for identifying genes involved in cellular proliferation said method comprising the steps of:

10 (a) introducing into cells of a cell population a construct comprising an inducible fusion promoter operably linked to a nucleic acid, said fusion promoter comprising at least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *λcI*O, wherein said at least one operator is positioned such that binding of
15 at least one repressor to said at least one operator represses transcription from said fusion promoter;

(b) inducing transcription of said nucleic acid from said inducible fusion promoter;

20 (c) identifying the cells in said cell population whose proliferation is reduced in response to the induction of transcription of said nucleic acid; and

(d) identifying the gene from a cell identified in step (c) to which at least a portion of said nucleic acid is complementary.

25 65. A method for identifying genes involved in cellular proliferation said method comprising the steps of:

(a) introducing into cells of a cell population a construct comprising an inducible fusion promoter operably linked to a nucleic acid, said fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter being linked to at least
30 one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and

lacIO, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;

5 (b) inducing transcription of said nucleic acid from said inducible fusion promoter;

(c) identifying the cells in said cell population whose proliferation is reduced in response to the induction of transcription of said nucleic acid; and

(d) identifying the gene from a cell identified in step (c) to which at least a portion of said nucleic acid is complementary.

10 66. The method of Claim 64, wherein said nucleic acid is random fragment of a microbial genome.

67. The method of Claim 64, wherein said nucleic acid encodes an aptamer.

15 68. The method of Claim 64, wherein said nucleic acid encodes a transcript that is complementary to a portion of a microbial genome.

69. The method of Claim 64, wherein said nucleic acid encodes a molecule that inhibits the proliferation of a microbe.

70. The method of Claim 64, wherein said at least one operator is *xylO*.

71. The method of Claim 70, wherein said fusion promoter is T5.

20 72. The method of Claim 70, wherein said fusion promoter further comprises a second operator.

73. The method of Claim 72, wherein said second operator is *lacO*.

74. The method of Claim 64, wherein said inducer is xylose.

75. The method of Claim 64, wherein said fusion promoter is titratable.

25 76. The method of Claim 64, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus*

30

epidermidis, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

77. The method of Claim 64, wherein said at least one gram-positive organism is *Staphylococcus aureus*.

5 78. The method of Claim 64, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

79. A method for identifying genes involved in cellular proliferation said method comprising the steps of:

10 (a) introducing into the genome of a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid, said fusion promoter comprising at least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *lacIO*, wherein said at least one operator is positioned such that binding of
15 at least one repressor to said at least one operator represses transcription from said fusion promoter;

20 (b) comparing the proliferation of said cell cultured in the presence of a first concentration of an inducer that induces transcription from said fusion promoter with the proliferation of said cell cultured in the presence of a concentration of said inducer that is less than said first concentration, wherein a difference in proliferation indicates that said fusion promoter modulates a gene that is required for proliferation; and

 (c) identifying the gene that is modulated by said fusion promoter.

25 80. A method for identifying genes involved in cellular proliferation said method comprising the steps of:

30 (a) introducing into the genome of a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid, said fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and

lacIO, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;

5 (b) comparing the proliferation of said cell cultured in the presence of a first concentration of an inducer that induces transcription from said fusion promoter with the proliferation of said cell cultured in the presence of a concentration of said inducer that is less than said first concentration, wherein a difference in proliferation indicates that said fusion promoter modulates a gene that is required for proliferation; and

10 (c) identifying the gene that is modulated by said fusion promoter.

81. The method of Claim 79, wherein said nucleic acid is random fragment of a microbial genome.

82. The method of Claim 79, wherein said nucleic acid encodes an aptamer.

15 83. The method of Claim 79, wherein said nucleic acid encodes a transcript that is complementary to a portion of a microbial genome.

84. The method of Claim 79, wherein said nucleic acid encodes a molecule that inhibits the proliferation of a microbe.

85. The method of Claim 79, wherein said at least one operator is *xyIO*.

20 86. The method of Claim 85, wherein said at least one promoter is T5.

87. The method of Claim 85, wherein said fusion promoter further comprises a second operator.

88. The method of Claim 87 wherein said second operator is *lacO*.

89. The method of Claim 79, wherein said inducer is xylose.

25 90. The method of Claim 79, wherein said fusion promoter is titratable.

91. The method of Claim 79, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*,
30 *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium*

tuberculosis, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

5 92. The method of Claim 79, wherein said at least one gram-positive organism is *Staphylococcus aureus*.

 93. The method of Claim 79, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

 94. A method for identifying a compound which reduces the activity or level of a gene product required for proliferation of a cell said method comprising:

10 (a) introducing into a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid that is complementary to at least a portion of a proliferation-required gene, said fusion promoter comprising at least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at least one operator
15 selected from the group consisting of *xyiO*, *tetO*, *trpO*, *malO* and *λclO*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;

 (b) sensitizing said cell by inducing transcription from said fusion
20 promoter;

 (c) contacting said sensitized cell with a compound; and

 (d) determining the degree to which said compound inhibits proliferation of said sensitized cell relative to a cell which has not been sensitized.

25 95. A method for identifying a compound which reduces the activity or level of a gene product required for proliferation of a cell said method comprising:

 (a) introducing into a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid that is complementary to at least a portion of a proliferation-required gene, said fusion promoter comprising at
30 least one promoter selected from the group consisting of T5, CP25, P32, P59,

P1P2 and PL, said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *λclO*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;

5 (b) sensitizing said cell by inducing transcription from said fusion promoter;

(c) contacting said sensitized cell with a compound; and

(d) determining the degree to which said compound inhibits proliferation of said sensitized cell relative to a cell which has not been sensitized.

10

96. The method of Claim 94, wherein said at least one operator is *xylO*.

97. The method of Claim 96, wherein said at least one promoter is T5.

98. The method of Claim 96, wherein said fusion promoter further comprises a second operator.

15

99. The method of Claim 98, wherein said second operator is *lacO*.

100. The method of Claim 94, wherein said inducer is xylose.

101. The method of Claim 94, wherein said fusion promoter is titratable.

102. The method of Claim 94, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylosis*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

25

103. The method of Claim 94, wherein said at least one gram-positive organism is *Staphylococcus aureus*.

104. The method of Claim 94, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

30

105. A compound identified using the method of Claim 94.

106. A method for inhibiting the activity or expression of a gene in an operon required for proliferation said method comprising:

- 5 (a) introducing into a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid that is complementary to at least a portion of a proliferation-required operon, said fusion promoter comprising at least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter; and
- 10 (b) inducing transcription from said fusion promoter.

107. A method for inhibiting the activity or expression of a gene in an operon required for proliferation said method comprising:

- 15 (a) introducing into a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid that is complementary to at least a portion of a proliferation-required operon, said fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter; and
- 20 (b) inducing transcription from said fusion promoter.

108. The method of Claim 106, wherein said at least one operator is *xylO*.

25 109. The method of Claim 108, wherein said at least one promoter is T5.

110. The method of Claim 108, wherein said fusion promoter further comprises a second operator.

111. The method of Claim 110, wherein said second operator is *lacO*.

112. The method of Claim 106, wherein said inducer is xylose.

30 113. The method of Claim 106, wherein said fusion promoter is titratable.

114. The method of Claim 106, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*,
5 *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

115. The method of Claim 106, wherein said at least one gram-positive
10 organism is *Staphylococcus aureus*.

116. The method of Claim 106, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

117. A method for identifying the biological pathway in which a proliferation-required gene or its gene product lies said method comprising:

15 (a) introducing into a test cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid that is complementary to at least a portion of a proliferation-required gene, said fusion promoter comprising at least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at least one
20 operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;

(b) inhibiting the activity of said proliferation-required gene or gene
25 product in said test cell by inducing transcription from said fusion promoter;

(c) contacting said test cell with a compound known to inhibit growth or proliferation of a cell, wherein the biological pathway on which said compound acts is known; and

(d) determining the degree to which said compound inhibits proliferation of said test cell relative to a cell which has not been contacted with said compound.

118. A method for identifying the biological pathway in which a proliferation-required gene or its gene product lies said method comprising:

(a) introducing into a test cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid that is complementary to at least a portion of a proliferation-required gene, said fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *lacO*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;

(b) inhibiting the activity of said proliferation-required gene or gene product in said test cell by inducing transcription from said fusion promoter;

(c) contacting said test cell with a compound known to inhibit growth or proliferation of a cell, wherein the biological pathway on which said compound acts is known; and

(d) determining the degree to which said compound inhibits proliferation of said test cell relative to a cell which has not been contacted with said compound.

119. The method of Claim 117, wherein said at least one operator is *xylO*.

120. The method of Claim 119, wherein said at least one promoter is T5.

121. The method of Claim 119, wherein said fusion promoter further comprises a second operator.

122. The method of Claim 121, wherein said second operator is *lacO*.

123. The method of Claim 117, wherein said inducer is xylose.

124. The method of Claim 117, wherein said fusion promoter is titratable.

125. The method of Claim 117, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*,
5 *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

126. The method of Claim 117, wherein said at least one gram-positive
10 organism is *Staphylococcus aureus*.

127. The method of Claim 117, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

128. A method for determining the biological pathway on which a test compound acts said method comprising:

15 (a) introducing into a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid that is complementary to at least a portion of a proliferation-required gene, wherein the biological pathway in which said gene or product of said gene lies is known, and wherein said fusion promoter comprises at least one promoter that is modified to have altered
20 activity in at least one gram-positive organism said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;

25 (b) inhibiting the activity of said proliferation-required in said cell by inducing transcription from said fusion promoter;

(c) contacting said cell with a test compound; and

(d) determining the degree to which said test compound inhibits proliferation of said cell relative to a cell containing a which does not contain
30 said construct.

129. A method for determining the biological pathway on which a test compound acts said method comprising:

- 5 (a) introducing into a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid that is complementary to at least a portion of a proliferation-required gene, wherein the biological pathway in which said gene or product of said gene lies is known, and wherein said fusion promoter comprises at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *lacIO*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;
- 10 (b) inhibiting the activity of said proliferation-required in said cell by inducing transcription from said fusion promoter;
- 15 (c) contacting said cell with a test compound; and
- (d) determining the degree to which said test compound inhibits proliferation of said cell relative to a cell containing a which does not contain said construct.

130. The method of Claim 128, wherein said at least one operator is *xylO*.

20 131. The method of Claim 130, wherein said at least one promoter is T5.

132. The method of Claim 130, wherein said fusion promoter further comprises a second operator.

133. The method of Claim 132, wherein said second operator is *lacO*.

134. The method of Claim 128, wherein said inducer is xylose.

25 135. The method of Claim 128, wherein said fusion promoter is titratable.

136. The method of Claim 128, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*,
30 *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium*

tuberculosis, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

5 137. The method of Claim 128, wherein said at least one gram-positive organism is *Staphylococcus aureus*.

138. The method of Claim 128, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

139. A method of manufacturing an antibiotic comprising the steps of:

10 (a) introducing into a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid that is complementary to at least a portion of a proliferation-required gene, said fusion promoter comprising at least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *lacIO*,
15 wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;

(b) sensitizing said cell by inducing transcription from said fusion promoter;

20 (c) contacting said sensitized cell with a compound;

(d) identifying a compound which substantially inhibits the proliferation of said sensitized cell relative to a cell which has not been sensitized; and

(e) manufacturing the compound so identified.

25 140. A method of manufacturing an antibiotic comprising the steps of:

(a) introducing into a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid that is complementary to at least a portion of a proliferation-required gene, said fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter being linked to at least one operator selected from
30

the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *λclO*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;

5 (b) sensitizing said cell by inducing transcription from said fusion promoter;

(c) contacting said sensitized cell with a compound;

(d) identifying a compound which substantially inhibits the proliferation of said sensitized cell relative to a cell which has not been sensitized; and

10 (e) manufacturing the compound so identified.

141. The method of Claim 139, wherein said at least one operator is *xylO*.

142. The method of Claim 141, wherein said at least one promoter is T5.

143. The method of Claim 141, wherein said fusion promoter further comprises a second operator.

15 144. The method of Claim 143, wherein said second operator is *lacO*.

145. The method of Claim 139, wherein said inducer is xylose.

146. The method of Claim 139, wherein said fusion promoter is titratable.

147. The method of Claim 139, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

148. The method of Claim 139, wherein said at least one gram-positive organism is *Staphylococcus aureus*.

149. The method of Claim 139, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

150. A method for identifying a gene which is required for proliferation of a prokaryotic cell said method comprising the steps of:

- 5 (a) replacing the native promoter of a gene in the chromosome of a prokaryotic cell having an enhanced frequency of homologous recombination with a regulatable fusion promoter comprising at least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO*, *lacO* and *λclO*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter; and
- 10 (b) identifying cells in which the extent of proliferation of said cell when said fusion promoter is active at a first level is substantially different than the extent of proliferation of said cell when said fusion promoter is active at a second level, said first level being greater than said second level.

15 151. A method for identifying a gene which is required for proliferation of a prokaryotic cell said method comprising the steps of:

- 20 (a) replacing the native promoter of a gene in the chromosome of a prokaryotic cell having an enhanced frequency of homologous recombination with a regulatable fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO*, *lacO* and *λclO*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter; and
- 25 (b) identifying cells in which the extent of proliferation of said cell when said fusion promoter is active at a first level is substantially different than the extent of proliferation of said cell when said fusion promoter is active at a second level, said first level being greater than said second level.

30 152. The method of Claim 150, wherein said fusion promoter is an inducible promoter.

153. The method of Claim 150, wherein the step of replacing said native promoter comprises introducing a linear nucleic acid comprising a 5' portion homologous to a first portion of said native promoter, a 3' portion homologous to a second portion of said native promoter and said fusion promoter disposed between said 5' portion and said 3' portion into said cell such that homologous recombination occurs between said 5' portion and said first portion of said native promoter and between said 3' portion and said second portion of said native promoter.

154. The method of Claim 153, wherein said linear nucleic acid is double stranded.

155. The method of Claim 153, wherein said linear nucleic acid is single stranded.

156. A method for identifying a compound which inhibits the proliferation of a prokaryotic cell said method comprising the steps of:

(a) replacing the native promoter of a gene in the chromosome of a prokaryotic cell having an enhanced frequency of homologous recombination with a regulatable fusion promoter comprising at least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO*, *lacO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter; and

(b) comparing the extent of proliferation of a first sample of said cell in the presence of said compound to the extent of proliferation of a second sample of said cell in the presence of said compound, wherein said first sample of said cell has a reduced activity of said fusion promoter activity relative the activity of said fusion promoter in said second sample of said cell and wherein said compound inhibits the proliferation of said cell if the extent of proliferation of said first sample of said cell is substantially less than the extent of proliferation of said second sample of said cell.

157. A method for identifying a compound which inhibits the proliferation of a prokaryotic cell said method comprising the steps of:

- 5 (a) replacing the native promoter of a gene in the chromosome of a prokaryotic cell having an enhanced frequency of homologous recombination with a regulatable fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter being linked to at least one operator selected from the group consisting of *xytO*, *tetO*, *trpO*, *malO*, *lacO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter; and
- 10 (b) comparing the extent of proliferation of a first sample of said cell in the presence of said compound to the extent of proliferation of a second sample of said cell in the presence of said compound, wherein said first sample of said cell has a reduced activity of said fusion promoter activity relative the activity of said fusion promoter in said second sample of said cell and wherein
- 15 said compound inhibits the proliferation of said cell if the extent of proliferation of said first sample of said cell is substantially less than the extent of proliferation of said second sample of said cell..

158. The method of Claim 156, wherein the step of replacing said native promoter comprises introducing a linear nucleic acid comprising a 5' portion homologous to a first portion of said native promoter, a 3' portion homologous to a second portion of said native promoter and said fusion promoter disposed between said 5' portion and said 3' portion into said cell such that homologous recombination occurs between said 5' portion and said first portion of said native promoter and

20 between said 3' portion and said second portion of said native promoter.

159. The method of Claim 158, wherein said linear nucleic acid is double stranded.

160. The method of Claim 158, wherein said linear nucleic acid is single stranded.

161. A method for identifying a gene which is required for proliferation of a prokaryotic cell said method comprising the steps of:

(a) introducing at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO*, *lacO* and *λc1O* into a prokaryotic cell having an enhanced frequency of homologous recombination such that said at least one operator regulates transcription of a target nucleic acid in the chromosome of said cell; and

(b) identifying cells in which the extent of proliferation of said cell when said target nucleic acid is transcribed at a first level is substantially different than the extent of proliferation of said cell when said target nucleic acid is transcribed at a second level, said first level being greater than said second level.

162. The method of Claim 161, wherein the step of introducing said at least one operator comprises introducing a linear nucleic acid comprising a 5' portion homologous to a first portion of the chromosome of said cell, a 3' portion homologous to a second portion of the chromosome of said cell and said at least one operator disposed between said 5' portion and said 3' portion into said cell such that homologous recombination occurs between the 5' portion and said first portion of the chromosome of said cell and between the 3' portion and said second portion of the chromosome of said cell.

163. The method of Claim 162, wherein said linear nucleic acid is double stranded.

164. The method of Claim 162, wherein said linear nucleic acid is single stranded.

165. A method of identifying a compound which inhibits the proliferation of a prokaryotic cell said method comprising the steps of:

(a) obtaining a prokaryotic cell in which transcription of a nucleic acid required for proliferation of said cell is regulated by at least one operator which has been introduced into the chromosome of said cell said at least one

operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO*, *lacO* and *λcI O*;

(b) sensitizing said cell by growing said cell under conditions in which the level of transcription of said gene is lower than that of a wild type cell;

5 (c) contacting said sensitized cell with said compound; and

(d) determining the degree to which said compound inhibits the growth of said sensitized cell relative to an unsensitized cell.

166. The method of Claim 165, wherein said cell has an enhanced frequency of homologous recombination.

10 167. The method of Claim 165, wherein said linear nucleic acid is single stranded.

168. The method of Claim 165, wherein said linear nucleic acid is double stranded.

15 169. A method of identifying a nucleic acid sequence having promoter activity in *Enterococcus faecalis* said method comprising the steps of:

(a) inserting a candidate nucleic acid into a vector comprising *lacL-lacM* reporter genes such that said candidate nucleic acid is upstream of the *lacL-lacM* reporter genes;

20 (b) introducing said vector comprising said candidate nucleic acid into *Enterococcus faecalis*;

(c) detecting expression of the *lacL-lacM* reporter genes, wherein expression of said *lacL-lacM* reporter genes indicates that said candidate nucleic acid sequence has promoter activity.

25 170. The method of Claim 169, wherein said detecting step comprises measuring β-galactosidase activity.

171. The method of Claim 169, wherein said candidate nucleic acid is a promoter modified to increase activity in a gram-positive organism.

172. The method of Claim 169, wherein said *lacL-lacM* reporter gene is derived from *Leuconostoc mesenteroides*.

30 173. The method of Claim 169, wherein said vector is pEPEF1.

Definitions

The following definitions are provided so as to facilitate the understanding of the invention as set out herein.

5 By "activity against a gene product" is meant having the ability to inhibit the function or to reduce the level or activity of the gene product in a cell. This includes, but is not limited to, inhibiting the enzymatic activity of the gene product or the ability of the gene product to interact with other biological molecules required for its activity, including inhibiting the gene product's assembly into a multimeric structure.

10 By "activity against a protein" is meant having the ability to inhibit the function or to reduce the level or activity of the protein in a cell. This includes, but is not limited to, inhibiting the enzymatic activity of the protein or the ability of the protein to interact with other biological molecules required for its activity, including inhibiting the protein's assembly into a multimeric structure.

15 By "activity against a nucleic acid" is meant having the ability to inhibit the function or to reduce the level or activity of the nucleic acid in a cell. This includes, but is not limited to, inhibiting the ability of the nucleic acid to interact with other biological molecules required for its activity, including inhibiting the nucleic acid's assembly into a multimeric structure.

20 By "activity against a gene" is meant having the ability to inhibit the function or expression of the gene in a cell. This includes, but is not limited to, inhibiting the ability of the gene to interact with other biological molecules required for its activity.

25 By "activity against an operon" is meant having the ability to inhibit the function or reduce the level of one or more products of the operon in a cell. This includes, but is not limited to, inhibiting the enzymatic activity of one or more products of the operon or the ability of one or more products of the operon to interact with other biological molecules required for its activity.

 By "antibiotic" is meant an agent which inhibits the proliferation of a cell or microorganism.

The term "aptamer" as used herein refers to a single-stranded or double-stranded oligodeoxyribonucleotide, oligoribonucleotide or modified derivatives of oligodeoxyribonucleotides or oligoribonucleotides that specifically binds and alters the biological function of a target molecule, which can be a protein, peptide and derivatives thereof.

By "biological pathway" is meant any discrete cell function or process that is carried out by a gene product or a subset of gene products. Biological pathways include anabolic, catabolic, enzymatic, biochemical and metabolic pathways as well as pathways involved in the production of cellular structures such as cell walls. Biological pathways that are usually required for proliferation of cells or microorganisms include, but are not limited to, cell division, DNA synthesis and replication, RNA synthesis (transcription), protein synthesis (translation), protein processing, protein transport, fatty acid biosynthesis, electron transport chains, cell wall synthesis, cell membrane production, synthesis and maintenance, and the like.

The term "expression" is defined as the production of a sense or antisense RNA molecule from a gene, gene fragment, genomic fragment, chromosome, operon or portion thereof. Expression can also be used to refer to the process of peptide or polypeptide synthesis.

An "expression vector" is defined as a vehicle by which a ribonucleic acid (RNA) is transcribed from a nucleic acid carried within the expression vehicle. The expression vector can also contain features that permit translation of a protein product from the transcribed RNA message expressed from the exogenous nucleic acid carried by the expression vector. Accordingly, an expression vector can produce an RNA molecule as its sole product or the expression vector can produce a RNA molecule that is ultimately translated into a protein product.

By "identifying a compound" is meant to screen one or more compounds in a collection of compounds such as a combinatorial chemical library, a natural product library or other library of chemical compounds or to characterize a single compound by testing the compound in a given assay and determining whether it exhibits the desired activity.

By "inducer" is meant an agent, solution or environmental condition which, when placed in contact with a cell or microorganism, increases transcription, or inhibitor and/or promoter clearance/fidelity, from a desired promoter.

By "inhibit activity of a gene or gene product" is meant having the ability to interfere with the function of a gene or gene product in such a way as to decrease expression of the gene, in such a way as to reduce the level or activity of a product of the gene or in such a way as to inhibit the interaction of the gene or gene product with other biological molecules required for its activity. Agents which inhibit the activity of a gene include agents that inhibit transcription of the gene, agents that inhibit processing of the transcript of the gene, agents that reduce the stability of the transcript of the gene, and agents that inhibit translation of the mRNA transcribed from the gene. In microorganisms, agents which inhibit the activity of a gene can act to decrease expression of the operon in which the gene resides or alter the folding or processing of operon RNA so as to reduce the level or activity of the gene product. The gene product can be a non-translated RNA such as ribosomal RNA, a translated RNA (mRNA) or the protein product resulting from translation of the gene mRNA. Of particular utility to the present invention are antisense RNAs that have activities against the operons or genes to which they specifically hybridize.

As used herein, "*lac* operator", "*lac* operator sequence", or "*lacO*" is intended to encompass the operator having SEQ ID NO.: 11 and any other *lac* operator sequences consistent with the intended uses described herein.

As used herein, "*λcl* operator", "*λcl* operator sequence", or "*λclO*" is intended to encompass all lambda operators to which the *λcl* repressor or variants thereof are capable of binding. Such operators include, but are not limited to, *λOR_I* (SEQ ID NO.: 54), *λOR₂* (SEQ ID NO.: 55), *λOR₃* (SEQ ID NO.: 56), *λOL_I* (SEQ ID NO.: 57), *λOL₂* (SEQ ID NO.: 58), *λOL₃* (SEQ ID NO.: 59).

As used herein, "*mal* operator", "*mal* operator sequence" or "*malO*" is intended to encompass the two *mal* operators MalMop (SEQ ID NO.: 60) and MalXop (SEQ ID NO.: 61), as described in Nieto et al., *J. Biol. Chem.* 272,30860-30865

(1997), the disclosure of which is incorporated herein by reference in its entirety, and any other *mal* operator sequences consistent with the intended uses described herein.

As used herein, "nucleic acid" means DNA, RNA, or modified nucleic acids. Thus, the terminology "the nucleic acid of SEQ ID NO: X" or "the nucleic acid comprising the nucleotide sequence" includes both the DNA sequence of SEQ ID NO: X, the complement thereof, and an RNA sequence in which the thymidines in the DNA sequence have been substituted with uridines in the RNA sequence and in which the deoxyribose backbone of the DNA sequence has been substituted with a ribose backbone in the RNA sequence.

As used herein, "polynucleotide" has the same meaning as nucleic acid.

As used herein, "proliferation-inhibiting" encompasses instances where the absence or substantial reduction of a gene transcript and/or gene product completely eliminates cell growth as well as instances where the absence of a gene transcript and/or gene product merely reduces cell growth. A proliferation-inhibiting antisense nucleic acid is one that can cause a reduction of a gene transcript and/or gene product that is sufficient to reduce or eliminate the growth or viability of the cell or microorganism.

As used herein, "proliferation-required" or "required for proliferation" encompasses instances where the absence or substantial reduction of a gene transcript and/or gene product completely eliminates cell growth as well as instances where the absence of a gene transcript and/or gene product merely reduces cell growth. A proliferation-required gene or gene family is one where, in the absence or substantial reduction of a gene transcript and/or gene product, growth or viability of the cell or microorganism is reduced or eliminated.

As used herein, "sub-lethal" means a concentration of an agent below the concentration required to inhibit all cell growth.

As used herein, "tetracycline analog" or "Tc analog" is intended to include compounds which are structurally related to tetracycline and which bind to the Tet repressor with a K_d of at least about 10^{-6} M. Preferably, the tetracycline analog binds with an affinity of about 10^{-9} M or greater. Examples of such tetracycline analogs include, but are not limited to, anhydrotetracycline (atc), doxycycline,

chlorotetracycline, oxytetracycline and others disclosed by Hlavka and Boothe, "The Tetracyclines," in Handbook of Experimental Pharmacology 78, R. K. Blackwood et al. (eds.), Springer-Verlag, Berlin, N.Y., 1985; L. A. Mitscher, "The Chemistry of the Tetracycline Antibiotics", Medicinal Research 9, Dekker, N.Y., 1978; Noyee
5 Development Corporation, "Tetracycline Manufacturing Processes" Chemical Process Reviews, Park Ridge, N.J., 2 volumes, 1969; R. C. Evans, "The Technology of the Tetracyclines", Biochemical Reference Series 1, Quadrangle Press, New York, 1968; and H. F. Dowling, "Tetracycline", Antibiotic Monographs, no. 3, Medical Encyclopedia, New York, 1955. Preferred Tc analog for high-level stimulation of
10 transcription is anhydrotetracycline. For use in prokaryotic organisms, a Tc analog can be chosen which has reduced antibiotic activity compared to Tc. Examples of such Tc analogues are anhydrotetracycline, epioxytetracycline and cyanotetracycline.

As used herein, "Tet repressor" is intended to describe a protein occurring in nature which represses transcription from tet operator sequences in prokaryotic cells in
15 the absence of Tc or Tc analog. Tetracycline repressors of the present invention may be obtained from any class of tetracycline repressor (e.g., TetR(A), TetR(B), TetR(C), TetR(D), TetR(E), TetR(G), TetR(H), TetR(J), TetR(Z), or combinations thereof). The nucleotide sequences of Tet repressors of representative members of the A, B, C, D and E classes are known (e.g., see Waters, S. H. et al. (1983) Nucl. Acids Res
20 11:6089-6105, Hillen, W. and Schollmeier, K. (1983) Nucl. Acids Res. 11:525-539 and Postle, K. et al. (1984) Nucl. Acids Res. 12:4849-4863, Unger, B. et al. (1984) Gene 31: 103-108, Unger, B. et al. (1984) Nucl Acids Res. 12:7693-7703 and Tovar, K. et al. (1988) Mol. Gen. Genet. 215:76-80, respectively).

As used herein, "tet operator", "tet operator sequence" or "tetO" is intended to
25 encompass all classes of tet operator sequences, e.g. (e.g., TetO(A) (SEQ ID NO.: 62), TetO(B) (SEQ ID NO.: 63), TetO(C) (SEQ ID NO.: 64), TetO(D) (SEQ ID NO.: 65), TetO(E) (SEQ ID NO.: 66), TetO(G), TetO(H), TetO(J), and TetO(Z). Nucleotide sequences of these five classes of tet operators are described in Waters, S. H. et al. (1983) cited supra, Hillen, W. and Schollenmeier, K. (1983) cited supra, Stuber, D.

and Bujard, H. (1981) Proc. Natl. Acad. Sci. USA 78:167-171, Unger, B. et al. (1984) cited supra and Tovar, K. et al. (1988) cited supra.

As used herein, "*trp* operator", "*trp* operator sequence" or "*trpO*" is intended to encompass all classes of *trp* operator sequences consistent with the intended uses described herein. One example is the *trp* operator of SEQ ID NO.: 67.

As used herein, "*xyl* operator", "*xyl* operator sequence" or "*xylO*" is intended to encompass all classes of *xyl* operator sequences consistent with the intended uses described herein. One example is the *xyl* operator of SEQ ID NO.: 68.

10

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a sequence map showing the Xyl-T5 fusion promoter (SEQ ID NO.: 1) showing the following elements: (1) -45 AT Box; (2) -35 RNA polymerase recognition site; (3) a 17 bp intervening sequence that can be replaced by a second operator; (4) -10 RNA polymerase recognition site; (5) the *xyl* operator; (6) multiple cloning sequence; (7A) first *rrnB* terminator; and (7B) second *rrnB* terminator.

FIG. 2 is a sequence map showing the Xyl-CP25 fusion promoter (SEQ ID NO.: 2) showing the following elements: (1) a portion of the *xylR* gene; (2) -35 RNA polymerase recognition site; (3) a 17 bp intervening sequence that can be replaced by a second operator; (4) -10 RNA polymerase recognition site; and (5) the *xyl* operator. Also shown are recognition sites for the restriction enzymes (a) *RsrII* and (b) *XhoI*.

FIG. 3A illustrates one embodiment of the present promoter replacement method in which a linear double stranded promoter replacement construct comprising a nucleic acid encoding a selectable marker, a transcriptional terminator, and a regulatable fusion promoter is inserted into the chromosome of a cell expressing the RecE and RecT gene products of the *Rac* prophage or gene products having analogous functions.

FIG. 3B illustrates one embodiment of the present promoter replacement method in which a linear single stranded promoter replacement construct comprising a 5' portion homologous to a target sequence, a 3' portion homologous to a target

sequence and a regulatable fusion promoter is inserted into the chromosome of a cell expressing the λ Beta and Gam proteins.

FIG. 4A illustrates one embodiment of the operator insertion method in which a linear double stranded regulatory element insertion construct comprising a nucleic acid encoding a selectable marker, a transcriptional terminator, and an operator is inserted into the chromosome of cells expressing the RecE and RecT gene products of the Rac prophage or gene products having analogous functions.

FIG. 4B illustrates one embodiment of the operator insertion method in which a linear single stranded regulatory element insertion construct comprising a 5' portion homologous to a target sequence, a 3' portion homologous to a target sequence and an operator is inserted into the chromosome of a cell expressing the λ Beta and Gam proteins.

FIG. 5 shows a plasmid map of the pXyl-T5 P15a *Staphylococcus* expression vector.

FIG. 6 shows a plasmid map for the pEPEF-X series of *Enterococcus* expression vectors.

FIG. 7 shows a bar graph depicting the ratio of RNA measured from xylose induced cells versus non-induced cells. RNA measurements were made in real time by RT-PCR and the samples were normalized for loading by using parallel 16S rRNA controls. Samples were prepared and analyzed in duplicate and the averages are plotted.

FIG. 8 shows a bar graph comparing the induced and uninduced activities of the Xyl-CP25, Xyl-P32, Xyl-P59 and Xyl-P1P2 fusion promoters in *Enterococcus faecalis*. The activity of each fusion promoter was determined by measuring the β -galactosidase activity produced by the expression of *lacL-lacM* reporter genes operably linked downstream of each fusion promoter.

FIG. 9 shows a bar graph comparing the induced and uninduced activities of the Xyl-T5 and Xyl-CP25 fusion promoters in *Enterococcus faecalis*. A comparison of the induced and uninduced activities of the Xyl-T5 fusion promoter in *Staphylococcus aureus* and *Enterococcus faecalis* is also shown. The activity of each

fusion promoter was determined by measuring the β -galactosidase activity produced by the expression of *lacL-lacM* reporter genes operably linked downstream of each fusion promoter.

FIG. 10 shows a bar graph depicting the activity of the Xyl-P59 fusion promoter over a range of xylose concentrations from 0 to 10%. The activity of Xyl-P59 was determined by measuring the β -galactosidase activity produced by the expression of *lacL-lacM* reporter genes operably linked downstream of this fusion promoter.

FIG. 11 shows a bar graph comparing the induced and uninduced activities of the Xyl-P59, Xyl-PL, and modified Xyl-PL (Xyl-PL10) fusion promoters in *Enterococcus faecalis*. The activity of each fusion promoter was determined by measuring the β -galactosidase activity produced by the expression of *lacL-lacM* reporter genes operably linked downstream of each fusion promoter.

FIG. 12 is an IPTG dose response curve in *E. coli* transformed with an IPTG-inducible plasmid containing either an antisense clone to the *E. coli* ribosomal protein *rplW* (AS-*rplW*) which is required for protein synthesis and essential for cell proliferation, or an antisense clone to the *elaD* (AS-*elaD*) gene which is not known to be involved in protein synthesis and which is also essential for proliferation.

FIG. 13A is a tetracycline dose response curve in *E. coli* transformed with an IPTG-inducible plasmid containing antisense to *rplW* (AS-*rplW*) in the absence (0) or presence of IPTG at concentrations that result in 20% and 50% growth inhibition.

FIG. 13B is a tetracycline dose response curve in *E. coli* transformed with an IPTG-inducible plasmid containing antisense to *elaD* (AS-*elaD*) in the absence (0) or presence of IPTG at concentrations that result in 20% and 50% growth inhibition.

FIG. 14 is a graph showing the fold increase in tetracycline sensitivity of *E. coli* transfected with antisense clones to essential ribosomal proteins L23 (AS-*rplW*) and L7/L12 and L10 (AS-*rplLrplJ*). Antisense clones to genes known to not be directly involved in protein synthesis, *atpB/E* (AS-*atpB/E*), *visC* (AS-*visC*), *elaD* (AS-*elaD*), *yohH* (AS-*yohH*), are much less sensitive to tetracycline.

FIG. 15 illustrates the results of an assay in which *Staphylococcus aureus* cells expressing an antisense nucleic acid complementary to the *gyrB* gene encoding the β subunit of gyrase were contacted with several antibiotics whose targets were known.

5

DETAILED DESCRIPTION OF THE INVENTION

The invention described herein involves the construction and use of fusion promoters that can be tightly and/or finely regulated (e.g., with a *xyl*, *tet*, *mal*, *lac*, *trp*, *λ c1*, or other suitable operators), constructs and cells having these promoters, methods of expression of RNAs, peptides and proteins, methods to identify genes essential for the growth of bacteria, and methods to identify compounds that inhibit bacterial growth.

Some embodiments of this invention contemplate the preparation of several different bacterial fusion promoters which contain at least one operator associated with a promoter sequence. The at least one operator is positioned such that the binding of at least one repressor to the operator represses transcription from the promoter. The promoter sequence may be a wildtype promoter obtained from a bacterium, virus or other microorganism or it may be a promoter sequence that has been modified to have altered activity in Gram-positive organisms. One embodiment, for example, is a fusion promoter comprising a promoter sequence modified to have altered activity in at least one Gram-positive organism linked to one or more operators including, but not limited to, the xylose operator ("*xylO*"), the tetracycline operator ("*tetO*"), the maltose operator ("*malO*"), lactose operator ("*lacO*"), the tryptophan operator ("*trpO*"), and the lambda C1 operator (" *λ c1O*"). These operators can be induced by the following inducers, respectively: xylose or analog thereof, tetracycline or analog thereof, maltose or analog thereof, lactose or analog thereof, tryptophan or analog thereof, and temperature.

In some embodiments, fusion promoters based on the bacteriophage T5 promoter and the xylose operator are contemplated. The T5 promoter has been shown to be capable of directing the efficient expression of genes primarily in *Escherichia coli*. (See e.g., Gentz et al., Proc. Natl. Acad. Sci. USA 78:4936-4940

30

(1981); Rommens et al., *Nucleic Acids Res.* 11:5921-5940 (1983); Wang et al., *Gene* 242:105-114 (2000); and U.S. Pat Nos. 4,495,280 and 4,689,406) and the xylose operator/repressor system has been shown to regulate expression in some Gram-positive bacteria. (See e.g., Dahl et al., *J. Mol. Biol.* 243:413-424 (1994);
5 Sizemore et al., *J. Bacteriol.* 174:3042-3048 (1992); Wieland et al., *Gene* 158:91-96 (1995); Kreuzer et al., *J. Bacteriol.* 171:3840-3845 (1989); Lokman et al., *J. Bacteriol.* 179:5391-5397 (1997); and Erbeznik et al., *J. Bacteriol.* 180:1103-1109 (1998)).

In a preferred embodiment, *xylO* is operably linked to a T5 promoter such that
10 transcription from the fusion promoter is inducible by agents that inhibit the binding of the xylose repressor (encoded by *xylR*) to the xylose operator (*xylO*). The *xylO* can be the *xylO* from *S. xylois* or *B. subtilis* or a number of other bacteria. This embodiment, referred to as the Xyl-T5 promoter, can also contain several other elements including, but not limited to, AT-rich boxes (AT Box) that enhance the
15 efficiency of gene expression in Gram-positive bacteria. The use of an AT-Box-enhanced T5 promoter has been described by D.J. Henner in "Expression of Heterologous Genes in *Bacillus subtilis*" *Methods Enzymol.*, 195:199 (1990), the disclosure of which is incorporated herein by reference in its entirety.

In some embodiments fusion promoters containing one or more AT Boxes are
20 contemplated. The AT Box is positioned such that it increases the level of transcription from the fusion promoter in Gram-positive bacteria. For the Xyl-T5 embodiment, a consensus -45 AT Box is shown in FIG. 1 at position (1). Many variations of the AT Box can be used. For example, the AT Box can have more or less adenine or thymine nucleotides or different sequences or combinations of adenine
25 and thymine molecules. In this respect, the diversity of the AT box is limited only by the ability of this AT rich region to facilitate transcription in Gram-positive bacteria. In other embodiments of the present invention, fusion promoters constructed from T5 promoter derivatives which have been modified to have altered activity in Gram-positive organisms are contemplated.

A preferred embodiment of the Xyl-T5 fusion promoter is shown in FIG.1. In italics, at positions (2) and (4) of FIG. 1 are shown the -35 and -10 RNA polymerase recognition sites (-35 Box and -10 Box), respectively. The 17 nucleotide long element shown at position (3) is a site in which a second operator can be inserted if desired.

5 The second operator is positioned such that the binding of a repressor to the second operator represses transcription from the promoter. In some embodiments, the second operator may be the *xylO*, *tetO*, *malO*, *lacO*, *trpO*, or *λc1O*. At position (5) is shown the *xyl* operator (*xylO*), which is placed just downstream of the -10 Box. The distance between *xylO* and the T5 promoter can be made shorter or longer provided the ability
10 to sustain repression of the T5 promoter is maintained. In some embodiments, a multiple cloning sequence (designated at position (6)) is also inserted downstream of the *xyl* operator to allow nucleic acids to be transcribed from the promoter when operably linked to the promoter. Many different multiple cloning sequences can be inserted at position (6) and the multiple cloning sequence shown is only one example.
15 Additionally, some embodiments can include a terminator sequence, to facilitate termination of the RNA transcribed from the promoter at a desired position (e.g., the first *rrnB* terminator shown at position (7A) and/or the second *rrnB* terminator shown at position (7B)). Preferably, the terminator sequence allows for Rho independent transcription termination.

20 The present invention also contemplates fusion promoters based on promoter sequences discovered in *Lactococcus lactis* linked to at least one operator. For example, the *Lactococcus lactis* promoters CP25 (Jensen and Hammer, *Appl. Environ. Microbiol.* 64:82-87 (1998), the disclosure of which is incorporated herein by reference in its entirety), P32 and P59 (Bruberg et al., *Appl. Microbiol. Biotechnol.*
25 42:108-115 (1994), the disclosure of which is incorporated herein by reference in its entirety), P1P2 (Nilsson and Johansen, *Biochimica Biophysica Acta.* 1219:141-144 (1994), the disclosure of which is incorporated herein by reference in its entirety), and PL (Madsen et al., *Appl. Environ. Microbiol.* 67:1128-1139, the disclosure of which is incorporated herein by reference in its entirety) can be fused to to at least one operator.
30 In some embodiments of the present invention, fusion promoters constructed from

derivatives of the above *Lactococcus lactis* sequences which have been modified to have altered activity in Gram-positive organisms are contemplated.

It will be appreciated that a regulatable fusion promoter suite can be constructed which provides a wide dynamic range of both basal and maximal transcriptional activities. Each fusion promoter within this suite can be tightly and
5 finely regulated over its own characteristic range of transcriptional activity. Furthermore, the regulatable fusion promoters within the suite can comprise different promoters operably linked to the same operators thus providing for a common means of regulation of the promoters within the suite. In some embodiments, the promoters
10 within the set of regulatable fusion promoters within the suite can be linked to at least one operator contained in a promoterless expression vector that is capable of replicating in one or more Gram-positive organisms. Accordingly, a regulatable fusion promoter having an appropriate basal level and/or maximal level of transcriptional activity can be selected from the suite for use in the appropriate
15 application.

In some embodiments, fusion promoters based on promoter sequences discovered in *Lactococcus* actively promote transcription in *Enterococcus faecalis* and other Gram-positive organisms. One embodiment is the Xyl-CP25 fusion promoter shown in FIG. 2. Position (1) of FIG. 2 shows a portion of the *xylR* gene located
20 upstream from the fusion promoter. Positions (2) and (4) show the -35 and -10 Boxes of the CP25 promoter sequence, respectively. The 17 nucleotide long element shown at position (3) is a site in which a second operator can be inserted if desired. The second operator is positioned such that the binding of a repressor to the second operator represses transcription from the promoter. In some embodiments, the second
25 operator may be the *xylO*, *tetO*, *malO*, *lacO*, *trpO*, or *λclO* operator. At position (5) is shown the *xyl* operator (*xylO*), which is placed just downstream of the -10 Box. The distance between *xylO* and the CP25 promoter can be made shorter or longer provided the ability to sustain repression of the CP25 promoter is maintained. The positions of unique RsrII and XhoI recognition sites flanking the promoter are indicated as (a) and
30 (b), respectively. In some embodiments, the RsrII and XhoI sites can be used to

exchange the CP25 promoter with other promoter sequences, such as promoters selected from SEQ ID NOs.: 36 - 45.

5 In other embodiments of the fusion promoters described herein, elements such as multiple cloning sites (MCS) and transcription terminators can be linked downstream. Preferably, the terminator sequence allows for Rho independent transcription termination. For example the *rrnB* terminator (See e.g., Orosz *et al.*, *Eur. J. Biochem.*, 201:653-659 (1991), the disclosure of which is incorporated by reference in its entirety) may be utilized.

10 Some fusion promoters of the present invention function in *Staphylococcus aureus* and *Enterococcus faecalis*. Other fusion promoters described herein function in Gram-positive organisms such as *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia*
15 *asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species.

In another aspect of the invention, two or more operators are linked to a promoter. The operators are positioned such that binding of repressors to the
20 operators represses transcription from the promoter. The operators may be identical or different. By controlling gene expression with multiple operators (e.g., using a plurality of operators that can be selectively modulated in specific organisms), genetic information can be safely transferred between bacterial species and expression can be selectively induced and regulated. For example, repressors that bind to one or more of
25 the operators may be expressed in a first bacterial species or strain but not in a second species. Advantageously, some embodiments maintain tight repression of a potentially lethal transcript in a first species or strain that is used to amplify the construct but the repression can be efficiently removed in a second species or strain in which transcription from the promoter is desired.

One embodiment, for example, concerns a fusion promoter that selectively enables the repression of transcription in *Escherichia coli* and *Staphylococcus* species. The "Xyl/Lac-T5" promoter comprises the *lacO* operably linked to the Xyl-T5 promoter so that transcription of the genetic information associated with the promoter can be induced in *Staphylococcus* by adding xylose or an analog thereof and/or in *Escherichia coli* by adding lactose or an analog thereof (e.g., IPTG). This selective regulation may be a consequence of the fact that *Staphylococcus* does not express the *lac* repressor that binds to the *lacO* while *Escherichia coli* does not express the *xylR* that binds to the *xylO*. In this manner, the construct can be efficiently amplified in *Escherichia coli* and retain the ability to express the desired transcript in *Staphylococcus* once the binding of *xylR* to *xylO* is inhibited by adding an inducer such as xylose or an analog thereof.

Similar multioperator constructs, which incorporate both *lacO* and *xylO* so as to provide regulatable expression in both *Escherichia coli* and Gram-positive organisms, can be made using promoter sequences described herein, including promoters selected from SEQ ID NOs.: 26 - 35.

By following the approaches described in the examples below, with modifications as would be apparent to one of skill in the art, the preparation of a fusion promoter comprising operators other than those described in the examples can be accomplished without undue burden or extensive experimentation.

The present invention also contemplates fusion promoters that have been modified to have altered activity in Gram-positive organisms. Modifications to fusion promoters can be made, for example, by synthesizing the modified DNA sequences, PCR based mutagenesis, site-directed mutagenesis techniques, DNA shuffling or any other techniques for making site specific mutations.

In some embodiments of the present invention, nucleotides in the -10, -35 or -45 Boxes of a promoter are changed to make those sequence elements look more or less like the consensus -10, -35 or -45 Box sequences in the desired Gram-positive organisms. Such modification can have the affect of either increasing or decreasing the activity of the fusion promoter.

It will be appreciated that a regulatable fusion promoter suite can be constructed which provides a wide dynamic range of both basal and maximal transcriptional activities. It will also be appreciated that modifying a regulatable fusion promoter so as to increase or decrease its activity can be used to expand the range of both basal and maximal transcriptional activities of a suite of regulatable fusion promoters. Each modified fusion promoter within this suite can be tightly and finely regulated over its own characteristic range of transcriptional activity. Furthermore, the modified regulatable fusion promoters within the suite can comprise different promoters operably linked to the same operators thus providing for a common means of regulation of the promoters within the suite. In some embodiments, the promoter within the set of modified regulatable fusion promoters within the suite can be linked to at least one operator contained in a promoterless expression vector that is capable of replicating in one or more Gram-positive organisms. Accordingly, a regulatable fusion promoter having an appropriate basal level and/or maximal level of transcriptional activity can be selected from the suite for use in the appropriate application.

In other embodiments, promoters are modified by altering the length of the region between the operator or transcription start site and the -10 Box, altering the length and/or composition of the region between the -10 Box and the -35 Box, and altering the length and/or composition of the AT rich region upstream of the -35 Box (i.e. the -45 Box). It will also be appreciated that modifications which affect promoter activity can include addition of elements such as the CRE region for catabolite repression in Gram-positive bacteria. These additional elements can be further modified so that they are more or less like their corresponding consensus sequence.

In some embodiments, modified fusion promoters can function in one or more Gram-positive organisms. For example, some modified fusion promoters of the present invention function in *Staphylococcus aureus* and *Enterococcus faecalis*. Other modified fusion promoters described herein function in Gram-positive organisms such as *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*,

Enterococcus faecium, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species.

In some embodiments of the present invention, the promoters described herein are used to initiate transcription of an RNA (sense or anti-sense) in bacteria, desirably Gram-positive bacteria. In another embodiment, random polynucleotides (e.g., nucleic acids generated by oligonucleotide synthesis, shearing of bacterial genomic DNA or enzymatic digestion, such as digestion with restriction endonucleases, DNase I or other nucleases) are operably linked to a promoter described herein. These constructs are transferred to bacteria and can be used to transcribe RNA.

In one embodiment of the present invention, a fusion promoter comprising an operator, such as *xylO*, and a promoter, such one selected from SEQ ID NOs.: 36 – 45, is operably linked to a nucleic acid to be transcribed from the promoter, the construct is introduced into bacteria, a solution of xylose or an analog thereof is provided (e.g., a 2.0% xylose solution), and induction of transcription proceeds. Preferred nucleic acids that are operably linked to the fusion promoter are at least 6, at least 10, at least 15, at least 20, at least 100, at least 200, at least 300, at least 400, at least 500, at least 800 or more than 800 nucleotides in length and, in some embodiments, the nucleic acids that are operably linked to the fusion promoter encode a transcript that is complementary to at least 6, at least 10, at least 15, at least 20, at least 100, at least 200, at least 300, at least 400, at least 500, at least 800 or more than 800 nucleotides within the genome of a Gram-positive organism, such as *Staphylococcus* or *Enterococcus*, which code for a gene product such as a protein or RNA. That is, nucleic acids that are operably linked to the fusion promoter can be in the "sense" or "antisense" orientation and can be at least 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83,

84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 125, 150, 175, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800 or more than 800 nucleotides in length.

In another embodiment, the promoters described herein are used in an expression construct to initiate the production of a polypeptide or protein in bacteria, desirably Gram-positive bacteria. In some aspects of this embodiment, the peptide or protein generated corresponds to an expression product encoded by a full-length gene, a portion of a full-length gene, or a random polynucleotide or fragment of DNA generated by oligonucleotide synthesis or enzymatic digestion of genomic DNA (e.g., bacteria, yeast, or mold).

In one embodiment of the present invention, a fusion promoter comprising an operator, such as *xylO*, and a promoter, such one selected from SEQ ID NOs.: 36 – 45, is operably linked to a nucleic acid to be transcribed from the promoter, the construct is introduced into bacteria, a solution of xylose or an analog thereof is provided (e.g., a 2.0% xylose solution), and induction of transcription proceeds. This expression method can be performed in Gram-positive or Gram-negative bacteria and are desirably performed in *Staphylococcus*, *Enterococcus* or other Gram-positive species.

The DNA encoding the desired molecule can be generated by oligonucleotide synthesis or enzymatic cleavage of DNA (e.g., random fragments of bacterial genomic DNA including, but not limited to, *Staphylococcus*) The peptides expressed by using a fusion promoter can be polypeptides, proteins, and fusion proteins. The peptides can be, for example, 3-20 amino acids in length, 20-100 amino acids in length, or 100, 200, 300, 400, 500, 600, 700, 800, or more amino acids in length. That is, the peptides can be at least 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 125, 150, 175, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800 or more than 800 amino acids in length.

Regulatable promoters providing a range of transcriptional levels in both the repressed and induced states, such as the promoters described herein, are desirable in applications where production of defined levels of transcript or protein is necessary. For example, promoters which can be used to tightly and/or finely regulate transcription at a desired level are desirable for use in cell-based assays which employ sensitized cells for screening potential antibiotic compounds (described in Examples 19-25). Specifically, it is desirable to use regulatable fusion promoters, such as those described herein, to modulate the sensitivity of cells to test compounds by altering the amount of a proliferation-inhibiting antisense nucleic acid transcribed by the promoter. Because the results of such cell-based assays are affected by the degree to which the cells are sensitized, the efficacy of these assays can be increased by using promoters having the ability to finely regulate the level of proliferation-inhibiting transcript over a wide range of transcript concentrations.

The present invention contemplates fusion promoters which can be used to achieve the desired level of nucleic acid transcription or protein expression desired for applications such as the cell-based gene and drug discovery methods described herein. The appropriate fusion promoter can be selected by first determining the approximate level of transcript production that is required for the assay. For example, if only a low level of transcription is necessary, a promoter with a low maximum transcription efficiency may be used. If a tightly controllable transcription over a wide range is required, a fusion promoter having nonleaky repression and a high maximum transcription efficiency may be used. The amount of actual transcription obtained from the fusion promoter can then be fine tuned by determining the concentration of inducer which achieves the desired effect. Inducer can be introduced to the cell cultures as a solid or a liquid. The inducer can also be an appropriate growth condition, including but not limited to, the absence of a regulatory molecule, a change in temperature or exposure to radiation. The inducer can comprise from 0% to greater than 25% of the concentration of the growth medium or can be an appropriate growth condition. In some embodiments, the inducer comprises 0%, 0.1%, 0.2%, 0.3%, 0.4%, 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1.0%, 1.5%, 2.0%, 3.0%, 4.0%, 5.0%, 6.0%, 7.0%,

8.0%, 9.0%, 10.0%, 12.0%, 15%, 17%, 20%, 25%, or greater than 25% of the concentration of the growth medium.

The present invention includes vectors containing fusion promoters operably linked to a reporter gene. Such expression vectors can be used to quantitate the transcriptional activity of the fusion promoter contained therein. In some
5 embodiments, a promoter can be inserted into the reporter vector as a cassette so as to form a fusion promoter construct with an operator located on the vector upstream of the reporter gene. In a preferred embodiment, the β -galactosidase encoding genes, *lacL-lacM*, from *Leuconostoc mesenteroides*, (Israelsen et al., *Appl. Environ.*
10 *Microbiol.* 61:2540-2547 (1995); Pedersen et al., *Mol. Gen. Genet.* 244:374-382 (1994), the disclosures of which are incorporated by reference in their entireties), are linked to a fusion promoter described herein. The present invention also contemplates methods for using the *lacL-lacM* genes as a reporter in *Enterococcus faecalis*.

Expression vectors containing fusion promoters linked upstream of a multiple
15 cloning site (MCS) and/or a transcription terminator are also contemplated by the present invention. In a preferred embodiment the transcription terminator is a rho-independent terminator such as *rrnB1t2*. In some embodiments of the present invention, a promoter sequence can be inserted into the expression vector as a cassette so as to form a fusion promoter construct with an operator located on the vector
20 upstream of the MCS. In other embodiments, a promoter linked to an operator can be inserted into the expression vector as a cassette just upstream of the MCS.

Expression vectors containing fusion promoters can be used to produce transcripts, such as antisense RNA, or for the expression of proteins. In some
25 embodiments, nucleic acids corresponding to proliferation-required genes or fragments thereof or random genomic fragments are inserted into the MCS in a sense or antisense orientation. Upon induction of the fusion promoter with an inducer, antisense or sense transcripts are produced. Expression of such transcripts from the fusion promoter can be tightly and/or finely regulated.

In a preferred embodiment of the present invention, reporter and expression
30 vectors contain both an origin of replication that is functional in *Escherichia coli*

and/or other Gram-negative organisms and an origin of replication that function in one or more Gram-positive organisms such as *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium teteni*, *Corynebacterium diptheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species. In some embodiments, an origin of replication is functional in *Staphylococcus aureus* and *Enterococcus faecalis*.

10 In some embodiments, nucleic acids that are operably linked to the fusion promoter encode a molecule (e.g., a sense RNA, anti-sense RNA, aptamer, or peptide) that inhibits proliferation of Gram-positive bacteria including, but not limited to, *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium teteni*, *Corynebacterium diptheriae*, *Enterococcus faecalis*,
15 *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species. Certain embodiments include methods of identifying genes that are essential
20 to bacterial proliferation ("essential genes") whereby the constructs described above are selectively induced in bacteria, such as pathogenic bacteria, at various levels and the impact on cell proliferation is monitored. For example, the fusion promoters of the present invention may be utilized in any of the methods described in United States Patent Application Serial Number 09/492,709, United States Patent Application Serial
25 Number 09/711,164, United States Patent Application Serial Number 09/741,699, and United States Patent Application Serial Number 09/815,242, the disclosures of which are incorporated by reference herein in their entireties.

One such method is practiced by introducing a candidate nucleic acid into a microorganism, wherein the candidate nucleic acid is operably linked to a fusion
30 promoter comprising a *xyl*, *tet*, *trp*, *mal*, *lac* or *lacI* operator and one of the promoter

sequences described herein. The fusion promoter may comprise a promoter that has been modified to alter its transcriptional activity in one or more Gram-positive bacteria including but not limited to *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diptheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species. In some embodiments, the candidate nucleic acid is generated by random oligonucleotide synthesis and in other embodiments, the candidate nucleic acid is generated by cleavage of genomic DNA from an organism (e.g., bacteria, viruses, fungi, mold, parasites, yeast, plants and animals). For example, the candidate nucleic acid may be a random fragment generated by methods such as DNaseI digestion, restriction endonuclease digestion or shearing. In some cases, the candidate nucleic acid will be made so that an antisense RNA, sense RNA, aptamer, peptide, or protein will be expressed upon induction of the fusion promoter. In some cases, the candidate nucleic acid will encode a transcript that corresponds to or is complementary to a gene present in the genome of an organism (e.g., bacteria, viruses, fungi, mold, parasites, yeast, plants and animals). In other cases, the candidate nucleic acid will encode a peptide that corresponds to or interacts with a protein in the bacteria.

In the embodiments in which an aptamer is transcribed upon induction of the fusion promoter, the aptamer may be a nucleic acid which directly binds to the target molecules. Alternatively, the aptamer may encode a peptide that binds to the target molecule. For example, the aptamer may encode a peptide of at least 4, 6, 8, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70 or more than 70 amino acids in length. Aptamers which encode peptides have been described in PCT publication WO 99/50462, the disclosure of which is incorporated herein by reference in its entirety.

For example, RNA aptamers corresponding to regions of Tat and Rev RNA were introduced into or expressed in a cell and were shown to be specific RNA

ligands for proteins that can specifically inhibit their translation. (Good *et al.*, 1997, Gene Therapy 4: 45-54). Similarly, Conrad *et al.* (J. Biol. Chem. 269: 32051, 1994) have shown that certain RNA aptamers can inhibit the β II isoenzyme of PKC. These RNA aptamers were selected from a pool of RNA molecules that contained a 120-
5 nucleotide randomized region.

In general, the aspects of the invention that employ aptamer technology concern the use of the fusion promoters described herein to drive expression of aptamers of varying lengths and compositions. A minimum of approximately 6 nucleotides, preferably 10, and more preferably 14 or 15 nucleotides, may be
10 necessary to effect specific binding to some target molecules. Thus, in some embodiments, fusion promoters of the invention are linked to oligonucleotides encoding aptamers that are 6, 10, 14, or 15 nucleotides in length. The only apparent limitations on the binding specificity of the target/aptamer complex concern sufficient sequence to be distinctive in the binding aptamer and sufficient binding capacity of the
15 target molecule to obtain the necessary interaction. Aptamers of binding regions containing sequences shorter than 10, e.g., 6-mers, are feasible if the appropriate interaction can be obtained in the context of the environment in which the target is placed. Thus, if there are few interferences by other materials, less specificity and less strength of binding may be required.

Several methods for selecting aptamer oligonucleotides directed against a specific protein are known in the art and these approaches can be employed to design constructs having the fusion promoters linked to an oligonucleotide that encodes an aptamer. Examples of these methods include those methods described by Mills *et al.* Proc. Natl. Acad. Sci. USA 58:224 (1967), Green *et al.* Nature 347:406 (1990), and
25 Tuerk and Gold Science 249:505 (1990), (all references are expressly incorporated by reference in their entireties), which show the use of *in vitro* evolution techniques to select RNA molecules with desired properties. Ellington and Szostak Nature 346:818 (1990), and Bock *et al.* Nature 355:564 (1992) (all references are expressly incorporated by reference in their entireties) also describe techniques useful for the
30 selection of DNA oligonucleotides. More teachings on the design and selection of

apamers can be found in U.S. Pat. Nos. 5,756,291; 5,874,567; 5,874,281; 5,998,596; and 5,965,352 (all references are expressly incorporated by reference in their entireties). The field has also been recently reviewed by Burke and Berzal-Herranz FASEB J 7:106 (1993), herein expressly incorporated by reference in its entirety.

5 Once the construct comprising the candidate nucleic acid operably linked to a promoter of the invention is made, it is transferred to host bacteria. Next, the proliferation of microorganisms transcribing a higher level of the candidate nucleic acid is compared to the proliferation of microorganisms that transcribe a lower level of the candidate nucleic acid or that do not transcribe the candidate nucleic acid. These
10 varying levels of transcription can be achieved by varying the amounts of inducer (e.g., xylose, tetracycline, maltose, the absence of tryptophan, temperature, exposure to ultraviolet radiation, or analogs thereof). The level of maximal induction that can be achieved will depend on the fusion promoter selected. A difference in cell proliferation in the sample transcribing a higher level of the candidate nucleic acid
15 compared to the sample expressing a lower level of the candidate nucleic acid indicates that the candidate nucleic acid encodes a nucleic acid that is complementary to an essential gene or encodes a peptide or protein that reduces proliferation of the microorganism.

 In some embodiments, the RNA, aptamer, peptide or protein reduces
20 proliferation of the microorganisms by interacting with another molecule required for proliferation. In this case, many techniques in molecular biology can be employed to identify the cell proliferation molecule that interacts with the aptamer, peptide, or protein and the gene encoding this molecule. Conventional one and two hybrid systems, for example, can be readily adapted to identify molecules that bind to an
25 aptamer, peptide, or protein described above, for example. Such approaches include:

- (1) the two-hybrid systems (Field & Song, *Nature* 340:245-246 (1989); Chien *et al.*, *Proc. Natl. Acad. Sci. USA* 88:9578-9582 (1991); and Young KH, *Biol. Reprod.* 58:302-311 (1998), all of which are
30 expressly incorporated by reference in their entirety);

- (2) the reverse two-hybrid system (Leanna & Hannink, *Nucl. Acid Res.* 24:3341-3347 (1996), herein incorporated by reference in their entirety);
- (3) the repressed transactivator system (Sadowski *et al.*, U.S. Pat. No. 5,885,779), herein incorporated by reference in their entirety);
- (4) phage display systems (Lowman HB, *Annu. Rev. Biophys. Biomol. Struct.* 26:401-424 (1997), herein incorporated by reference in their entirety); and
- (5) GST/HIS pull down assays, mutant operators (Granger *et al.*, WO 98/01879) and the like (*See also* Mathis G., *Clin. Chem.* 41:139-147 (1995); Lam K.S. *Anticancer Drug Res.*, 12:145-167 (1997); and Phizicky *et al.*, *Microbiol. Rev.* 59:94-123 (1995), all of which are expressly incorporated by reference in their entirety).

In one embodiment, the present invention utilizes an antisense-based method to identify proliferation-required sequences. Generally, a library of nucleic acids from a given source are subcloned or otherwise inserted immediately downstream of an inducible fusion promoter on an appropriate vector, such as one of the expression vectors described herein, thus forming an expression library. It is generally preferred that expression is directed by a regulatable fusion promoter such that expression level can be adjusted by addition of variable concentrations of an inducer molecule or of an inhibitor molecule to the medium. Temperature activated fusion promoters, such as promoters regulated by temperature sensitive repressors, such as the lambda C₁₈₅₇ repressor, are also envisioned. Although the insert nucleic acids may be derived from the chromosome of the cell or microorganism into which the expression vector is to be introduced, because the insert is not in its natural chromosomal location, the insert nucleic acid is an exogenous nucleic acid for the purposes of the discussion herein.

Once generated, the expression library containing the nucleic acids is introduced into a population of cells (such as the organism from which the nucleic acid sequences were obtained or an organism other than the organism from which the sequences were

obtained) to search for genes that are required for bacterial proliferation. Because the library molecules are foreign, in context, to the population of cells, the expression vectors and the nucleic acid segments contained therein are considered exogenous nucleic acid.

5 Expression of the nucleic acid fragments in the test population of cells containing the expression library is then activated. Activation of the expression vectors consists of subjecting the cells containing the vectors to conditions that result in the expression of the nucleic acids carried by the expression library. The test population of cells is then assayed to determine the effect of expressing the nucleic acid fragments on
10 the test population of cells. Those expression vectors that negatively impact the growth of the cells upon induction of expression of the random sequences contained therein are identified, isolated, and purified for further study.

 A variety of assays are contemplated to identify nucleic acid sequences that negatively impact growth upon expression. In one embodiment, growth in cultures
15 expressing nucleic acid sequences and growth in cultures not expressing these sequences is compared. Growth measurements are assayed by examining the extent of growth by measuring optical densities. Alternatively, enzymatic assays can be used to measure bacterial growth rates to identify nucleic acid sequences of interest. Colony size, colony morphology, and cell morphology are additional factors used to evaluate growth
20 of the host cells. Those cultures that fail to grow or grow at a reduced rate under expression conditions are identified as containing an expression vector encoding a nucleic acid fragment that negatively affects a proliferation-required gene.

 In some embodiments, genomic fragments of an organism are operably linked to an inducible fusion promoter in a vector and assayed for growth inhibition activity.
25 Examples below describe the examination of libraries of genomic fragments cloned into vectors comprising inducible fusion promoters. Vectors containing fusion promoters, such as those described herein, can be used. Upon transformation of these vectors into a host organism and induction, the vectors produce an RNA molecule corresponding to the subcloned genomic fragments. In those instances where a genomic fragment is in an
30 antisense orientation with respect to the fusion promoter, the antisense transcript

produced can be complementary to at least a portion of a sense messenger RNA (mRNA) encoding a proliferation-required gene product such that the antisense transcript interacts with at least a portion of the proliferation-required sense mRNA produced by the host organism thereby decreasing the translation efficiency or the level of the proliferation-required sense mRNA and thus decreasing production of the proliferation-required protein encoded by the sense mRNA molecule. In cases where the sense mRNA encodes a protein required for proliferation, the host organism containing a vector from which transcription from the fusion promoter has been induced fails to grow or grows at a substantially reduced rate. Additionally, in cases where the transcript produced is complementary to at least a portion of a non-translated RNA and where that non-translated RNA is required for proliferation, host cells containing a vector from which transcription from the fusion promoter has been induced also fail to grow or grow at a substantially reduced rate.

In cases where the candidate nucleic acid encodes an antisense nucleic acid comprising a nucleotide sequence that is complementary to a gene required for proliferation or a portion thereof, the gene sequence of the coding fragment of the proliferation-required gene or a portion thereof can be determined by direct sequencing.

Determination of the gene(s) corresponding to the nucleotide sequence can be achieved by comparing the obtained sequence data with databases containing known protein and nucleotide sequences from various microorganisms. Thus, initial gene identification can be made on the basis of significant sequence similarity or identity to either characterized or predicted from *Staphylococcus aureus* and *Enterococcus faecalis* genes or their encoded proteins and/or homologues in other species.

The number of nucleotide and protein sequences available in database systems has been growing exponentially for years. For example, the complete nucleotide sequences of *Caenorhabditis elegans* and several bacterial genomes, including *E. coli*, *Aeropyrum pernix*, *Aquifex aeolicus*, *Archaeoglobus fulgidus*, *Bacillus subtilis*, *Borrelia burgdorferi*, *Chlamydia pneumoniae*, *Chlamydia trachomatis*, *Clostridium tetani*, *Corynebacterium diphtheria*, *Deinococcus radiodurans*, *Haemophilus*

influenzae, *Helicobacter pylori* 26695, *Helicobacter pylori* J99, *Methanobacterium thermoautotrophicum*, *Methanococcus jannaschii*, *Mycobacterium tuberculosis*, *Mycoplasma genitalium*, *Mycoplasma pneumoniae*, *Pseudomonas aeruginosa*, *Pyrococcus abyssi*, *Pyrococcus horikoshii*, *Rickettsia prowazekii*, *Synechocystis* PCC6803, *Thermotoga maritima*, *Treponema pallidum*, *Bordetella pertussis*, *Campylobacter jejuni*, *Clostridium acetobutylicum*, *Mycobacterium tuberculosis* CSU#93, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Pseudomonas aeruginosa*, *Pyrobaculum aerophilum*, *Pyrococcus furiosus*, *Rhodobacter capsulatus*, *Salmonella typhimurium*, *Streptococcus mutans*, *Streptococcus pyogenes*, *Ureaplasma urealyticum* and *Vibrio cholera* are available. This nucleotide sequence information is stored in a number of databanks, such as GenBank, the National Center for Biotechnology Information (NCBI), the Genome Sequencing Center (<http://genome.wustl.edu/gsc/salmonella.shtml>), and the Sanger Centre (http://www.sanger.ac.uk/projects/S_typhi) which are publicly available for searching.

A variety of computer programs are available to assist in the analysis of the sequences stored within these databases. FASTA, (W. R. Pearson (1990) "Rapid and Sensitive Sequence Comparison with FASTP and FASTA" Methods in Enzymology 183:63- 98), Sequence Retrieval System (SRS), (Etzold & Argos, SRS an indexing and retrieval tool for flat file data libraries. Comput. Appl. Biosci. 9:49-57, 1993) are two examples of computer programs that can be used to analyze sequences of interest. In one embodiment of the present invention, the BLAST family of computer programs, which includes BLASTN version 2.0 with the default parameters, or BLASTX version 2.0 with the default parameters, is used to analyze nucleotide sequences.

BLAST, an acronym for "Basic Local Alignment Search Tool," is a family of programs for database similarity searching. The BLAST family of programs includes: BLASTN, a nucleotide sequence database searching program, BLASTX, a protein database searching program where the input is a nucleic acid sequence; and BLASTP, a protein database searching program. BLAST programs embody a fast algorithm for sequence matching, rigorous statistical methods for judging the significance of matches,

and various options for tailoring the program for special situations. Assistance in using the program can be obtained by e-mail at blast@ncbi.nlm.nih.gov. tBLASTX can be used to translate a nucleotide sequence in all three potential reading frames into an amino acid sequence.

5 Bacterial genes are often transcribed in polycistronic groups. These groups comprise operons, which are a collection of genes and intergenic sequences under common regulation. The genes of an operon are transcribed on the same mRNA and are often related functionally. Given the nature of the screening protocol, it is possible that the identified exogenous nucleic acid corresponds to a gene or portion thereof with or
10 without adjacent noncoding sequences, an intragenic sequence (i.e. a sequence within a gene), an intergenic sequence (i.e. a sequence between genes), a nucleotide sequence spanning at least a portion of two or more genes, a 5' noncoding region or a 3' noncoding region located upstream or downstream from the actual nucleotide sequence that is required for bacterial proliferation. Accordingly, it is often desirable to determine
15 which gene(s) that is encoded within the operon is individually required for proliferation.

 In one embodiment of the present invention, an operon is identified and then dissected to determine which gene or genes are required for proliferation. Operons can be identified by a variety of means known to those in the art. For example, the
20 RegulonDB DataBase described by Huerta et al. (*Nucl. Acids Res.* 26:55-59, 1998), which may also be found on the website http://www.cifn.unam.mx/Computational_Biology/regulondb/, the disclosures of which are incorporated herein by reference in their entireties, provides information about operons in *Escherichia coli*. The Subtilist database
25 (<http://bioweb.pasteur.fr/GenoList/SubtiList>), (Moszer, I., Glaser, P. and Danchin, A. (1995) *Microbiology* 141: 261-268 and Moszer, I (1998) *FEBS Letters* 430: 28-36, the disclosures of which are incorporated herein in their entireties), may also be used to predict operons. This database lists genes from the fully sequenced, Gram-positive bacteria, *Bacillus subtilis*, together with predicted promoters and terminator sites.
30 This information can be used in conjunction with the *Staphylococcus aureus* genomic

sequence data to predict operons and thus produce a list of the genes affected by the antisense nucleic acids of the present invention. The TIGR microbial database has an incomplete version of the *Enterococcus faecalis* genome http://www.tigr.org/cgi-bin/BlastSearch/blast.cgi?organism=e_faecalis. One can take a nucleotide sequence and BLAST it for homologs.

5 A number of techniques that are well known in the art can be used to dissect the operon. Analysis of RNA transcripts by Northern blot or primer extension techniques are commonly used to analyze operon transcripts. In one aspect of this embodiment, gene disruption by homologous recombination is used to individually inactivate the genes of an operon that is thought to contain a gene required for proliferation.

10 Several gene disruption techniques have been described for the replacement of a functional gene with a mutated, non-functional (null) allele. These techniques generally involve the use of homologous recombination. One technique using homologous recombination in *Staphylococcus aureus* is described in Xia et al. 1999, Plasmid 42: 144-149, the disclosure of which is incorporated herein by reference in its entirety. This technique uses crossover PCR to create a null allele with an in-frame deletion of the coding region of a target gene. The null allele is constructed in such a way that nucleotide sequences adjacent to the wild type gene are retained. These homologous sequences surrounding the deletion null allele provide targets for homologous recombination so that the wild type gene on the *Staphylococcus aureus* chromosome can be replaced by the constructed null allele. *Enterococcus faecalis* genes can be disrupted by recombining in a non-replicating plasmid that contains an internal fragment to that gene (Leboeuf, C., L. Leblanc, Y. Auffray and A. Hartke. 2000. J. Bacteriol. 182:5799-5806, the disclosure of which is incorporated herein by reference in its entirety).

25 The crossover PCR amplification product is subcloned into a suitable vector having a selectable marker, such as a drug resistance marker. In some embodiments the vector may have an origin of replication which is functional in *Escherichia coli* or another organism distinct from the organism in which homologous recombination is to occur, allowing the plasmid to be grown in *Escherichia coli* or the organism other than

30

that in which homologous recombination is to occur, but may lack an origin of replication functional in *Staphylococcus aureus* or *Enterococcus faecalis* such that selection of the selectable marker requires integration of the vector into the homologous region of the *Staphylococcus aureus* or *Enterococcus faecalis* chromosome. Usually a single crossover event is responsible for this integration event such that the *Staphylococcus aureus* or *Enterococcus faecalis* chromosome now contains a tandem duplication of the target gene consisting of one wild type allele and one deletion null allele separated by vector sequence. Subsequent resolution of the duplication results in both removal of the vector sequence and either restoration of the wild type gene or replacement by the in-frame deletion. The latter outcome will not occur if the gene should prove essential. A more detailed description of this method is provided below. It will be appreciated that this method may be practiced with any genes identified as required for proliferation in any of the organisms described herein.

Bacteria that harbor the expression vectors described herein can be induced to transcribe a desired level of RNA or to express a desired level of peptide, which affects a gene that is essential for the viability and/or growth of the cell and renders the cells more sensitive to compounds that reduce cellular proliferation. This technique is referred to as Essential Gene Interference (EGI). In one embodiment, for example, an RNA, such as an antisense RNA, or peptide generated from the constructs described above interacts with an essential gene of a bacteria, RNA, or product thereof and when the level of transcript or peptide is increased in the cell, viability and/or cell growth is compromised. Thus, the genes that are affected by the transcript or peptide either directly or indirectly (e.g., by interacting with the RNA or protein that correspond to the gene) causing a reduction in viability and/or cell growth are identified as essential genes. These essential genes are then used as drug discovery targets for high throughput drug screening.

The promoters described herein can also be used to replace endogenous promoters so as to create cells having specific regulatable genes. By this approach, an essential gene is identified (e.g., using the techniques described herein) and the endogenous promoter is replaced with a regulatable promoter of the invention (e.g.,

Xyl-T5, Xyl-T5-DD, Xyl-CP25, Xyl-P32, Xyl-P59, Xyl-P59*, Xyl-P1P2, Xyl-P1P2*, Xyl, PL, and XylPL10) using ET cloning technology. This technique is referred to as Rapid Replacement of Genomic Promoters (promoter replacement). This method employs the Polymerase Chain Reaction (PCR) to replace the promoter of the target gene without interrupting the endogenous Shine-Dalgarno sequence. As little as 60bp homology to the target promoter can be used. The correct integration of the cassette is confirmed by colony PCR and correct integrants are tested for growth defects when an inducer is titrated away. The inability to grow in the absence of inducer confirms the essentiality of the target gene. Additionally, the bacterial strains created using this approach can be used in high throughput drug screening methodologies.

A similar technique to promoter replacement can be used to insert an operator into a genomic promoter region so as to make the genomic promoter regulatable. This technique is referred to as operator insertion. As little as 60bp homology to the target promoter region can be used. The correct integration of the cassette is confirmed by colony PCR and correct integrants are tested for growth defects when an inducer is titrated away. The inability to grow in the absence of inducer confirms the essentiality of the target gene. Additionally, the bacterial strains created using this approach can be used in high throughput drug screening methodologies.

In some embodiments, promoter replacement and operator insertion methods are conducted in cells which have an increased frequency of homologous recombination. For example, the organism may lack or have a reduced level or activity of one or more exonucleases which would ordinarily degrade the DNA to be inserted into the chromosome, thereby effectively increasing the stability of the linear DNA in the organism.

In still further embodiments, the strains may both lack or have reduced levels of exonucleases and constitutively or conditionally express proteins involved in mediating homologous recombination. For example, Gram-positive bacterial strains in which the activity of an exonuclease which degrades linear nucleic acids, such as exonuclease V of the RecBCD recombination pathway or an exonuclease with an analogous function, has been reduced or eliminated, may be used. Such strains

include those having mutations in the *recB*, *recC*, or *recD* genes or genes that express proteins having analogous functions, that enhance the frequency of homologous recombination. In some embodiments, the strains have mutations in more than one of the *recB*, *recC*, or *recD* genes or genes that express proteins having analogous functions, that enhance the frequency of homologous recombination. For example the strains may have mutations in both the *recB* and *recC* genes or genes that express proteins having analogous functions.

It will be appreciated that Gram-positive organisms having reduced or eliminated levels of one or more proteins analogous in function to *RecB*, *RecC*, or *RecD* may be used with the methods described herein. In some embodiments, Gram-positive organisms having reduced or eliminated levels of one or more proteins analogous in function to *RecB*, *RecC* or *RecD* may be selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species. In some embodiments, it may be desirable to use *Escherichia coli* having reduced or eliminated levels of one or more of *RecB*, *RecC* or *RecD* proteins as a surrogate host in the methods described herein.

The methods may also be performed in Gram-positive bacterial strains in which the activity of a the *RecET* recombinase system of the *Rac* prophage, or a recombinase system having an analogous function, has been activated. Such strains are analogous to *Escherichia coli* strains which carry an *sbcA* mutation. The *RecE* gene of the *Rac* prophage encodes *ExoVIII* a 5'-3' exonuclease, while the *RecT* gene of the *Rac* prophage encodes a single stranded DNA binding protein which facilitates renaturation and D-loop formation. Thus, the gene products of the *RecE* and *RecT* genes or proteins with analogous functions facilitate homologous recombination. The *RecE* and *RecT* genes are on the same operon and are not normally expressed.

Similarly, genes having a function analogous to the RecE and RecT may not be expressed. However, strains having a mutation analogous to *sbcA* would activate the expression of RecE and RecT or genes having analogous function. In some embodiments, the methods may be performed in strains which carry mutations in the *recB* and *recC* genes or genes that express proteins having analogous functions, as well as the mutation in the gene analogous in function to *sbcA*.

In some embodiments, the methods may be performed in Gram-positive bacterial strains in which recombination via the RecF pathway or a pathway having an analogous function has been enhanced. One example of such strains are those having a mutation analogous in function to the *Escherichia coli sbcB* mutation.

It will be appreciated that the *recE* and *recT* gene products, or proteins with analogous functions may be constitutively or conditionally expressed in Gram-positive organisms. In some embodiments, these proteins may be conditionally or constitutively expressed in Gram-positive organisms such as *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species. In some embodiments, Gram-positive strains may be genetically engineered to express the RecE and *recT* genes of the λ prophage. For example, plasmids encoding these gene products may be introduced into the organism. If desired, the coding sequences encoding these gene products may be optimized to reflect the codon preferences of the organism in which they are to be expressed. In some embodiments, it may be desirable to use *Escherichia coli* constitutively or conditionally expressing the RecE and RecT gene products as a surrogate host in the methods described herein.

In further embodiments, the methods may be conducted in cells which utilize the Red system of bacteriophage lambda (λ) or analogous systems from other phages to enhance the frequency of homologous recombination. The Red system contains

three genes, γ , β and *exo* whose products are the Gam, Bet and Exo proteins (see Ellis et al. PNAS 98:6742-6746, 2001, the disclosure of which is incorporated herein by reference in its entirety). The Gam protein inhibits the RecBCD exonuclease V, thus permitting Bet and Exo to gain access to the ends of the DNA to be integrated and facilitating homologous recombination. The Bet protein is a single stranded DNA binding protein. The Exo protein is a double-stranded DNA dependent 5'-3' exonuclease. Thus, expression of the λ red proteins or proteins having analogous functions facilitates homologous recombination.

It will be appreciated that the λ Bet, Gam and Exo proteins, or proteins with analogous functions may be constitutively or conditionally expressed in Gram-positive organisms. In some embodiments, these proteins may be conditionally or constitutively expressed in Gram-positive organisms such as *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species. For example, plasmids encoding these gene products may be introduced into the organism. If desired, the coding sequences encoding these gene products may be optimized to reflect the codon preferences of the organism in which they are to be expressed. In some embodiments, it may be desirable to use *Escherichia coli* constitutively or conditionally expressing the λ Bet, Gam and Exo gene products as a surrogate host in the methods described herein.

In some embodiments, the cells may have an increased frequency of homologous recombination as a result of more than one of the aforementioned characteristics. In some embodiments, the enhanced frequency of recombination may be a conditional characteristic of the cells which depends on the culture conditions in which the cells are grown. For example, in some embodiments, expression of the λ Red Gam, Exo, and Bet proteins or *recE* and *recT* proteins may be constitutive or

regulated. Thus, the cells may have an increased frequency of homologous recombination as a result of any combination of the aforementioned characteristics. For example, in some embodiments, the cell may carry the sbcA and recBC mutations or analogous mutations.

5 In some embodiments, a linear double stranded DNA containing the regulatable fusion promoter, operator or other regulatory element to be inserted into the chromosome of the prokaryotic organism is introduced into an organism constitutively or conditionally expressing the recE and recT or the λ Bet, Gam and Exo proteins or proteins with analogous functions as described above. In some
10 embodiments, the double stranded DNA may be introduced into an organism having the recBC and sbcA mutations or analogous mutations.

 In other embodiments, a single stranded DNA containing the regulatable fusion promoter, operator or other regulatory element to be inserted into the chromosome of the prokaryotic organism is introduced into an organism constitutively
15 or conditionally expressing the λ Bet protein or a protein with an analogous function. In some embodiments the single stranded DNA is introduced into an organism constitutively or conditionally expressing both the λ Bet and Gam proteins or proteins with analogous functions. In further embodiments, the single stranded DNA is introduced into an organism constitutively or conditionally expressing the λ Bet, Gam
20 and Exo proteins or proteins with analogous functions.

 It will be appreciated that any method of enhancing the frequency of homologous recombination may be used in conjunction with the present invention and that the methods of the present invention are not limited to use in cells having the aforementioned characteristics.

25 The 5' and 3' regions of the linear nucleic acid to be integrated via homologous recombination may have any length of homology to the target sequence in the chromosome of the organism which permits homologous recombination to occur. In some embodiments, the nucleic acid to be integrated via homologous recombination is generated by PCR. In such embodiments, it is desirable for the PCR
30 amplicon to be relatively short. Accordingly, in some embodiments, the 5' and 3'

regions of the amplicon which have homology to the target sequence in the chromosome of the organism, may each be more than 120 nucleotides in length, at least 120 nucleotides in length, at least 100 nucleotides in length, at least 90 nucleotides in length, at least 80 nucleotides in length, at least 70 nucleotides in length, at least 60 nucleotides in length, at least 50 nucleotides in length, at least 40 nucleotides in length, at least 30 nucleotides in length or less than 30 nucleotides in length.

As discussed above, the linear double stranded DNA or linear single stranded DNA may be introduced into a suitable strain having an enhanced frequency of recombination.

In one embodiment of the present invention, genes required for proliferation are identified by methods in which the native promoter of a gene is replaced with a regulatable fusion promoter using homologous recombination. In such methods, a linear promoter replacement construct which comprises a 5' portion homologous to a first portion of the native promoter, a 3' portion homologous to a second portion of the native promoter and a regulatable fusion promoter disposed between the 5' portion and the 3' portion is introduced into a cell containing the target gene. After replacement of the native promoter in the target gene with the regulatable fusion promoter via homologous recombination, the extent of proliferation of cells in which the regulatable fusion promoter has replaced the native promoter under conditions in which the regulatable fusion promoter is active at a first level is compared to the extent of proliferation of the cells under conditions in which the regulatable fusion promoter is active at a second level which is lower than the first level. If the candidate gene is required for proliferation, the extent of proliferation of the cells grown under conditions where the fusion promoter is active at the higher level will be greater than the extent of proliferation of cells grown under conditions where the fusion promoter is active at the lower level.

The regulatable fusion promoter in the promoter replacement construct may be any of the regulatable fusion promoters described herein. For example, in some embodiments, the regulatable fusion promoter may be selected from the group

consisting of Xyl-T5, Xyl-T5-DD, Xyl-CP25, Xyl-P32, Xyl-P59, Xyl-P59*, Xyl-P1P2, Xyl-P1P2*, Xyl-PL, and Xyl-PL10.

In some embodiments, the linear promoter replacement construct may further comprise a fusion promoter operably linked to a nucleic acid which encodes an identifiable marker or a selectable marker. The promoter which is operably linked to a nucleic acid encoding an identifiable marker or a selectable marker and the nucleic acid which encodes the identifiable marker or selectable marker may be 5' of the regulatable fusion promoter. The identifiable marker may be any protein whose expression is readily detectable. For example, the identifiable marker may be β -galactosidase, or any of the markers familiar to those skilled in the art which allow identification of cells expressing the marker. Likewise, the selectable marker may be any protein which allows selection of cells which express the protein. For example, the selectable marker may confer resistance to an antibiotic or permit growth under a particular set of culture conditions. A variety of selectable markers are familiar to those skilled in the art, including genes which confer resistance to antibiotics such as chloramphenicol, ampicillin, kanamycin and tetracycline.

In some embodiments, a transcriptional terminator, such as the *rrnB* terminator may be positioned between the 3' end of the nucleic acid encoding the identifiable marker or selectable marker and the 5' end of the regulatable fusion promoter.

The promoter replacement method may be performed in any desired organism. For example, in some embodiments, the method may be practiced in Gram-positive bacteria. In certain embodiments, the cell to be used with any of the methods described herein is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species.

In some embodiments, the promoter replacement method is conducted in cells which have an increased frequency of homologous recombination. For example, in some embodiments, the method is conducted in cells which carry the *sbcA* and *recBC* mutations, or in cells having mutations which confer similar phenotypes. In other
5 embodiments, the cells may express one or more proteins in the λ Red recombination system. For example, in some embodiments the cells may constitutively or conditionally express the λ Gam, Bet, and Exo proteins or other proteins having similar functions. In other embodiments, the cells may constitutively or conditionally express the λ Bet protein or a protein having a similar function.

10 In some embodiments, a double stranded promoter replacement construct is introduced into a cell comprising the *recBCD* and *sbcA* mutations, *recBC* and *sbcA* mutations or mutations with analogous consequences. The double stranded promoter replacement construct may or may not comprise a gene encoding a selectable or identifiable marker, but in preferred embodiments the double stranded promoter
15 replacement construct includes a gene encoding a selectable or identifiable marker. In other embodiments, a double stranded promoter replacement construct is introduced into a cell constitutively or conditionally expressing the *recE* and *recT* proteins or proteins with analogous functions. In still other embodiments, a double stranded promoter replacement construct is introduced into a cell constitutively or conditionally
20 expressing the λ Gam, Bet, and Exo proteins or other proteins having similar functions. In such embodiments, cells in which the promoter replacement construct has integrated are identified by obtaining colonies which express the identifiable marker or by growing the cells in appropriate selective media. If desired, homologous integration may be confirmed by performing a Southern blot or amplification reaction
25 using a probe or primers which will yield a known result if homologous recombination has occurred.

In other embodiments, a promoter replacement construct may be introduced into the cell as a single stranded DNA. Preferably, the single stranded promoter replacement construct does not comprise a gene which encodes an identifiable or
30 selectable marker, but in some embodiments a gene which encodes an identifiable or

selectable marker may be included in the construct. In some embodiments, the single stranded nucleic acid may be generated by *in vitro* synthesis. In other embodiments, a circular single stranded DNA, such as M13, comprising the promoter replacement construct may be fragmented by sonication and the fragments may be introduced into
5 the cell. In still other embodiments, a single stranded promoter replacement construct is introduced into a cell constitutively or conditionally expressing the λ Bet protein, the Bet and Gam proteins, the Bet, Gam and Exo proteins or other proteins having similar functions. Cells in which integration has occurred at the desired location are identified by performing colony hybridization, amplification reactions, or other
10 methods familiar to those skilled in the art.

As discussed above, the 5' and 3' portions of the linear promoter replacement construct which have homology to the promoter region of the target gene may have any length which permits homologous recombination to occur. Accordingly, in some embodiments, the 5' and 3' regions of the amplicon which have homology to the
15 promoter region of the target gene in the chromosome of the organism, may each be more than 120 nucleotides in length, at least 120 nucleotides in length, at least 100 nucleotides in length, at least 90 nucleotides in length, at least 80 nucleotides in length, at least 70 nucleotides in length, at least 60 nucleotides in length, at least 50 nucleotides in length, at least 40 nucleotides in length, at least 30 nucleotides in length
20 or less than 30 nucleotides in length.

In some embodiments, the promoter replacement construct is introduced into the cell as a linear nucleic acid. In particular, in some embodiments, the promoter replacement construct is generated by an conducting amplification reaction, such as a PCR reaction, to generate a linear amplicon. In some aspects of this embodiment, the
25 PCR reaction is performed on a linearized plasmid. In a preferred aspect of this embodiment, the amplicon contains about 60 bp of homology to the target gene at both the 5' and 3' ends of the construct.

Thus, one embodiment of the invention relates to a method for identifying a gene which is required for proliferation of a cell comprising replacing the native
30 promoter of a gene with a regulatable fusion promoter and identifying cells in which

the extent of proliferation of the cell when the fusion promoter is active at a first level is substantially different than the extent of proliferation of the cell when the fusion promoter is active at a second level, the first level being greater than the second level. In some embodiments, the cell has an enhanced frequency of homologous recombination. In some embodiments, the fusion promoter is an inducible promoter. In other embodiments, the level of transcription from the promoter varies with the level of inducer present in the medium in which the cell is grown. In certain embodiments, the step of replacing the native promoter comprises introducing a linear nucleic acid comprising a 5' portion homologous to a first portion of the native promoter, a 3' portion homologous to a second portion of the native promoter and the regulatable promoter disposed between the 5' portion and the 3' portion into a cell such that homologous recombination occurs between the 5' portion and the first portion of the native promoter and between the 3' portion and the second portion of the native promoter. The nucleic acid may, in some embodiments, further comprise a gene encoding a selectable marker. The regulatable fusion promoter may be any of the fusion promoters described herein. In some embodiments, the 5' portion comprises at least 60 nucleotides homologous to the first portion of the native promoter and the 3' portion comprises at least 60 nucleotides homologous to the second portion of the native promoter. Certain embodiments include those in which the nucleic acid is linear. In some embodiments, the step of identifying cells comprises identifying cells which proliferate to a greater extent in the presence of inducer than in the absence of inducer. In other embodiments, the Shine Delgarno sequence of said native promoter is not replaced.

FIG. 3A illustrates one embodiment of the present promoter replacement method in which a linear double stranded promoter replacement construct comprising a nucleic acid encoding a selectable marker, a transcriptional terminator, and a regulatable fusion promoter is inserted into the chromosome of a cell expressing the RecE and RecT gene products of the Rac prophage or gene products having analogous functions. FIG. 3B illustrates one embodiment of the present promoter replacement method in which a linear single stranded promoter replacement construct comprising a

5' portion homologous to a target sequence, a 3' portion homologous to a target sequence and a regulatable fusion promoter is inserted into the chromosome of a cell expressing the λ Beta and Gam proteins.

As discussed below, the promoter replacement method may also be used to obtain cells in which the level or activity of a gene product required for proliferation is reduced. Such cells may be used in cell-based assays to identify compounds which inhibit proliferation.

In another embodiment of the present invention, genes required for proliferation are identified by methods in which an operator is integrated into the promoter of a target gene or into the target gene itself via homologous recombination. The operator may be any nucleic acid which allows the activity or level of the gene product encoded by the target gene to be modulated. For example, in some embodiments, the operator may be recognized by a repressor. In such methods, a linear operator insertion construct which comprises a 5' portion homologous to a first portion of the target promoter or gene, a 3' portion homologous to a second portion of the target promoter or gene and an operator disposed between the 5' portion and the 3' portion is introduced into a cell containing the target gene. After integration of the operator, expression of the target gene is under the control of the operator. For example, in some embodiments, the operator may be recognized by a transcriptional repressor. Numerous operators recognized by transcriptional repressors, such as the operators described herein, including the *lac* operator, the *tet* operator, the *xyl* operator, *λ cl* operator, the *mal* operator, or the *trp* operator, may be used.

After insertion of the operator in the target promoter or gene via homologous recombination, the extent of proliferation of cells in which the operator has been inserted under conditions in which the promoter is active at a first level is compared to the extent of proliferation of the cells under conditions in which the promoter is active at a second level which is lower than the first level. If the candidate gene is required for proliferation, the extent of proliferation of the cells grown under conditions where the promoter is active at the higher level will be greater than the extent of proliferation of cells grown under conditions where the promoter is active at the lower level.

The operator may be integrated at any location in which the activity or level of the gene product encoded by the target gene is modulated by the operator. For example, the operator may be integrated at any location in which expression of the target gene will be regulated by the effector which acts on the operator. For example, if the operator is integrated into a bacterial cell, the operator may be placed upstream of the -35 box, between the -35 box and -10 box, or downstream of the -10 box (e.g. overlapping the transcription start site). In some embodiments, the operator is integrated so as to maintain the appropriate relationships on the DNA helix between the transcription factors which direct expression from the promoter which is placed under the control of the operator.

As discussed above, preferably the cells into which the operator is to be integrated have an enhanced frequency of homologous recombination. In some embodiments, the operator may be introduced into the cell as linear double stranded DNA. Such DNA may be generated by an amplification reaction, such as a PCR reaction. In some aspects of this embodiment, the PCR reaction is performed on a linearized plasmid. In such embodiments, the cells may constitutively or conditionally express the λ Red recombinase system, comprising the Gam, Exo, and Bet proteins or an analogous system from another phage. Alternatively, the cells may comprise the recBCD and sbcA mutations, recBC and sbcA mutations or analogous mutations. In some embodiments, the cells constitutively or conditionally express the recE and recT proteins or proteins with analogous functions. In some embodiments, the cells comprise the recBCD and sbcA mutations, recBC and sbcA mutations, or analogous mutations, and constitutively or conditionally express the recE and recT proteins, or analogous proteins.

In some embodiments, the operator may be introduced into the cells as linear single stranded DNA. Such DNA may be synthesized *in vitro* or generated by sonication of a circular single stranded nucleic acid such as an M13 construct comprising the operator insertion construct. In some embodiments, the cells may constitutively or conditionally express the Bet protein of the λ Red recombinase system or an analogous protein from another phage. In still further embodiments, the

expression of the proteins in the λ Red recombinase system, the Bet protein from the λ Red recombinase system, or analogous proteins from other phage may be conditional.

In some embodiments, the cells may be any of the cells described above which have an increased frequency of homologous recombination or any other cells having
5 an increased frequency of homologous recombination which are familiar to those of skill in the art.

In some embodiments, the 5' and 3' portions of the linear operator insertion construct which have homology to the promoter region of the target gene or the target gene may have any length which permits homologous recombination to occur.
10 Accordingly, in some embodiments, the 5' and 3' regions of the construct which have homology to the promoter region of the target gene or the target gene, may each be more than 120 nucleotides in length, at least 120 nucleotides in length, at least 100 nucleotides in length, at least 90 nucleotides in length, at least 80 nucleotides in length, at least 70 nucleotides in length, at least 60 nucleotides in length, at least 50
15 nucleotides in length, at least 40 nucleotides in length, at least 30 nucleotides in length or less than 30 nucleotides in length. For example, in some embodiments, the 5' and 3' regions of the construct which have homology to the target promoter or gene may be about 30 nucleotides in length.

Alternatively, in some embodiments, the operator insertion construct may be a
20 single stranded nucleic acid, such as a synthetic oligonucleotide. Preferably, the single stranded nucleic acid is introduced into cells constitutively or conditionally expressing a protein which stabilizes single stranded DNA, such as the λ Bet protein or an analogous protein, as described in Ellis et al., PNAS 98: 6742-6746, 2001, the disclosure of which is incorporated herein by reference in its entirety. After
25 introduction of the operator insertion construct into a suitable cell, cells in which the operator has integrated by homologous recombination are identified. The cells may be identified by performing an amplification reaction or Southern blot. Alternatively, to identify cells in which expression of a gene required for proliferation is regulated by the inserted operator, cells which grow well under conditions in which the promoter
30 which is regulated by the operator is active at a higher level but not under conditions

in which the promoter is active at a lower level are identified. For example, cells which grow on media containing an inducer which increases the level of transcription from the promoter but not on media in which transcription is repressed may be identified. In some embodiments, the operator may be the *lac* operator and cells
5 which grow on media containing IPTG but not on media lacking IPTG may be identified. In some embodiments, the operator may be the *tet* operator and cells which grow on media containing tetracycline but not on media lacking tetracycline may be identified. In some embodiments, the operator may be the *xyl* operator and cells which grow on media containing xylose but not on media lacking xylose may be
10 identified. In some embodiments, the operator may be the *mal* operator and cells which grow on media containing maltose but not on media lacking maltose may be identified. In some embodiments, the operator may be the *trp* operator and cells which grow on media lacking tryptophan but not on media containing tryptophan may be identified. In some embodiments, the operator may be the *λcl* operator and cells
15 which grow at the permissive temperature but not at the restrictive temperature may be identified.

The operator insertion method provides a fast and inexpensive way to engineer directed insertions into the chromosome. In some embodiments, the operator insertion construct may lack a nucleic acid encoding an identifiable or selectable marker. The
20 method allows insertions of only the desired nucleic acid sequence without the necessity of introducing additional nucleic acid sequences such as sequences encoding identifiable or selectable markers. In addition, in embodiments where the native promoter is placed under the control of an operator, the expression level under conditions in which the promoter is not repressed or activated will be the
25 physiological level.

The operator insertion method may be performed in any desired prokaryotic organism. For example, in some embodiments, the method may be practiced in Gram-positive bacteria. In certain embodiments, the cell to be used with any of the methods described herein is selected from the group consisting of *Bacillus anthracis*,
30 *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium*

tetani, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*,
Lactococcus lactis, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium*
tuberculosis, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus*
epidermidis, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus*
5 *mutans* or any species falling within the genera of any of the above species.

In some embodiments, the operator insertion method is conducted in cells
which have an increased frequency of homologous recombination. For example, in
some embodiments, the method is conducted in cells which carry the *sbcA* and *recBC*
mutations, or in cells having mutations which confer similar phenotypes. In other
10 embodiments, the cells may constitutively or conditionally express one or more
proteins in the λ Red recombination system. For example, in some embodiments the
cells may constitutively or conditionally express the λ Gam, Bet, and Exo proteins or
other proteins having similar functions. In other embodiments, the cells may
constitutively or conditionally express the λ Bet protein or a protein having a similar
15 function.

In some embodiments, the promoter region for a target gene/operon is
identified either experimentally or bioinformatically. An oligonucleotide is designed
which matches at least 90, at least 70, at least 60, or at least 50 residues of the
promoter region. An operator is engineered to be present in the middle of the
20 oligonucleotide sequence. Thus, the operator will have at least 45, at least 35, at least
30, or at least 25 nucleotides of homology to the target region on each side. Examples
of operators include the *lac*, *tet*, *xyl*, *mal*, *trp*, or *λ cl* operators, though any other
operator known in the art can function. The oligonucleotide can be designed such that
the operator is inserted: i) upstream of the -35 box, ii) in between the -35 and the -10
25 box, or iii) downstream of the -10 box.

In some embodiments, the oligonucleotides are introduced into cells that
conditionally or constitutively express the λ Red (or homologous phage)
recombination system, which consists of the λ Gam, Exo, and Bet proteins (Ellis et
al., PNAS 98:6742-6746 (2001)). This recombination system efficiently recombines
30 oligonucleotide DNA into the bacterial chromosome. Specifically, in some

embodiments, the cells constitutively or conditionally express the λ Bet protein, which is a single strand binding protein that promotes annealing of complementary single strand and can mediate strand exchange, or another protein with an analogous function. Transformants are then plated on non-selective plates. To ascertain if a transforming is a recombinant, it is replica plated on plates that contain and lack the appropriate inducer. The plates are screened for an expected phenotype. For example, if the *xyI* operator is inserted downstream of the -10 box of an essential gene, then one would predict that recombinants would only be able to grow in the presence of inducer (xylose). For genes of known or expected phenotypes, e.g., those resulting in growth, transformants could be rapidly screened by plating on inducing (de-repressing) plates and then replica plating transformants on plates containing or lacking inducer. Those transformants that do not grow in the absence of inducer would contain the desired operator insertion.

Recombinants could also be verified by Southern or PCR analysis. This method is amenable to high throughput and thus many targeted recombinants could be constructed in a short time.

The operator insertion method also has the advantage of being portable to other bacteria by conditionally inducing the expression of the λ Bet protein. Additionally, this method could be employed to test if a gene is essential by screening for conditional growth, as discussed herein.

In some embodiments, the operator insertion method is conducted in cells which have an increased frequency of homologous recombination. For example, in some embodiments, the method is conducted in cells which carry the *sbcA* and *recBC* mutations, or in cells having mutations which confer similar phenotypes. In other embodiments, the cells may constitutively or conditionally express one or more proteins in the λ Red recombination system. For example, in some embodiments the cells may constitutively or conditionally express the λ Gam, Bet, and Exo proteins or other proteins having similar functions. In other embodiments, the cells may constitutively or conditionally express the λ Bet protein or a protein having a similar function.

In some embodiments, a double stranded operator insertion construct is introduced into a cell comprising the recBCD and sbcA mutations, recBC and sbcA mutations or mutations with analogous consequences. The double stranded operator insertion construct may or may not comprise a gene encoding a selectable or identifiable marker, but in preferred embodiments the double stranded operator insertion construct includes a gene encoding a selectable or identifiable marker. In other embodiments, a double stranded operator insertion construct is introduced into a cell constitutively or conditionally expressing the recE and recT proteins or proteins with analogous functions. In still other embodiments, a double stranded operator insertion construct is introduced into a cell constitutively or conditionally expressing the λ Gam, Bet, and Exo proteins or other proteins having similar functions. In such embodiments, cells in which the operator insertion construct has integrated are identified by obtaining colonies which express the identifiable marker or by growing the cells in appropriate selective media. If desired, homologous integration may be confirmed by performing a Southern blot or amplification reaction using a probe or primers which will yield a known result if homologous recombination has occurred.

In other embodiments, a single stranded operator insertion construct is introduced into the cell. Preferably, the single stranded operator insertion construct does not comprise a gene which encodes an identifiable or selectable marker, but in some embodiments a gene which encodes an identifiable or selectable marker may be included in the construct. In some embodiments, the single stranded nucleic acid may be generated by *in vitro* synthesis. In other embodiments, a circular single stranded DNA, such as M13, comprising the operator insertion construct may be fragmented by sonication and the fragments may be introduced into the cell. In still other embodiments, a single stranded operator insertion construct is introduced into a cell constitutively or conditionally expressing the λ Bet protein, the Bet and Gam proteins, the Bet, Gam and Exo proteins or other proteins having similar functions. Cells in which integration has occurred at the desired location are identified by performing colony hybridization, amplification reactions, or other methods familiar to those skilled in the art.

FIG. 4A illustrates one embodiment of the operator insertion method in which a linear double stranded regulatory element insertion construct comprising a nucleic acid encoding a selectable marker, a transcriptional terminator, and an operator is inserted into the chromosome of cells expressing the RecE and RecT gene products of the Rac prophage or gene products having analogous functions. FIG. 4B illustrates one embodiment of the operator insertion method in which a linear single stranded regulatory element insertion construct comprising a 5' portion homologous to a target sequence, a 3' portion homologous to a target sequence and an operator is inserted into the chromosome of a cell expressing the λ Beta and Gam proteins.

As discussed below, the operator insertion method may also be used to obtain cells in which the level or activity of a gene product required for proliferation is reduced. Such cells may be used in cell-based assays to identify compounds which inhibit proliferation.

In still another embodiment, a vector containing an outward-directed inducible promoter is randomly integrated into the genome of bacteria and clones containing these promoters that exhibit a reduction in viability and/or cell growth are identified. This technique is referred to as Random Insertional Cis-Antisense (random promoter insertion). This strategy involves the use of constructs that contain an inducible promoter (e.g., one of the promoters described herein) and a selectable marker to randomly integrate into the genome of a bacteria. When such constructs are integrated into or near an essential gene, induction of transcription from the inducible promoter will frequently reduce or eliminate normal expression of the essential gene. This event is particularly likely if the vector integrates such that transcription from the inducible promoter is in an antisense orientation relative to normal transcription of the endogenous gene. The integration in such cases is likely to be 3' to the gene, within the 3' untranslated region of the gene, within the coding region of the gene, or in the case of eukaryotes, within an intron of the gene. Induction of transcription in such integrants will dramatically interfere with normal sense transcription of the gene.

Since the expression of essential genes is required for proliferation of the organism, when such integration events occur within or near an essential gene,

induction of transcription will be lethal or will severely impair proliferation of the organism. Such conditional lethal or conditional growth clones can be readily isolated and the genes into which the vector has integrated can be determined by the cloning and sequencing techniques described herein. In this manner, the random promoter
5 insertion method can be used to discover essential bacterial genes.

Dominant negative mutants are another class of conditional lethal mutants that can be identified by using the random promoter insertion technique. These mutants can be generated when the vector integrates such that transcription from the inducible promoter is in the sense orientation relative to normal transcription of the endogenous
10 gene. In this case the integration site can be 5' to the gene, within the 5' untranslated region of the gene, within the coding region of the gene, or in the case of eukaryotes, within an intron of the gene. Because the dominant negative mutants will involve certain classes of essential genes, the dominant negative mutants are useful for identifying a subset of essential genes. Such conditional dominant negative mutants
15 can also be employed in a cell-based assay.

By one approach, a vector having the outward-directed inducible promoter is linearized by cutting with a restriction endonuclease and the linearized vector is transformed into bacteria. Sites of integration will be random but in many cases, the inducible promoter is oriented such that it produces an antisense RNA. Transformants
20 are robotically picked, grown, and gridded onto inducing and non-inducing media (e.g., media having and lacking xylose, media having and lacking tetracycline, and media having and lacking maltose). The gridded plates are then screened for transformants that exhibit a growth defect on inducing media relative to the phenotype on non-inducing media. Transformants that have a reduced viability and/or cell
25 growth in the presence of the inducer harbor a promoter that has integrated proximal to an essential gene. Once identified, the clone is sequenced to determine the identity of the essential gene. The genes that are affected by the random promoter insertion either directly or indirectly (e.g., by interacting with the RNA or protein that corresponds to the gene) causing a reduction in viability and/or cell growth are used as
30 drug discovery targets for high throughput drug screening.

In some embodiments of the present invention, complementary to genes required for the proliferation of a host organism from which the antisense molecules were originally obtained may be used to identify homologous antisense nucleic acids and the coding nucleic acids complementary thereto from cells or microorganisms other than the original host organism, to inhibit the proliferation of cells or microorganisms other than the original host organism by inhibiting the activity or reducing the amount of the identified homologous coding nucleic acid or homologous polypeptide in the cell or microorganism other than the original host organism, or to identify compounds which inhibit the growth of cells or microorganisms other than the original host organism as described below. For example, antisense nucleic acids complementary to proliferation-required genes from original host organism may be used to identify compounds which inhibit the growth of Gram-positive organisms such as *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species.

In one embodiment of the present invention, antisense nucleic acids complementary to the sequences identified as required for proliferation or portions thereof are operably linked to a regulatable fusion promoter, such as one of the fusion promoters described herein, contained in a vectors capable of replicating within a species other than the species from which the proliferation-required sequences were obtained. Expression vectors which contain a fusion promoter and one or more origins of replication that are functional in one or more organisms, such as the expression vectors described herein, may be used. For example, the vector may be capable of replicating and/or producing transcripts in one or more of the following Gram-positive organisms selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium*

diphtheriae, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling
5 within the genera of any of the above species.

Following the subcloning of the antisense nucleic acids, which are complementary to proliferation-required sequences or portions thereof from the host organism from which the stabilized antisense nucleic acids were originally obtained, into an expression vector under control of a regulatable fusion promoter capable of
10 replicating and producing stabilized transcripts in a second cell or microorganism of interest (i.e. a cell or microorganism other than the one from which the identified nucleic acids were obtained), the antisense nucleic acids are conditionally transcribed to test for bacterial growth inhibition. The coding sequences complementary to the antisense nucleic acids from the original host that, when transcribed, inhibit growth of
15 the second cell or microorganism are compared to the known genomic sequence of the second cell or microorganism to identify the homologous gene from the second organism. If the homologous sequence from the second cell or microorganism is not known, it may be identified and isolated by hybridization to the proliferation-required sequence of interest from the original host organism or by amplification using PCR
20 primers based on the proliferation-required nucleotide sequence of interest. In this way, nucleotide sequences which may be required for the proliferation of the second cell or microorganism may be identified. For example, the second microorganism may be selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium*
25 *diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species.

The homologous nucleic acid sequences from the second cell or microorganism which are identified as described above may then be operably linked to a promoter, such as one of the regulatable fusion promoter described herein, in an antisense orientation and introduced into the second cell or microorganism. The techniques described herein for identifying genes required for proliferation may thus be employed to determine whether the identified nucleotide sequences from a second cell or microorganism inhibit the proliferation of the second cell or microorganism. For example, the second microorganism may be selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species. If the stabilized antisense nucleic acid inhibits the proliferation of the second cell or microorganism, a cell-based assay, such as the one described herein, can be used to test and identify candidate antibiotic compounds.

In another embodiment of the present invention, screening of candidate antibiotic compounds can be performed directly by using the antisense molecule isolated from the original host organism. In this embodiment, an antisense nucleic acid comprising a nucleic acid complementary to the proliferation-required sequences from the original host organism or a portion thereof is transcribed in the second host from a regulatable fusion promoter such as one of the fusion promoters described herein. If the antisense molecule is transcribed so as to sufficiently alter the level or activity of a nucleic acid required for proliferation of the second host, the second host may be used directly in a cell-based assay, such as those described herein, to identify candidate antibiotic compounds.

The techniques for identifying drug discovery targets described above can be used to generate bacterial strains for high throughput drug screening methodologies. Because the promoters described herein can be tightly and/or finely regulated over a

wide range of maximal and basal transcription levels, one can selectively synthesize a precise amount of transcript or peptide sufficient to sensitize the cells without achieving complete lysis of cell population. For example, bacterial strains comprising the fusion promoters described herein are created using the EGI, promoter replacement, operator insertion, and random promoter insertion techniques. By
5 carefully adjusting the amount of inducer in the growth media, an optimal balance of slowed growth and sensitivity to compounds can be achieved. Libraries of test compounds are then screened against the sensitized cells, the effect on cell growth is monitored, and compounds that further reduce viability and/or cell growth are
10 identified. Through subsequent rounds of drug design, medicinal chemistry, and analysis, optimal antibiotics are developed using this inventive method.

Other embodiments concern an *in vivo* system for determining whether a gene is required for infection of a host by a bacterium. Such a system can also be used for validating the drug discovery targets and lead compounds identified by the approaches
15 described above. Accordingly, bacterial strains (e.g., including bacterial strains such as *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*,
20 *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species) having essential genes under the control of the regulatable promoters described herein are created using the EGI technology, promoter replacement technology, operator insertion technology or the random promoter insertion
25 technology and these bacteria are used to inoculate a suitable host animal (e.g., neutropenic mice). Two hours after inoculation, an experimental group of animals are either provided a solution containing inducer (if the EGI technology or random promoter insertion technology was used to create the compromised cells) or a solution containing a minimal amount of inducer (if the promoter replacement technology or
30 operator insertion technology was used to create the compromised cells).

Accordingly, the bacteria are induced to generate the transcript or peptide that interacts with the essential gene or gene product or transcription of an endogenous essential gene is reduced. Subsequently, the amount of viable bacteria remaining in the animal or tissue sample from the animal is determined. By using this approach, one can rapidly evaluate bacterial strains carrying essential genes that can be selectively downregulated *in vivo*. Additionally, one can use these animal models to screen lead compounds *in vivo* so as to identify new antibiotics.

10

EXAMPLE 1

Construction of Fusion Promoters Having a Single Operator

This example describes the construction of the fusion promoters that can be used to drive transcription of an RNA and/or expression of a peptide or protein in bacteria in a tightly regulated fashion. Many of the promoters described herein can function in multiple bacterial species including Gram-positive organisms such as *Staphylococcus* and *Enterococcus* species.

For construction of the Xyl-T5 fusion promoter, which is functional, for example, in *Staphylococcus aureus*, a *S. aureus*/*E. coli* shuttle vector, referred to as pRNLX2, was initially constructed. First, the multiple cloning site was removed from the *S. aureus* vector pRN5548, obtained from Richard Novick at the Skirball Institute at New York University, by digesting the vector with *HindIII*. The remaining vector sequence was self-ligated, which resulted in the formation of a new vector referred to as pRNHIII. Next, the pRNHIII vector was digested with *SalI* and the DNA was treated with Calf intestine alkaline phosphatase (CIP). Subsequently, the *SalI* linearized and CIP-treated pRNHIII vector was ligated to a 2.6 kb *XhoI*/*SalI* fragment of pLex5BA, which contains the plasmid origin and ampicillin resistance marker. (Diederich, L.*et al.*, *BioTechniques* 16: 916-923 (1994), the disclosure of which is incorporated herein by reference in its entirety). This vector, referred to as pRNLX2, contains both a *S. aureus* and an *E. coli* origin of replication. Thus, the vector can replicate in both Gram-negative and Gram-positive bacteria.

A *xyl* expression system was then constructed using the pRNLX2 shuttle vector. First, the *xyl* repressor gene and promoter were obtained from the plasmid pWH942 using PCR with the flanking primers 5'TGGCATTCTACGACTATAAC 3' (SEQ ID No.: 3) and 5'GCGAATTCTACATTTTAGTTGGTTAATTT 3' (SEQ ID No.: 4). (Schnappinger, D. *et al.*, *FEMS Microbiology Letters* 129: 121-128 (1995), the disclosure of which is incorporated herein by reference in its entirety). These primers contain *EcoRI* sites near their 5' ends and result in a PCR amplicon with trimmable *EcoRI* sites. The amplicons were digested with *EcoRI* and, subsequently, ligated into the pRN5548 vector, which had been linearized by digestion with *EcoRI* and *SwaI*. The resulting plasmid, called pXylRN, was used as a source for the *xyl* fragment for the final construct. The *xyl* fragment was excised from pXylRN with *EcoRI* and *PshAI* and was ligated to pRNLX2 that had been linearized with *EcoRI* and *PvuII*. The structure of the resultant plasmid was confirmed by sequencing the insert from both sides.

In the final step, the *xyl* promoter from pKLO20 was removed and replaced by the Xyl-T5 promoter. The *xyl* promoter and operator of pKLO20 is flanked by an *RsrII* site at its 5' end and the *EcoRI* site of the multiple cloning site at its 3' end. The promoter and operator were removed by digesting the plasmid with *RsrII* and *EcoRI*. The larger vector fragment of this digestion was purified from the smaller promoter fragment by preparative gel electrophoresis and a pair of oligonucleotides were designed having the following sequences:

5'GACCGTCATAAAAAATTTATTTGCTTTCAGGAAAATTTTCTGTATAATA
GATTCAAGTTAGTTTGTTTATTAAATTAACCACTAAAATGTAG3' (SEQ ID
No.: 5)

5'AAATTCTACATTTTAGTTGGTTAATTTAATAAACAACTAACTTGAATCTAT
TATACAGAAAAATTTTCCTGAAAGCAAATAAATTTTTATGACG3' (SEQ ID
No.: 6).

These oligonucleotides were annealed and, thereby, formed a double-stranded fragment. This fragment included: the RsrII sticky end (underlined), an AT rich region, the T5 promoter, the *xylO*, and the *EcoRI* sticky end (*italicized*). This fragment was then combined with the purified *RsrII-EcoRI* vector fragment of pKLO20 and ligated to form the pXyl-T5 vector.

To facilitate the construction of fusion promoters that function in *Enterococcus faecalis*, a reporter plasmid having a swappable promoter module located between a *xyl* repressor gene (*xylR*) and *xyl* operator (*xylO*) was designed. The *Lactococcus lactis* promoter probe vector pAK80 (described in Israelsen, et al. *Appl. Environ. Microbiol.* 61, 2540-47 (1995), the disclosure of which is incorporated by reference in its entirety), which contains replicons functional in both *Escherichia coli* (p15A origin) and *Enterococcus faecalis* (PCT1138 replicon) as well as a selectable marker for erythromycin resistance (*erm*), was digested at the unique HindIII and BamHI polylinker sites just upstream from the promoterless *Leuconostoc mesenteroides* β -galactosidase reporter genes (*lacL-lacM*). After digestion, the 11 kb vector fragment was purified from the 26 bp polylinker by preparative gel electrophoresis then ligated to a 1535 bp gel purified HindIII/BamHI fragment from pEP25 (SEQ ID NO: 7) which contains the CP25 promoter from *Lactococcus lactis* (SEQ ID NO.: 38) flanked upstream by the xylose repressor gene (*xylR*) and downstream by the xylose operator sequence (*xylO*). A portion of the resulting ligation mixture was transformed into competent *Escherichia coli* XL-1 Blue cells (Stratagene, La Jolla, CA) and 20 μ l, 75 μ l or 100 μ l of the transformation mixture was plated on BYGT medium containing 20 μ g/ml erythromycin. Isolated erythromycin-resistant transformants were picked and streaked to obtain single colony isolates. Plasmid DNA was then purified from representative single colony transformants. The presence of the chimeric plasmid pEPEF1 (SEQ ID NO.: 8) was confirmed by digesting purified plasmid preparations with HindIII/BamHI and detecting an excised 1535 bp fragment, which corresponds to the fragment containing the Xyl-CP25 fusion promoter, by gel electrophoresis.

FIG. 2 shows a region of pEPEF1 that contains the Xyl-CP25 fusion promoter. Immediately adjacent to the 5' end of the CP25 promoter and just downstream of *xylR*

lies a unique RsrII recognition site (element (a) in FIG. 2). A unique recognition site for XhoI (element (b) in FIG. 2) lies immediately adjacent to the 3' end of the CP25 promoter just upstream from *xylO*. The placement of these two unique restriction sites allow the CP25 promoter to be removed from pEPEF1 and replaced with promoter
 5 sequences from *Lactococcus* or other microbial species. For example, the CP25 promoter can be replaced with promoters having SEQ ID NOs.: 36 - 45.

In one example, a pEPEF1 derivative was created by replacing the CP25 promoter with the P59 promoter from *Lactococcus lactis*. Two complementary DNA oligonucleotides corresponding to the sequence of the P59 promoter were synthesized
 10 (SEQ ID NOs.: 9 and 10). Each sequence included a 5'-overhang end complementary to either an RsrII or XhoI overhang.

5'GACCGACATTAAATTCTTGACAGGGAGAGATAGGTTTGATAGAATATAA
 TAGTTGTC3' (SEQ ID NO.: 9)

15

5'TCGAGACAACTATTATATTCTATCAAACCTATCTCTCCCTGTCAAGAATT
 TAATGTCG3' (SEQ ID NO.: 10)

The underlined nucleotides comprise the 5'-overhang of the RsrII recognition
 20 site whereas the italicized nucleotides correspond to the 5'-overhang of the XhoI recognition site. The double stranded P59 RsrII/XhoI fragment was constructed by combining an equimolar amount of SEQ ID NOs.: 9 and 10, heating the mixture to 90°C to denature any secondary structure, and allowing the complementary sequences to anneal by slowly cooling the solution to room temperature. The double stranded
 25 P59 promoter sequence was then ligated into gel-purified, RsrII/XhoI-digested pEPEF1 reporter vector thereby creating the Xyl-P59 fusion promoter. This reporter vector was termed pEPEF1-P59.

It will be appreciated that any promoter sequence, including the promoters of
 30 SEQ ID NOs.: 36 - 45, can be synthesized to include an RsrII overhang at one end and a XhoI overhang at the other. Promoter sequences can be wildtype promoters known

from *Lactococcus lactis* and other Gram-positive species or promoters that have been modified to alter their transcription efficiency in gram-positive organisms. Example 7 provides a description of promoters that have been modified to alter their transcription efficiency in *Staphylococcus aureus* or *Enterococcus faecalis*.

- 5 The next example describes the construction of fusion promoters that have multiple operators.

EXAMPLE 2

Construction of Fusion Promoters Having a Plurality of Operators

- 10 This example describes the construction of fusion promoters comprising a plurality of operators. Fusion promoters such as Xyl/Lac-T5 and Xyl/Lac-P59 can be constructed by using a cloning strategy that is similar to the promoter swapping procedure described in Example 1. It will be appreciated that this procedure may be used to generate fusion promoters having two or more operators selected from the
- 15 group consisting of *xylO*, *tetO*, *trpO*, *malO*, *λc1O* and *lacO*.

- The *lac* operon is contained in many vectors including pGEM-11Zf(-), which is commercially available from Promega Corp. The sequence of the pGEM-11Zf(-) vector can be found in the NCBI database, Accession number X65314, the disclosure of which is incorporated herein by reference in its entirety. The *lac* operator
- 20 comprises the 17bp sequence: (TTGTTATCCGCTCACAA) (SEQ ID No. 11), which can be inserted in place of the 17 bp intervening sequence between the -35 Box and -10 Box of a variety of promoters, including the promoters of SEQ ID NOs.: 26 - 35.

- In one example, a Xyl/Lac-T5 promoter can be made by replacing the 17 bp sequence, which comprises element (3) in FIG. 1, with the 17 bp *lac* operator
- 25 sequence. To make the Xyl/Lac-T5 fusion promoter, two oligonucleotides are synthesized and annealed to create a double stranded fragment. The two oligonucleotides are provided below:

- 5'GACCGTCATAAAAAATTTATTTGCTTTGTGAGCGGATAACAATTATA
- 30 ATAGATTCAAGTTAGTTTGTATTAAATTAACCAACTAAAATGTAG3'
- (SEQ ID No.: 12)

5' AATTCTACATTTTAGTTGGTTAATTTAATAAAACAACTAACTTGAATCT
 ATTATAATTGTTATCCGCTCACAAAGCAAATAAATTTTTATGACG3'
 (SEQ ID No.: 13).

5

The resulting double stranded fragment includes: the *RsrII* sticky end (underlined portion of SEQ ID NO.: 12), an AT rich region found to increase the level of transcription from the T5 promoter in Gram-positive bacteria, the T5 promoter, the *lac* operator (*lacO*), the xylose operator (*xyIO*), and the *EcoRI* sticky end (italicized portion of SEQ ID NO.: 13). This double stranded fragment can be combined with the purified *RsrII/EcoRI* vector fragment of pKLO20 and ligated to form the vector pXyl/LacT5.

In a similar example, a Xyl/Lac-P59 promoter can be constructed by replacing the 17 bp sequence between the -35 Box and -10 Box regions (nucleotides 18 - 34) of SEQ ID NO.: 40) of P59 with the 17 bp *lac* operator sequence (SEQ ID NO.: 11). To make the Xyl/Lac-P59 fusion promoter, two oligonucleotides are synthesized and annealed to create a double stranded fragment. The two oligonucleotides are provided below:

20 5' GACCGACATTAAATTCTTGACATTGTGAGCGGATAACAATAGAAATATAA
 TAGTTGTC3' (SEQ ID NO.: 14)

5' *TCGAGACA*ACTATTATATTCTATTGTTATCCGCTCACAAATGTCAAGAATT
 TAATGTCG-3' (SEQ ID NO.: 15).

25

The underlined nucleotides comprise the 5'-overhang of the *RsrII* recognition site whereas the italicized nucleotides correspond to the 5'-overhang of the *XhoI* recognition site. The double stranded P59/Lac fusion promoter formed by the annealing of these two oligonucleotides is then ligated with purified, *RsrII/XhoI*-digested pEPEF1, which contains the *xyIO* operator, to form the vector pEPEF1-Lac/P59.

30

It will be appreciated that there are many equivalent techniques that can be used to construct fusion promoters having a plurality of operators. It will also be appreciated that operators other than *lacO* and promoters other than those described herein may be used in multi-operator constructs. For example, the fusion promoters
5 may comprise a plurality of operators selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO*, *λclO* or *lacO*. Furthermore, one of ordinary skill in the art will recognize that an operator may be positioned in a various locations with respect to the promoter provided that the operator retains sufficient ability to regulate transcription.

The next example describes the construction of expression vectors containing
10 regulatable fusion promoters.

EXAMPLE 3

Multipurpose Expression Vectors Containing Fusion Promoters

In this section, the construction and use of expression vectors comprising
15 fusion promoters is described. Because these expression vectors can accept a wide range of fusion promoters each having a range of transcriptional activity, these vectors are useful for finely modulating the amount of nucleic acid or polypeptide expressed inside a host cell throughout a wide range of concentrations.

It will be appreciated that a regulatable fusion promoter suite for use in
20 expression vectors can be constructed which provides a wide dynamic range of both basal and maximal transcriptional activities. Each fusion promoter within this suite can be tightly and finely regulated over its own characteristic range of transcriptional activity. Furthermore, the regulatable fusion promoters within the suite can comprise different promoters operably linked to the same operators thus providing for a
25 common means of regulation of the promoters within the suite. For example, the promoters within the set of regulatable fusion promoters within the suite can be linked to at least one operator contained in a promoterless expression vector that is capable of replicating in one or more Gram-positive organisms. In other embodiments, a promoter operator fusion construct may be inserted into the vector as a cassette.
30 Accordingly, a regulatable fusion promoter having an appropriate basal level and/or

maximal level of transcriptional activity can be selected from the suite for use in the appropriate application. As described below, fusion promoters can be quickly and easily exchanged in the expression vector as needed.

Fusion promoters prepared as described in the preceding sections can be incorporated into existing expression vectors using conventional techniques in molecular biology. Embodiments include, for example, expression vectors having promoters of SEQ ID NOs.: 36 – 45 linked to operators such as *xylO*, *xylO/lacO* or one or more operators including, but not limited to *tetO*, *malO*, *lacO*, *trpO*, and *λc1O*. In some embodiments, the expression vectors comprising the fusion promoters allow for the transcription and/or expression of a peptide protein or transcript in a plurality of Gram-positive and Gram-negative bacterial species (e.g., *Escherichia coli*, *Staphylococcus* species, and *Enterococcus* species).

These vectors can also comprise any of the following elements: genes encoding assayable markers such as enzymes that produce detectable products, genes encoding selectable markers such as polypeptides that control drug resistance, one or more unique restriction sites positioned downstream of the promoter such that inserts cloned into the site can be transcribed from the promoter, one or more transcription terminators such as a rho independent or rho dependent terminators which are positioned downstream of the unique cloning sites so as to terminate transcription of the nucleic acid insert, an origin of replication functional in Gram-negative species such as *Escherichia coli*, and an origin of replication functional in Gram-positive species such as *Staphylococcus* or *Enterococcus*.

For example, pXyl-T5 P15a is an expression vector which comprises the Xyl-T5 fusion promoter (see FIG. 5). This vector is 6858 bp in length and can be used to transcribe RNA and/or express a peptide or protein in Gram-positive bacteria including, but not limited to, *Staphylococcus* species. This vector comprises the Xyl-T5 fusion promoter positioned between the *xylR* gene and a multiple cloning site having unique restriction sites (e.g., *EcoRI*, *SacI*, *XhoI*, *KpnI*, *SmaI*, *XmaI*, *BamHI*, *XbaI*, and *Sall* sites), the beta-lactamase gene from pLex5BA the P15a replicon (which maintains the vector at approximately 15 copies/cell in *E. coli*), and the pC194

Staphylococcus replicon (which includes *palA*, *palB*, the pc194 origin of replication, the *rep* gene, and a chloramphenicol acetyl transferase gene that imparts chloramphenicol resistance in Gram-positive bacteria).

5 The pXyl-T5 p15a vector was constructed from pXyl-T5 by replacing the *E. coli* ColE1 origin of pXyl-T5, which allows replication of the vector at 100-200 copies per cell, with the *P15a* origin of pACYC184, which allows replication of the vector at approximately 15 copies per cell in *E. coli*. (Chang, A.C.Y. and Cohen, S.N., *J. Bacteriol.* 134:1141-1156 (1978), the disclosure of which is incorporated herein by reference in its entirety). The ColE1 origin lies within two closely linked *NotI* sites.

10 To obtain the Xyl plasmid having the ColE1 origin, the pXyl-T5 vector was digested with *NotI* and the larger vector fragment was purified by preparative gel electrophoresis. Next, a derivative of plasmid pACYC184, which includes the *P15a* origin as a similar *NotI* cassette, was digested with *NotI* and the smaller P15a-encoding fragment was gel purified. (Diederich, L.*et al.*, *BioTechniques* 16: 916-923

15 (1994), the disclosure of which is incorporated herein by reference in its entirety). The two purified *NotI* fragments were combined and ligated to form the vector pXyl-T5 P15a. Table 1 lists the components of the pXyl-T5 P15a vector and their locations.

TABLE 1

Type	Start Position	End Position	Name	Description
Gene	176	1327	<i>xylR</i>	The gene encoding repressor protein XylR
Promoter	1360	1455	A:T rich T5 promoter	The modified T5 promoter with upstream A:T rich box
Region	1455	1494	Multiple Cloning site	Multiple cloning site includes following unique restriction sites: <i>EcoRI</i> , <i>SacI</i> , <i>XhoI</i> , <i>KpnI</i> , <i>SmaI</i> , <i>XmaI</i> , <i>BamHI</i> , <i>XbaI</i> , <i>Sall</i>
Gene	2903	2043	<i>ampR</i>	The beta-lactamase gene from pLex5BA
Region	3037	3852	P15a	A replicon that allows maintenance of vector at approximately 15 copies per cell in <i>E. coli</i> hosts. Inserted as <i>NotI</i> cassette at the Start and End numbers of the vector sequence.
Region	4302	6659		<i>S. aureus</i> replicon from pC194. Includes the five features described below:
Region	4462	4302	<i>C palA</i>	With <i>palB</i> , imperfect direct repeats
Gene	5242	4592	<i>C Cat</i>	Gram-positive Chloramphenicol acetyl transferase gene-imparts chloramphenicol resistance in <i>S. aureus</i> .
Region	5562	5399	C ori	pC194 origin of replication
Gene	5699	6388	<i>rep</i>	gene for rep protein
Region	6659	6626	<i>C palB</i>	

A plasmid map of pXyl-T5 P15a is provided in FIG. 5. The nucleotide
5 sequence of pXyl-T5 P15a is provided in SEQ ID NO.: 16.

It will be appreciated that corresponding expression vectors which include fusion promoters comprising either single or multiple operators can be constructed using conventional techniques in molecular biology. For example, pXyl/Lac-T5 p15a can be constructed by digesting pXyl-T5 p15a with RsrII and EcoRI so as to remove
10 the Xyl-T5 fusion promoter. After purification, the linear vector fragment is ligated to an RsrII/EcoRI fragment comprising the pXyl/Lac-T5 promoter the construction of which was described in Example 2.

In another example, expression vectors functional in both *Escherichia coli* and *Enterococcus faecalis* were constructed by replacing the *lacL-lacM* reporter genes of

pEPEF1 with the multiple cloning site (MCS) and *rrnBt1t2* terminator of pLexP20. Specifically, pEPEF1 was digested with SmaI and SalI to remove the 3808 basepair fragment containing the *lacL-lacM* reporter genes. Following digestion, the large vector fragment was gel purified, the SalI overhanging end was made blunt with T4 DNA polymerase, and the fragment was circularized by ligating the blunt ends. The resulting plasmid, termed pEPEF2, was then digested with BamHI, gel purified, and blunted with T4 DNA polymerase. The ClaI/BstBI fragment containing the MCS and *rrnBt1t2* terminator was removed from pLexP20, gel purified, blunted using T4 DNA polymerase, then ligated to the blunt ended pEPEF2. A portion of the resulting ligation mixture was transformed into competent *Escherichia coli* and aliquots of the transformation mixture were plated on medium containing 20 µg/ml erythromycin. Isolated erythromycin-resistant transformants were picked and streaked to obtain single colony isolates. Plasmid DNA was then purified from representative single colony transformants. The presence and orientation of the MCS/terminator fragment was determined by PCR and DNA sequencing, respectively. A plasmid comprising the MCS/terminator fragment oriented such that the MCS was located adjacent to *xylO* was termed pEPEF3 (SEQ ID NO.: 17).

It will be appreciated that expression vectors such as those described above can be used to construct derivative expression vectors which include fusion promoters comprising either single or multiple operators. For example, the CP25 promoter can be removed from pEPEF3 by digestion of this vector with RsrII and XhoI. After purification, the linear vector fragment is ligated to an RsrII/XhoI fragment comprising P59. A pEPEF based expression vector having a generic promoter, termed pEPEF-X is shown in FIG. 6. Accordingly, many expression vectors derived from either pXyl-T5 or pEPEF3 can be constructed by exchanging the promoter sequence contained therein with the desired promoter sequence. In some embodiments, an expression vector is constructed comprising a promoter selected from SEQ ID NOs.: 36 - 45. pEPEF-X based expression vectors containing the *Lactococcus lactis* derived promoter sequences described herein, include pEPEF9 (SEQ ID NO.: 46), pEPEF14 (SEQ ID NO.: 47), pEPEF14* (SEQ ID NO.: 48), P1P2/EF14 (SEQ ID NO.: 49),

pEPEF18 (SEQ ID NO.: 50), pEPEF20 (SEQ ID NO.: 51), and pEPEF22 (SEQ ID NO.: 52).

Several methods of transcribing an RNA and/or expressing a peptide or protein are also embodiments included herein. More specifically, the fusion promoters described herein can be used to transcribe a sense or anti-sense RNA or can be used to express a recombinant peptide or protein in a cell.

The next example describes experiments that were performed to evaluate the ability of the Xyl-T5 promoter to transcribe RNA.

10

EXAMPLE 4

Activity of Fusion Promoters by Real-Time RT-PCR

To analyze the strength of the Xyl-T5 promoter that comprised the T5 promoter operably linked to an AT Box and an *xylO* operator, transcript accumulation experiments were conducted using real-time RT-PCR (Reverse Transcriptase Polymerase Chain Reaction). RT-PCR detects a specific RNA by PCR amplification of a reverse transcriptase synthesized cDNA copy of the RNA. With real time RT-PCR, the accumulation of PCR amplified DNA is measured at each amplification step using a specific fluorogenically labeled oligonucleotide probe. The probe anneals to the newly formed cDNA product and is subsequently cleaved by the 5' nuclease activity of the Taq DNA Polymerase during DNA amplification. Cleavage of the probe results in an increase in fluorescent dye signal, which is used to monitor accumulation of the specific PCR product at each PCR amplification step. The PCR cycle time at which logarithmic amplification is reached is proportional to the initial amount of RNA template in the reaction.

The levels of transcription from the Xyl-T5 promoter were evaluated in the following experiment. Fresh overnight colonies of *Staphylococcus aureus* RN4420 containing plasmid a plasmid having the Xyl-T5 promoter were added to 10 ml of Luria Broth (LB) that was supplemented with 15 µg/ml chloramphenicol (LB CM15). The cells were grown at 37°C with shaking until early log phase (OD600 between 0.1 and 0.2) at which point they were diluted 1/10 into pre-warmed LB CM15. After one hour of growth at 37°C with shaking, three 1 ml aliquots of cells were pelleted and

quick frozen on dry ice. The remaining suspension was split into two flasks. Xylose was added to a final inducer concentration of 2.0% in one flask, whereas the second flask remained non-induced (control). The resulting transcript is approximately 300 nucleotides in length and corresponds to vector sequence.

5 Immediately after xylose induction (2 minute time point) and at 10, 20, 30 and 60 minute time points, (three) 1 ml aliquots of cells were pelleted from each flask (with and without xylose) and quick frozen on dry ice. Cell pellets were stored at -80°C. Total RNA was purified from each sample using a Qiagen 96 well RNeasy kit. Approximately, 1 ng of RNA (two samples for each time point) was used as template
10 for RT-PCR with an oligonucleotide primer set (forward and reverse primers, and a fluorogenic probe) for specific detection of the 3' tail of the Xyl-T5 transcript. The reaction consisted of 300 nM each of the forward and reverse primers; 100 nM fluorogenic probe; 0.75 U AmpliTaq Gold (PE Biosystems); 7.5 U MultiScribe Reverse Transcriptase (PE Biosystems), 1x Buffer A (PE Biosystems); 5.8 mM MgCl,
15 and 333 nM of each dNTP. The RT-PCR conditions were 48°C for 30 minutes, 95°C for 10 minutes, followed by 40 cycles of 95°C for 15 seconds and 60°C for 1 minute. RT-PCR reactions and detection were done in an ABI Prism 7700 Sequence Detection System.

 The results of the experiment described above are shown in FIG. 7. Within 2
20 minutes of xylose induction, greater than a 30 fold excess of RNA was detected, as compared to the non-induced controls. At 60 minutes, greater than a 130 fold excess of RNA was detected in the xylose induced samples, as compared with non-induced bacteria. This example demonstrates that the fusion promoters described herein can be used to efficiently transcribe an RNA. Further, the data described herein verify that
25 in the absence of xylose or an analog thereof Xyl-T5 promoter-driven transcription can be effectively repressed.

 It will be appreciated that the fusion promoters described above can be used to construct a regulatable fusion promoter suite which provides a wide dynamic range of both basal and maximal transcriptional activities. Each fusion promoter within this
30 suite can be tightly and finely regulated over its own characteristic range of

transcriptional activity. Furthermore, the regulatable fusion promoters within the suite can comprise different promoters operably linked to the same operators thus providing for a common means of rapidly regulating the promoters within the suite. In some embodiments, the promoters within the set of regulatable fusion promoters within the suite can be linked to at least one operator contained in a promoterless expression vector that is capable of replicating in one or more Gram-positive organisms. Accordingly, a regulatable fusion promoter having an appropriate basal level and/or maximal level of transcriptional activity can be selected from the suite for use in the appropriate application.

The example below demonstrates the relative strengths of the fusion promoters described herein.

EXAMPLE 5

Activity of Fusion Promoters by Expression of Reporter Genes

This example describes the use of reporter gene constructs to measure the activity of fusion promoters such as those constructed in Examples 1 and 2. In one experiment, the pXyl-T5 and pXyl-T5 P15a vectors were tested to determine the relative level of expression of a reporter gene in both *Escherichia coli* and *Staphylococcus aureus*. To test the ability of the Xyl-T5 promoter to induce transcription and express a heterologous polypeptide, a construct was made in which the reporter gene was operably linked to the Xyl-T5 promoter. A pXyl-T5 vector was digested with SacI and Sall, and the linearized vector was purified by preparative gel electrophoresis. The linearized vector fragment was then ligated with a fragment containing the reporter gene. This construct was named pXyl-T5-Reporter. The pXyl-T5-Reporter vector containing the reporter gene under the control of the Xyl-T5 promoter was then introduced into *S. aureus* and the cells were cultured in the presence and absence of 2.0% xylose. The transcription and translation of the gene product encoded by the reporter gene was evident when xylose was provided in the growth medium but in the absence of an agent that inhibits the binding of the xylose repressor to the xylose operator, production of the gene product encoded by the reporter gene was not detectable.

When the vector containing the reporter gene under the control of the Xyl-T5 promoter was transformed in *Escherichia coli*, it was noted that expression of the gene product encoded by the reporter gene was very high, and uncontrollable either in the presence or absence of xylose. This may be because *Escherichia coli* does not express the XylR repressor. In those embodiments where it is desirable that *Escherichia coli* serve as a surrogate host, pXyl-T5 p15a, which has a reduced copy number in *Escherichia coli* can be used.

To evaluate the effect of the P15a origin on the level of reporter gene expression in *Staphylococcus aureus* and *Escherichia coli* in the presence and absence of xylose, a construct similar to pXyl-T5-Reporter was made, using the pXyl-T5 p15a vector. This construct was denoted pXyl-T5 p15a-Reporter. A comparison of the induction of reporter gene expression of pXyl-T5-Reporter and pXyl-T5 p15a-Reporter showed that the two vectors functioned identically when transformed and expressed in *Staphylococcus aureus*, but the P15a variant showed a markedly reduced expression of the reporter gene when transformed in *Escherichia coli*. The demonstrates that the p15a origin of replication decreases reporter gene expression by reducing the copy number of the expression vector thereby reducing the number Xyl-T5 promoters in the cell.

Fusion promoters contained on pEPEF1 and its derivatives were evaluated to determine their strength in *Enterococcus faecalis* by measuring the β -galactosidase activity produced by expression of the *lacL-lacM* reporter genes under induced and uninduced conditions. In one experiment, derivatives of pEPEF1 were prepared by replacing CP25 with the *Lactococcus lactis* derived promoters P32 (SEQ ID NO.: 39), P59 (SEQ ID NO.: 40), or P1P2 (SEQ ID NO.: 42) as described in Example 1. pEPEF1 and the resulting derivatives were each separately transformed into electrocompetent *Enterococcus faecalis* and the cells were cultured in the absence and the presence of 5% xylose. The β -galactosidase activity produced by the expression of the reporter genes was measured essentially as described in Israelsen et al. *Appl. Environ. Microbiol.* 61:2540-47 (1995), the disclosure of which is incorporated herein by reference in its entirety, with the following modifications. After incubation with or

without xylose for four hours, the cells were harvested by centrifugation then resuspended in 1 ml of Z buffer comprising 0.06 M $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 0.04 M $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$, 0.01 M KCl, 0.001 M $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.05 M β -mercaptoethanol, at pH 7.0, with 0.6 ml of glass beads (150-212 micron diameter). The cells were lysed
5 using a bead beater at maximum speed for 1 minute. After centrifugation, the supernatant was removed, vortexed with one drop of 0.1% SDS and 2 drops of chloroform at high speed for 10 seconds, then incubated with 200 μl of a 4 mg/ml *o*-nitrophenyl- β -D-galactopyranoside (ONPG) solution at 37°C. After approximately 20 minutes, the reaction was stopped by adding 250 μl of 1M sodium carbonate. β -
10 galactosidase activity was quantitated spectrophotometrically. All assays were performed in triplicate.

FIG. 8 displays the amount of β -galactosidase activity produced by each fusion promoter in *Enterococcus faecalis* in the presence and absence of xylose. Promoters CP25 and P32 both produce very little β -galactosidase activity in the absence of
15 xylose; however, in the presence of 5% xylose, both of these promoters express enough β -galactosidase to produce approximately 25 Miller units of activity. The activities of both P59 and P1P2 are much greater than that of either CP25 or P32. In the presence of xylose, P59 and P1P2 each produce between 125 and 150 Miller units of β -galactosidase activity. In the absence of xylose, P59 is strongly repressed having
20 less than 4% of its induced activity. By contrast, P1P2 produces about 20% of its induced activity even in the absence of xylose.

The above data shows that the fusion promoters described herein differ in the levels of transcription which they provide and the levels of transcription under non-inducing conditions. Some promoters are highly active under induced conditions
25 whereas others show only a modest increase in activity upon induction. Furthermore, some promoters described herein may be strongly repressed in the absence of xylose whereas the activity of other promoters may only be partially inhibited.

It will be appreciated that the fusion promoters described above can be used to construct a regulatable fusion promoter suite which provides a wide dynamic range of
30 both basal and maximal transcriptional activities. Each fusion promoter within this

suite can be tightly and finely regulated over its own characteristic range of transcriptional activity. Furthermore, the regulatable fusion promoters within the suite can comprise different promoters operably linked to the same operators thus providing for a common means of regulation of the promoters within the suite. In some
5 embodiments, the promoters within the set of regulatable fusion promoters within the suite can be linked to at least one operator contained in a promoterless expression vector that is capable of replicating in one or more Gram-positive organisms. Accordingly, a regulatable fusion promoter having an appropriate basal level and/or maximal level of transcriptional activity can be selected from the suite for use in the
10 appropriate application.

In another experiment, the activity of the CP25 fusion promoter was compared to the activity of Xyl-T5. To produce a pXyl-T5 p15a vector containing the *lacL-lacM* reporter, the purified 4 kb HindIII/Sall fragment of pAK80 which contains the *lacL-lacM* reporter construct was ligated into purified pXyl-T5 p15a which had been
15 linearized using HindIII and Sall. The resulting construct was termed pXyl-T5 p15a LacLM. To compare the activity of the T5 fusion promoter with that of CP25, pXyl-T5 p15a LacLM and pEPEF1 were transformed separately into *Enterococcus faecalis* and the cells were cultured in the presence of 1% and 5% xylose and in its absence. The β -galactosidase activity produced by the expression of the reporter genes was
20 measured using the Miller assay as described above.

FIG. 9 displays the amount of β -galactosidase activity produced by each fusion promoter under induced and uninduced conditions. In the absence of xylose, both the fusion promoters are strongly repressed in *Enterococcus faecalis*. When induced with 1% xylose, CP25 has about twice the activity of Xyl-T5 in *Enterococcus faecalis*. At
25 5% xylose, CP25 produces about 35 Miller units of activity compared to about 22 Miller units by Xyl-T5.

To determine whether there is a difference in the level of gene expression from pXyl-T5 p15a in *Staphylococcus aureus* and *Enterococcus faecalis*, pXyl-T5 p15a LacLM was also transformed into *Staphylococcus aureus* which was grown in the
30 presence of 1% and 5% xylose and in its absence. The β -galactosidase activity

produced by the expression of the reporter genes was measured using the Miller assay as described above.

FIG. 9 shows that the activity of Xyl-T5 is similar in *Staphylococcus aureus* and *Enterococcus faecalis*. When induced with 5% xylose in each organism, the Xyl-T5 fusion produces approximately 22 Miller units of β -galactosidase activity.

The above data shows that the fusion promoters disclosed herein have the ability to function in a variety of Gram-positive organisms and to maintain their characteristic activities when transferred from one host to another. This result is in contrast to the alteration in activity that is seen when the promoter is transferred to Gram-negative organisms. One example is the unregulated activity of Xyl-T5 in *Escherichia coli*.

The example below demonstrates that the activity of the fusion promoters described herein can be finely regulated over a range of inducer concentrations. In other words, the activity of these fusion promoters is titratable.

15

EXAMPLE 6

Titratable Expression from Fusion Promoters

This example shows that the activity of promoters described herein is titratable with inducer thus allowing a reproducible, incremental modulation of promoter activity by altering the concentration of inducer. Each fusion promoter described herein possesses its own characteristic activity over a range of xylose concentrations. Within this range, the amount of RNA transcribed from a fusion promoter can be finely regulated in response to the concentration of xylose used to induce the promoter.

In one example, the derivative of pEPEF1 containing the P59 promoter (pEPEF1-P59) was transformed into *Enterococcus faecalis* and the transformants were cultured in the presence of increasing concentrations of xylose (from 0.1% to 10%) and in its absence. The β -galactosidase activity produced by the expression of the *lacL-lacM* reporter genes was measured using the Miller assay as described in

Example 5. FIG. 10 shows that the activity of the P59 promoter increases in response to induction with increasing concentrations of xylose.

The next example describes modifications to fusion promoters that effect their transcriptional activity.

5

EXAMPLE 7

Modifications of Fusion Promoters that Alter Transcription Efficiency

This example describes the construction of fusion promoters that have been modified to have altered transcriptional activity in Gram-positive organisms. More specifically, modified fusion promoters, which are derived from wildtype promoters, 10 have been constructed by creating mutations in the promoter sequence which either increase or diminish the strength of the promoter.

In one example, the strong *Lactococcus lactis* promoter, PL (SEQ ID NO.: 44), was modified to increase its activity in Gram-positive organisms. First, a regulatable 15 PL fusion promoter was constructed by linking the PL promoter to a xylose operator using the methods described previously. Examination of the PL sequence revealed that changing the C at position 45 of SEQ ID NO.: 44 to an A would make the -10 Box region of promoter PL identical to -10 Box consensus site reported for Gram-positive organisms (TATAAT). PL10, a derivative of PL which incorporates this 20 single basepair modification, was constructed by annealing the complementary synthetic oligonucleotides of SEQ ID NOs.: 18 and 19 using the procedure described in Example 1.

5' GACCGTTTCGTGAACTTTTTGTTGACAAAGATAAAAACACATGATATA
25 ATTAAATCAC3' (SEQ ID NO.: 18)

5' TCGAGTGATTTAATTATATCATGTGTTTTTATCTTTGTCAACAAAAAAGTT
CACGAAACG3' (SEQ ID NO.: 19)

The underlined nucleotides comprise the 5'-overhang of the RsrII recognition site whereas the italicized nucleotides correspond to the 5'-overhang of the XhoI recognition site. The double stranded promoter formed by the annealing of these two oligonucleotides was then ligated with purified, *RsrII/XhoI*-digested pEPEF1 to form a pEPEF1 derivative containing the Xly-PL10 fusion promoter. This derivative was termed pEPEF1-PL10. To determine the effect of this modification, the strengths of PL10, PL and P59 were compared by separately transforming pEPEF1 derivatives containing the appropriate promoter into *Enterococcus faecalis* then culturing the cells in the absence or presence of 5% xylose. The β -galactosidase activity produced by the expression of the *lacL-lacM* reporter genes was measured using the Miller assay as described in Example 5.

FIG. 11 compares the strengths of the PL, PL10 and P59 fusion promoters expressing the *lacL-lacM* reporter genes. When induced with 5% xylose, the activity of PL10 is about 20% greater than that of the wildtype PL promoter. Both PL and PL10 have substantially greater activity than P59 under induced conditions. In the absence of xylose, P59 is almost completely repressed whereas both PL and PL10 show significant basal activity.

In another example, P59 was modified to increase its activity in Gram-positive organisms. Examination of the P59 sequence revealed that changing the G at position 37 of SEQ ID NO.: 40 to a T would make the -10 Box region of promoter P59 identical to -10 Box consensus site in Gram-positive organisms. Additionally it appeared that addition of an AT rich region upstream of the -35 Box might enhance promoter activity. A derivative of P59 which incorporates these modifications was constructed by annealing the complementary synthetic oligonucleotides of SEQ ID NOs.: 20 and 21 using the procedure described in Example 1.

5'GACCGAAAAATGACAGTTTATTCTTGACAGGGAGAGATAGGTTTGATAT
AATATAATAGTTGTC3' (SEQ ID NO.: 20)

5' TCGAGACA ACTATTATATTATATCAAACCTATCTCTCCCTGTCAAGAATA
 AACTGTCATTTTTCG3' (SEQ ID NO.: 21)

5 The underlined nucleotides comprise the 5'-overhang of the *RsrII* recognition site whereas the italicized nucleotides correspond to the 5'-overhang of the *XhoI* recognition site. The double stranded promoter formed by the annealing of these two oligonucleotides was then ligated with purified, *RsrII/XhoI*-digested pEPEF1, transformed into *E. faecalis*, and assayed as previously described. When compared to the unmodified P59, the derivative had about 20% higher activity.

10 It will be appreciated that modified fusion promoters, such as those described above, can be used to construct a regulatable fusion promoter suite which provides a wide dynamic range of both basal and maximal transcriptional activities. Modifying regulatable fusion promoters so as to increase their activity expands the range of both basal and maximal transcriptional activities of a suite of regulatable fusion promoters.

15 Each modified fusion promoter within this suite can be tightly and finely regulated over its own characteristic range of transcriptional activity. Furthermore, the modified regulatable fusion promoters within the suite can comprise different promoters operably linked to the same operators thus providing for a common means of regulation of the promoters within the suite. In some embodiments, the modified

20 promoters within the set of regulatable fusion promoters within the suite can be linked to at least one operator contained in a promoterless expression vector that is capable of replicating in one or more Gram-positive organisms. Accordingly, a regulatable fusion promoter having an appropriate basal level and/or maximal level of transcriptional activity can be selected from the suite for use in the appropriate

25 application.

In yet another example, the Xyl-T5 fusion promoter was modified to decrease its activity in Gram-positive organisms. By changing four nucleotides in the -35 Box of SEQ ID NO.: 36 (G to T at position 18, C to A at position 19, T to C at position 20 and T to A at position 21) and one in the -10 Box of SEQ ID NO.: 36 (A to T at

30 position 43) the Xyl-T5 promoter could be altered to include -10 and -35 elements that

are identical to those of the weak *xylA* promoter. (For a description of the *xylA* promoter, see Sizemore *et al.*, *Mol. Gen. Genet.* 227:377-384 (1991) and Schnappinger, D. *et al.*, *FEMS Microbiology Letters* 129: 121-128 (1995), the disclosures of which are incorporated herein by reference in their entireties). Xyl-T5-
 5 DD, a derivative of Xyl-T5 which incorporates these modifications was constructed by annealing the complementary synthetic oligonucleotides of SEQ ID NOs.: 22 and 23 using the procedure described in Example 1.

5'GACCGTCATAAAAAATTTATTTTACATCAGGAAAATTTTCTGTATATTA
 10 GATTCAAGTTAGTTTGTATTAAATTAACCAACTAAAATGTAG3' (SEQ ID NO.: 22)

5'AA^{TT}CTACATTTTAGTTGGTTAATTTAATAAACAACTAACTTGAATCTA
 15 ATATACAGAAAAATTTTCCTGATGTAAATAAATTTTTATGACG3' (SEQ ID NO.: 23)

The underlined nucleotides comprise the 5'-overhang of the *RsrII* recognition site whereas the italicized nucleotides correspond to the 5'-overhang of the *EcoRI* recognition site. The double stranded promoter formed by the annealing of these two
 20 oligonucleotides was then ligated with purified, *RsrII/EcoRI*-digested pXyl-T5-Reporter (prepared as described in Example 5). The pXyl-T5-Reporter vector was then introduced into *S. aureus* and the cells were cultured in the presence and absence of 2.0% xylose and expression levels were measured. Xyl-T5-DD had only about 10% of the activity of unmodified Xyl-T5. The nucleotide sequence of pXyl-T5-DD
 25 P15a, a Xyl-T5-DD based expression vector, is provided in (SEQ ID NO.: 53).

It will be appreciated that modified fusion promoters, such as those described above, can be used to construct a regulatable fusion promoter suite which provides a wide dynamic range of both basal and maximal transcriptional activities. Modifying regulatable fusion promoters so as to decrease their activity expands the range of both
 30 basal and maximal transcriptional activities of a suite of regulatable fusion promoters.

Each modified fusion promoter within this suite can be tightly and finely regulated over its own characteristic range of transcriptional activity. Furthermore, the modified regulatable fusion promoters within the suite can comprise different promoters operably linked to the same operators thus providing for a common means of regulation of the promoters within the suite. In some embodiments, the modified promoters within the set of regulatable fusion promoters within the suite can be linked to at least one operator contained in a promoterless expression vector that is capable of replicating in one or more Gram-positive organisms. Accordingly, a regulatable fusion promoter having an appropriate basal level and/or maximal level of transcriptional activity can be selected from the suite for use in the appropriate application.

It will be appreciated that the methods described herein can be used to create promoter sequence modifications that either enhance or reduce promoter activity. These modifications can be made either alone or in combination. Among these changes are modifications of nucleotides within the promoter's -35 and -10 Box elements which make the promoter more or less like the consensus -35 Box and -10 Box sequences (TTGACA and TATAAT, respectively). Other modifications include altering the length of the region between the operator or transcription start site and the -10 Box, altering the length and/or composition of the region between the -10 Box and the -35 Box, and altering the length and/or composition of the AT rich region upstream of the -35 Box (i.e. the -45 Box). It will also be appreciated that modifications which affect promoter activity can include addition of elements such as the CRE region for catabolite repression. These additional elements can be further modified so that they are more or less like their corresponding consensus sequence. Promoters which incorporate one or more modifications similar to those described above are represented by SEQ ID NOs.: 37, 41, 43 and 45.

The experiments described in the Examples 4 -7 verify that the fusion promoters disclosed herein can be used to tightly and finely regulate transcription of an RNA or expression of a peptide or a protein throughout a range of activity levels.

It will be appreciated that both modified and unmodified fusion promoters, such as those described above, can be used to construct a regulatable fusion promoter suite which provides a wide dynamic range of both basal and maximal transcriptional activities. Modifying regulatable fusion promoters so as to increase or decrease their activity expands the range of both basal and maximal transcriptional activities of a suite of regulatable fusion promoters. The addition of unmodified fusion promoters having unique basal and/or maximal transcriptional activities also expands the dynamic range of the suite. Each regulatable fusion promoter within this suite can be tightly and finely regulated over its own characteristic range of transcriptional activity. Furthermore, the regulatable fusion promoters within the suite can comprise different promoters operably linked to the same operators thus providing for a common means of regulation of the promoters within the suite. In some embodiments, the promoters within the set of regulatable fusion promoters within the suite can be linked to at least one operator contained in a promoterless expression vector that is capable of replicating in one or more Gram-positive organisms. Accordingly, a regulatable fusion promoter having an appropriate basal level and/or maximal level of transcriptional activity can be selected from the suite for use in the appropriate application. Such a suite of promoters is useful in the methods of identifying genes required for proliferation or methods of identifying compounds which inhibit proliferation described herein.

The examples below discusses methods that use fusion promoters to identify essential genes and compounds having potential antibiotic activity.

EXAMPLE 8

Identification of Genes Required for Cellular Proliferation by Expressing Proliferation- Inhibiting Antisense Nucleic Acids from a Fusion Promoter

This example describes the examination of a library of candidate antisense nucleic acid operably linked to fusion promoters to identify essential genes in *Staphylococcus aureus* and *Enterococcus faecalis*. Although the following examples

utilize the fusion promoters Xyl-T5, CP25 and P59, it will be appreciated that any of the fusion promoters described herein may be used.

It will be appreciated that a regulatable fusion promoter suite which provides a wide dynamic range of both basal and maximal transcriptional activities can be used to facilitate the discovery of proliferation-required genes. Each fusion promoter within this suite can be tightly and finely regulated over its own characteristic range of transcriptional activity. Furthermore, the regulatable fusion promoters within the suite can comprise different promoters operably linked to the same operators thus providing for a common means of regulation of the promoters within the suite. In some embodiments, the promoters within the set of regulatable fusion promoters within the suite can be linked to at least one operator contained in a promoterless expression vector that is capable of replicating in one or more Gram-positive organisms. Accordingly, a regulatable fusion promoter having an appropriate basal level and/or maximal level of transcriptional activity can be selected from the suite based on the expression level of the proliferation-required genes that are targeted for discovery.

In the methods for identifying genes encoding gene products required for cellular proliferation using antisense nucleic acids expressed from an inducible fusion promoter, random genomic fragments are obtained from the organism in which it is desired to identify genes required for cellular proliferation. The random genomic fragments may be generated by a partial digestion with a restriction enzyme, mechanical shearing, using techniques such as sonication and nebulization, or DNaseI digestion. The random genomic fragments are operably linked to a regulatable fusion promoter in a vector, such as one of the expression vectors described herein. In those instances where the inserted genomic fragments are in an antisense orientation with respect to the promoter, the transcript produced is complementary to at least a portion of an mRNA encoding a gene product such that they interact with sense mRNA produced from various genes and thereby decrease the translation efficiency or the level of the sense messenger RNA (mRNA) thus decreasing production of the protein encoded by these sense mRNA molecules. In cases where the sense mRNA encodes a protein required for proliferation, cells grown under inducing conditions fail to grow or grow at

a substantially reduced rate. Additionally, in cases where the transcript produced is complementary to at least a portion of a non-translated RNA and where that non-translated RNA is required for proliferation, cells grown under inducing conditions also fail to grow or grow at a substantially reduced rate. In contrast, cells grown under non-inducing conditions grow at a normal rate.

The genes to which the antisense nucleic acids are complementary are then identified and utilized in the methods of the present invention. Thus, to identify genes required for cellular proliferation, the extent of proliferation of cells containing the vectors in the presence of an inducer which induces transcription from the regulatable fusion promoter is compared to the extent of proliferation of cells in the absence of the inducer. Those cells which grow well in the absence of the inducer but exhibit significantly reduced proliferation in the presence of the inducer contain a vector encoding an antisense nucleic acid complementary to at least a portion of a gene required for cellular proliferation.

Use of the above method to identify genes required for cellular proliferation in *E. coli*, *Staphylococcus aureus*, *Salmonella typhimurium*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* and *Enterococcus faecalis* has been described in the following U.S. Patent Applications, the disclosures of which are incorporated herein by reference in their entireties: U.S. Patent Application Serial Number 09/815,242, filed March 21, 2001; U.S. Patent Application Serial Number 09/492,709, filed January 27, 2000; U.S. Patent Application Serial Number 09/711,164, filed November 9, 2000; U.S. Patent Application Serial Number 09/741,669, filed December 19, 2000; U.S. Patent Application Serial Number 09/815,242, filed March 21, 2001; and U.S. Patent Application Serial Number 09/948,993, filed September 06, 2001. The methods used to identify genes required for cellular proliferation in *Staphylococcus aureus*, and *Enterococcus faecalis* are summarized below.

Nucleic acids involved in proliferation of *Staphylococcus aureus* were identified using the EGI technique as follows. A shotgun library of *Staphylococcus aureus* genomic fragments was cloned into the vector pXylT5-P15a, which harbors the Xyl-T5 inducible promoter. The vector was linearized at a unique *Bam*HI site

immediately downstream of the XylT5 promoter/operator. The linearized vector was treated with shrimp alkaline phosphatase to prevent reclosure of the linearized ends. Genomic DNA isolated from *Staphylococcus aureus* strain RN450 was fully digested with the restriction enzyme *Sau3A* or alternatively, partially digested with DNase I
5 and "blunt-ended" by incubating with T4 DNA polymerase. Random genomic fragments between 200 and 800 base pairs in length were selected by gel purification. The size-selected genomic fragments were added to the linearized and dephosphorylated vector at a molar ratio of 0.1 to 1, and ligated to form a shotgun library.

10 The ligated products were transformed into electrocompetent *E. coli* strain XL1-Blue MRF' (Stratagene) and plated on LB medium with carbenicillin at 100 µg/ml. Resulting colonies numbering 5×10^5 or greater were scraped and combined, and were then subjected to plasmid purification.

The purified library was then transformed into electrocompetent
15 *Staphylococcus aureus* strain RN4220. Resulting transformants were plated on LBG agar with chloramphenicol at 15 µg/ml in order to generate 100 to 150 platings at 500 colonies per plating. The colonies were subjected to robotic picking and arrayed into wells of 384 well culture dishes. Each well contained 100µl of LBG + CM15 liquid medium. Inoculated 384 well dishes were incubated 16 hours at 37°C, and each well
20 was robotically gridded onto solid LBG + CM15 medium with or without 2% xylose. Gridded plates were incubated 16 hours at 37°C, and then manually scored for arrayed colonies that were growth-compromised in the presence of xylose. Such growth sensitivity occurred at a frequency of one in sixty arrayed colonies.

Arrayed colonies that were growth-sensitive on medium containing 2% xylose,
25 yet were able to grow on similar medium lacking xylose, were subjected to further growth sensitivity analysis as follows: Colonies from the plate lacking xylose were manually picked and inoculated into individual wells of a 96 well culture dish containing LBG + CM15, and were incubated for 16 hours at 37°C. These cultures were robotically diluted 1/100 into fresh medium and allowed to incubate for 4 hours
30 at 37°C, after which they were subjected to serial dilutions in a 384 well array and

then gridded onto media with and without 2% xylose. After growth for 16 hours at 37°C, the arrays of serially diluted spots that resulted were compared between the two media. Spots that grew similarly on both media were scored as a negative and corresponding colonies were no longer considered. Spots on xylose medium that
5 failed to grow to the same serial dilution compared to those on the non-xylose plate were given a score based on the differential, i.e. should the spots on xylose only grow to a serial dilution of -4 while they were able to grow to -8 on the non-xylose plate, then the corresponding transformant colony received a score of "4" representing the log difference in growth observed.

10 Nucleic acids involved in proliferation of *E. faecalis* were identified as follows. A shotgun library of *E. faecalis* genomic fragments was cloned into either pEPEF3 or pEPEF14, which contain the CP25 or P59 promoter, respectively, regulated by the xyl operator/repressor. The vector was linearized at a unique *Sma*I site immediately downstream of the promoter/operator. The linearized vector was
15 treated with alkaline phosphatase to prevent reclosure of the linearized ends. Genomic DNA isolated from *E. faecalis* strain OG1RF was partially digested with DNase I and "blunt-ended" by incubating with T4 DNA polymerase. Random genomic fragments between 200 and 800 base pairs in length were selected by gel purification. The size-selected genomic fragments were added to the linearized and dephosphorylated vector
20 at a molar ratio of 2 to 1, and ligated to form a shotgun library.

The ligated products were transformed into electrocompetent *E. coli* strain TOP10 cells (Invitrogen) and plated on LB medium with erythromycin (Erm) at 150 µg/mL. Resulting colonies numbering 5×10^5 or greater were scraped and combined, and were then subjected to plasmid purification.

25 The purified library was then transformed into electrocompetent *E. faecalis* strain OG1RF. Resulting transformants were plated on Todd-Hewitt (TH) agar with erythromycin at 10 µg/mL in order to generate 100 to 150 platings at 500 colonies per plating. The colonies were subjected to robotic picking and arrayed into wells of 384 well culture dishes. Each well contained 100 µL of THB + Erm 10 µg/mL.
30 Inoculated 384 well dishes were incubated 16 hours at room temperature, and each

well was robotically gridded onto solid TH agar + Erm with or without 5% xylose. Gridded plates were incubated 16 hours at 37 °C, and then manually scored for arrayed colonies that were growth-compromised in the presence of xylose.

Arrayed colonies that were growth-sensitive on medium containing 5% xylose, yet were able to grow on similar medium lacking xylose, were subjected to further growth sensitivity analysis. Colonies from the plate lacking xylose were manually picked and inoculated into individual wells of a 96 well culture dish containing THB + Erm 10, and were incubated for 16 hours at 30 °C. These cultures were robotically diluted 1/100 into fresh medium and allowed to incubate for 4 hours at 37 °C, after which they were subjected to serial dilution on plates containing 5% xylose or plates lacking xylose. After growth for 16 hours at 37 °C, the arrays of serially diluted spots that resulted were compared between the two media. Colonies that grew similarly on both media were scored as a negative and corresponding colonies were no longer considered. Colonies on xylose medium that failed to grow to the same serial dilution compared to those on the non-xylose plate were given a score based on the differential. For example, colonies on xylose medium that only grow to a serial dilution of -4 while they were able to grow to -8 on the non-xylose plate, then the corresponding transformant colony received a score of "4" representing the log difference in growth observed.

Of the 1.4 million *Enterococcus faecalis* colonies screened using the CP25 fusion promoter, about 1 of every 3750 colonies had reduced proliferation ability whereas of the 390,000 colonies screened using the P59 fusion promoter, about 1 of every 390 colonies had reduced proliferation ability.

Thus, it will be appreciated that use of a regulatable fusion promoter from the suite of regulatable fusion promoters which has a higher level of transcriptional activity may increase the frequency at which proliferation-required genes are identified and may allow identification of proliferation-required genes which would not be identified using weaker promoters. Such a fusion promoter would be expected to produce a larger number of colonies having reduced proliferation ability per

colonies screened. Furthermore, proliferation-required genes from new gene classes would be expected.

Expression vectors were purified from *Staphylococcus aureus* or *Enterococcus faecalis* colonies identified as having reduced proliferation upon induction of the fusion promoter. The nucleic acids contained on the expression vectors under control of the fusion promoter were isolated for subsequent nucleotide sequence determination and further characterization.

It will be appreciated that the random genomic fragments used in the above methods can be generated using restriction enzymes and endonucleases other than those specifically described or by any other procedure. In addition, random genomic fragments may be generated by mechanical shearing. Sonication and nebulization are two such techniques commonly used for mechanical shearing of DNA.

The next example describes experiments that use the Rapid Replacement of Genomic Promoters technique to identify essential genes for use as targets for drug discovery.

EXAMPLE 9

Replacement of Genomic Promoters for Proliferation-Required Genes with Inducible Fusion Promoters

Nucleic acids involved in proliferation of *Staphylococcus aureus* or *Enterococcus faecalis* are identified using the Rapid Replacement of Genomic Promoters technique described in U.S. Patent Application Serial Number 09/948993, filed September 6, 2001, the disclosure of which is incorporated herein by reference in its entirety, and U.S. Provisional Patent Application Serial Number 60/230347, filed September 6, 2000, the disclosure of which is incorporated herein by reference in its entirety. Initially, a target gene is selected. The gene may be a gene which is known or suspected to be essential for proliferation or a gene which has not been characterized with respect to its essentiality for proliferation. Identification of the essential gene can be accomplished by using the EGI technique or other methods including, but not limited to, genes shown to be essential in the literature or genes of

unknown essentiality that are predicted to be essential by bioinformatics. Additionally, one can target essential operons containing multiple genes.

Next, the native promoter of the target gene or operon is replaced with a regulatable promoter, such as the fusion promoters described herein using "ET cloning" technology. Briefly, this is accomplished by engineering the one of the fusion promoters described herein such that it is 3' of a selectable marker (e.g., a CAT gene). This entire cassette is amplified by PCR. This PCR product becomes the template for another round of PCR using primers with 80bp of homology to the target promoter and 20bp of homology to the fusion promoter-containing cassette. However, as little as 60bp homology to the target promoter can be used. The region of homology is chosen such that the cassette will replace the promoter of the target gene without effecting the endogenous Shine-Delgarno sequence. This construct is then transformed into competent cells containing various levels of inducer (e.g. xylose or analog thereof) to assure that the correct level of expression is achieved to allow survival. In some embodiments, the native promoter in a strain having an enhanced frequency of homologous recombination may be replaced with a regulatable promoter.

The correct integration of the cassette can be confirmed by colony PCR as described in U.S. Patent Application Serial Number 09/948993, filed September 6, 2001, and U.S. Provisional Patent Application entitled Rapid PCR Method for Determination of Whether a Gene is Essential, Serial Number 60/230347, filed September 6, 2000, attorney docket number ELITRA.022PR, the disclosures of which are incorporated herein by reference in their entireties. Preferably, correct integrants are tested for growth defects as inducer is titrated away. The inability to grow in the absence of inducer will confirm that the target was essential. These strains can then be induced to a level determined to be the optimal balance between slowed growth and sensitivity to inhibition of the target protein for use in a cell-based assay for compounds with antibiotic activity or may be used to identify genes required for proliferation as described herein.

The next example describes experiments that use an operator insertion technique to identify essential genes for use as targets for drug discovery.

EXAMPLE 10**Regulation of Genomic Promoters for Proliferation-Required Genes by Operator****Insertion**

5 In this example, insertion of an operator into the promoter of a proliferation-required gene is described. A target promoter which drives the expression of a target gene or operon is identified. The target gene may be a gene which is known to be required for proliferation, suspected to be required for proliferation, or a gene which has not been characterized with respect to whether it is required for proliferation. An
10 oligonucleotide comprising a *xyl*, *tet*, *trp*, *mal*, *λcl* or *lac* operator flanked on each side by 40 nucleotides homologous to the target promoter is synthesized. The 40 nucleotide flanking sequences are determined based on the desired location for intergration of the *xyl* operator into the promoter. The single stranded oligonucleotide construct is then transformed into a bacterium having an enhanced frequency of
15 homologous recombination. For example, the bacterium may express the λ Beta and Gam proteins. The cells in the transformation mixture are diluted and plated on medium containing xylose. Colonies in which the operator has integrated into the target promoter are identified by colony PCR. The identified colonies are grown in medium containing or lacking inducer. The colonies proliferate on medium
20 containing inducer but fail to grow on medium lacking inducer, thereby indicating the target promoter directs transcription of a gene encoding a gene product required for proliferation. Strains in which a gene encoding a gene product required for proliferation is under the control of the *xyl*, *tet*, *trp*, *mal*, *λcl* or *lac* operator may also be used to identify compounds which inhibit proliferation in the cell-based assays
25 described herein.

It will be appreciated that operators other than the *xyl*, *tet*, *trp*, *mal*, *λcl* or *lac* operator can be used with the methods described above.

The next example describes experiments that use the Random Insertional Cis-Antisense technique to identify essential genes for use as targets for drug discovery.

30

EXAMPLE 11**Insertional Inactivation of Proliferation-Required Genes by Random Insertion of an Inducible Fusion Promoter**

Nucleic acids involved in proliferation of *Staphylococcus* or *Enterococcus* are
5 identified using the Random Insertional Cis-Antisense technique described in U.S.
Provisional Patent Application Serial Number 60/230,403, filed September 6, 2000,
the disclosure of which is incorporated herein by reference in its entirety. In one
embodiment, a vector containing an outward-directed inducible fusion promoter is
randomly integrated into the genome of a Gram-positive organism such as *S. aureus* or
10 *E. faecalis* or the other Gram-positive organisms described herein and clones
containing promoters in cis-antisense orientation with regard to essential genes are
identified. Accordingly, a vector containing an inducible promoter, such as one of the
fusion promoters described herein, is first engineered such that the inducible promoter
reads into a multiple cloning site (MCS). The genetic marker is juxtaposed directly 3'
15 of the MCS. This positioning helps ensure that the transforming vector will not be
degraded by exonucleases (transformants in which the vector is degraded will not be
recovered because the genetic marker will be lost). The vector is linearized by cutting
with any single restriction enzyme in the MCS. Using different enzymes to linearize
may increase the randomness of integration of the vector. The linearized vector is
20 then transformed into the appropriate bacterial strain.

The integrative transformation frequency of linear DNA varies widely among
bacterial strains. In some strains, linear vectors usually transform integratively at a
significantly higher frequency than circular vectors. This is especially true in bacterial
strains that have been modified to have an enhanced frequency of homologous
25 recombination. In strains in which the frequency of integrative transformation of
linear vectors is low, circular vectors often transform integratively at a higher
frequency. The mechanism of integration is largely unknown but it is contemplated
that integration occurs via nonhomologous recombination. Sites of integration appear
to be entirely random. Transformants are robotically picked, grown, and are
30 robotically gridded onto inducing and noninducing (inducing or not inducing

transcription of the promoter) media. These gridded plates are then screened for transformants that have a pronounced growth defect on inducing media relative to the phenotype on noninducing media. Such transformants may have the vector integrated into their genome in several possible locations in an essential gene: the 3' untranslated
5 region or any part of the coding or noncoding region that does not result in disruption of function of the gene product. In most cases the inducible promoter is oriented such that it produces an antisense RNA from the target gene. Thus, when the promoter is induced the expression of the essential gene is reduced. Such inhibition can occur by one of several mechanisms including binding of the antisense RNA to the mRNA,
10 colliding RNA polymerases, RNA interference (RNAi) or a combination of such phenomenon.

The identity of the essential gene is then determined by sequencing the junction of the vector with the essential gene, the junction where the inducible promoter is juxtaposed with genomic DNA. This assay can also be used to create
15 bacterial strains for cell-based assays that evaluate candidate antibiotic compounds as described herein.

The next example describes approaches that were used to determine the identity of the newly discovered essential genes.

20

EXAMPLE 12

Nucleotide Sequence Determination of Genes that are Affected by Proliferation- Inhibiting Antisense Nucleic Acids Expressed from Fusion Promoters

With reference to experiments that were performed using the antisense technique, this example describes approaches that were used to determine the nucleic
25 acid sequence of newly discovered essential genes. It should be understood that the teachings herein can also be used to determine the nucleic acid sequence of essential genes identified using the promoter replacement, operator insertion or Random Insertional Cis-Antisense technique.

Plasmids from transformant colonies that received a dilution plating score of
30 "2" or greater were isolated to obtain the genomic DNA insert responsible for growth

inhibition as follows. *Staphylococcus aureus* were grown in standard laboratory media (LB or TB with 15 ug/ml Chloramphenicol to select for the plasmid). Growth was carried out at 37°C overnight in culture tubes or 2 ml deep well microtiter plates. Lysis was performed as follows. Cultures (2-5 ml) were centrifuged and the cell pellets resuspended in 1.5 mg/ml solution of lysostaphin (20 µl/ml of original culture) followed by addition of 250 µl of resuspension buffer (Qiagen). Alternatively, cell pellets were resuspended directly in 250 µl of resuspension buffer (Qiagen) to which 5-20 µl of a 1 mg/ml lysostaphin solution were added. DNA was isolated using Qiagen miniprep kits or Wizard (Qiagen) miniprep kits according to the instructions provided by the manufacturer.

The genomic DNA inserts were then amplified from the purified plasmids by PCR as follows. Approximately, 1 :1 of Qiagen purified plasmid was put into a total reaction volume of 25 :1 Qiagen Hot Start PCR mix. The following primers were used in the PCR reaction:

pXylT5F: CAGCAGTCTGAGTTATAAAATAG (SEQ ID NO: 24)

LexL TGT TTTATCAGACCGCTT (SEQ ID NO: 25)

PCR was carried out in a PE GenAmp with the following cycle times:

- Step 1. 95 °C 15 min
- Step 2. 94 °C 45 sec
- Step 3. 54 °C 45 sec
- Step 4. 72 °C 1 minute
- Step 5. Return to step 2 (29 times)
- Step 6. 72 °C 10 minutes
- Step 7. 4 °C hold

The PCR products were cleaned using Qiagen Qiaquick PCR plates according to the manufacturer's instructions. The amplified genomic DNA inserts were then subjected to automated sequencing.

For *E. faecalis*, plasmids from transformant colonies that received a dilution plating score of "2" or greater were isolated to obtain the genomic DNA insert responsible for growth inhibition as follows. *E. faecalis* were grown in THB 10 µg/ml Erm at 30°C overnight in 100 µl culture wells in microtiter plates. To amplify insert
5 DNA 2 µl of culture were placed into 25 µl Qiagen Hot Start PCR mix. PCR reactions were in 96 well microtiter plates. Primers to vector regions flanking the insert were used in the PCR reaction. PCR was carried out in a PE GenAmp with the following cycle times:

- 10 Step 1. 95° C 15 min
- Step 2. 94° C 45 sec
- Step 3. 54° C 45 sec
- Step 4. 72° C 1 minute
- Step 5. Return to step 2, 29 times
- 15 Step 6. 72° C 10 minutes
- Step 7. 4° C hold

The PCR products were cleaned using Qiagen Qiaquick PCR plates according to the manufacturer's instructions. The purified PCR products were then directly
20 cycle sequenced with Qiagen Hot Start PCR mix. PCR was carried out in a PE GenAmp with the following cycle times:

- Step 1. 94° C 15 min
- Step 2. 96° C 10 sec
- 25 Step 3. 50° C 5 sec
- Step 4. 60° C 4 min
- Step 5. Return to step 2, 24 times
- Step 6. 4° C hold

The PCR products were cleaned using Qiagen Qiaquick PCR plates according to the manufacturer's instructions. The amplified genomic DNA inserts were then subjected to automated sequencing.

5 The example below describes techniques that were used to compare the sequences identified by the approaches above to sequences present in publicly available databases so as to better characterize the essential genes.

EXAMPLE 13

Comparison Of Isolated Nucleic Acids to Known Sequences

10 With reference to experiments that were performed using the antisense technique, this example describes approaches that were used to characterize the nucleic acid sequence of newly discovered essential genes in *S. aureus* and *E. faecalis*. It should be understood that the teachings herein can also be used to determine the nucleic acid sequence of essential genes in other Gram-positive cells and essential genes identified in
15 Gram-positive cells using the promoter replacement, operator insertion or Random Insertional Cis-Antisense technique.

The nucleic acid sequences of the subcloned fragments from *S. aureus* or *E. faecalis* obtained from the expression vectors containing the Xyl-T5 and CP25 fusion promoters discussed above were compared to known sequences from *S. aureus* or *E.*
20 *faecalis* and other microorganisms as follows. First, to confirm that each clone originated from one location on the chromosome and was not chimeric, the sequences of the selected clones were compared against the *S. aureus* or *E. faecalis* genomic sequences to align the clone to the correct position on the chromosome. The NCBI BLASTN v 2.0.9 program was used for this comparison, and the incomplete
25 *Staphylococcus aureus* genomic sequences licensed from TIGR, as well as the NCBI nonredundant GenBank database were used as the source of genomic data. The *E. faecalis* sequences were compared to a proprietary database.

The BLASTN analysis was performed using the default parameters except that the filtering was turned off. No further analysis was performed on inserts which
30 resulted from the ligation of multiple fragments.

In general, antisense molecules and their complementary genes are identified as follows. First, all possible full length open reading frames (ORFs) are extracted from available genomic databases. Such databases include the GenBank nonredundant (nr) database, the unfinished genome database available from TIGR and the PathoSeq database developed by Incyte Genomics. The latter database comprises over 40 annotated bacterial genomes including complete ORF analysis. If databases are incomplete with regard to the bacterial genome of interest, it is not necessary to extract all ORFs in the genome but only to extract the ORFs within the portions of the available genomic sequences which are complementary to the clones of interest.

Computer algorithms for identifying ORFs, such as GeneMark, are available and well known to those in the art. Comparison of the clone DNA to the complementary ORF(s) allows determination of whether the clone is a sense or antisense clone. Furthermore, each ORF extracted from the database can be compared to sequences in well annotated databases including the GenBank (nr) protein database, SWISSPROT and the like. A description of the gene or of a closely related gene in a closely related microorganism is often available in these databases. Similar methods are used to identify antisense clones corresponding to genes encoding non-translated RNAs.

In some cases, for example, the NCBI BLASTN 2.0.9 computer algorithm was used and the default parameters were used with the exception that filtering was turned off. The default parameters for the BLASTN and BLASTX analyses were:

Expectation value (e)=10
 Alignment view options: pairwise
 Filter query sequence (DUST with BLASTN, SEG with others)=T
 Cost to open a gap (zero invokes behavior)=0
 Cost to extend a gap (zero invokes behavior)=0
 X dropoff value for gapped alignment (in bits) (zero invokes behavior)=0
 Show GI's in defines=F
 Penalty for a nucleotide mismatch (BLASTN only)=!3
 Reward for a nucleotide match (BLASTN only)=1
 Number of one-line descriptions (V)=500
 Number of alignments to show (B)=250
 Threshold for extending hits=default
 Perform gapped alignment (not available with BLASTX)=T
 Query Genetic code to use=1
 DB Genetic code (for TBLAST[nx] only)=1

Number of processors to use=1
 SeqAlign file
 Believe the query defline=F
 Matrix=BLOSUM62
 5 Word Size= default
 Effective length of the database (use zero for the real size)=0
 Number of best hits from a region to keep=100
 Length of region used to judge hits=20
 Effective length of the search space (use zero for the real size)=0
 10 Query strands to search against database (for BLAST[nx] and TBLASTX), 3 is
 both, 1 is top, 2 is bottom=3
 Produce HTML output=F

Those ORFs found to be complementary to the cloned nucleic acid sequences
 discussed above were evaluated with regard to orientation. Antisense clones were
 15 identified as those clones for which transcription from the inducible promoter would
 result in the expression of an RNA antisense to the complementary ORF.

The gene descriptions in the PathoSeq database derive from annotations
 available in the public sequence databases described above. It will be appreciated that
 ORFs may also be identified using databases other than PathoSeq. For example, the
 20 ORFs may be identified using the methods described in U.S. Provisional Patent
 Application Serial Number 60/191,078, filed March 21, 2000, the disclosure of which
 is incorporated herein by reference in its entirety.

In the antisense-based screening experiments in *Enterococcus faecalis* as
 described in Examples 8 and 9, 256 proliferation-inhibiting antisense sequences
 25 corresponding to 61 unique ORFs were discovered using the CP25 fusion promoter
 whereas 790 proliferation-inhibiting antisense sequences corresponding to 150 unique
 ORFs were discovered using the P59 fusion promoter.

The example below describes approaches that may be used to identify operons or
 downstream genes associated with the essential genes identified by the methods above.
 30

EXAMPLE 14

Identification of Genes and their Corresponding Operons that are Affected by
Proliferation-Inhibiting Antisense Nucleic Acids Expressed from Fusion Promoters

With reference to experiments that were performed using the antisense technique, this example describes approaches that may be used to identify operons or downstream genes associated with the newly discovered essential genes in *S. aureus* and *E. faecalis*. It should be understood that the teachings herein can also be used to identify operons or downstream genes associated with the essential genes in other Gram-positive cells and essential genes identified in Gram-positive cells using the promoter replacement, operator insertion or Random Insertional Cis-Antisense technique

Once the genes involved in *S. aureus* and *E. faecalis* proliferation are identified as described above, the operons in which these genes lie may be identified by comparison with known microbial genomes. Since bacterial genes are expressed in a polycistronic manner, the antisense inhibition of a single gene in an operon might affect the expression of all the other genes on the operon or the genes downstream from the single gene identified. Accordingly, each of the genes contained within an operon may be analyzed for their effect on proliferation.

Operons are predicted by looking for all adjacent genes in a genomic region that lie in the same orientation with no large noncoding gaps in between. First, full length ORFs complementary to the antisense molecules are identified as described above. Adjacent ORFs are then identified and their relative orientation determined either by directly analyzing the genomic sequences surrounding the ORFs complementary to the antisense clones or by extracting adjacent ORFs from the collection obtained through whole genome ORF analysis described above followed by ORF alignment. Operons predicted in this way may be confirmed by comparison to the arrangement of the homologous genes in the *Bacillus subtilis* complete genome sequence, as reported by the genome database compiled at Institut Pasteur Subtilist Release R15.1 (June 24, 1999) which can be found at <http://bioweb.pasteur.fr/GenoList/SubtiList/>. The *Bacillus subtilis* genome is the only fully sequenced and annotated genome from a Gram-positive microorganism, and appears to have a high level of similarity to *Staphylococcus aureus* both at the level of conservation of gene sequence and genomic organization including operon structure. Annotation of some of the DNA sequences in some of the aforementioned

databases is lacking, but comparisons may be made to *E. coli* using tools such as BLASTX. Public or proprietary databases may be used to analyzed *E. faecalis* sequences as well as the databases listed above.

5 Once the full length ORFs and/or the operons containing them have been identified using the methods described above, they can be obtained from a genomic library by performing a PCR amplification using primers at each end of the desired sequence. Those skilled in the art will appreciate that a comparison of the ORFs to homologous sequences in other microorganisms will facilitate confirmation of the start and stop codons at the ends of the ORFs.

10 In some embodiments, the primers may contain restriction sites which facilitate the insertion of the gene or operon into a desired vector. For example, the gene may be inserted into an expression vector and used to express the proliferation-required protein as described below. Other methods for obtaining the full length ORFs and/or operons are familiar to those skilled in the art. For example, natural
15 restriction sites may be employed to insert the full length ORFs and/or operons into a desired vector.

The next example describes approaches that may be used to identify individual genes within an operon that was characterized as being required for proliferation.

20

EXAMPLE 15

Identification of Individual Genes within an Operon Required for Proliferation

With reference to experiments that were performed using the antisense technique, this example describes approaches that may be used to identify individual genes within an operon that has been characterized as being required for proliferation
25 of a Gram-positive organism such as *S. aureus* or *E. faecalis* or the other Gram-positive organisms listed herein. It should be understood that the teachings herein can also be used to identify individual genes within an operon that were characterized as being required for proliferation in other Gram-positive cells and can be used in conjunction with the promoter replacement, operator insertion or Random Insertional Cis-Antisense
30 technique.

The strategy described in this example seeks to determine if a targeted gene within an operon is required for cell proliferation by replacing the targeted gene in the chromosome with an in-frame deletion of the coding region of the targeted gene. Deletion inactivation of a chromosomal copy of a gene in *Staphylococcus aureus* or
5 *Enterococcus faecalis* can be accomplished by integrative gene replacement. The principles of this method were described in Xia, M., et al. 1999 Plasmid 42:144-149 and Hamilton, C. M., et al 1989. *J. Bacteriol.* 171: 4617-4622, the disclosures of which are incorporated herein by reference in their entireties. In this approach, a mutant allele of the targeted gene is constructed by way of an in-frame deletion and
10 introduced into the chromosome using a suicide vector. This results in a tandem duplication comprising a deleted (null) allele and a wild type allele of the target gene. Cells in which the vector sequences have been deleted are isolated using a counter-selection technique. Removal of the vector sequence from the chromosomal insertion results in either restoration of the wild-type target sequence or replacement of the wild
15 type sequence with the deletion (null) allele. *E. faecalis* genes can be disrupted using a suicide vector that contains an internal fragment to a gene of interest. With the appropriate selection this plasmid will homologously recombine into the chromosome (Nallapareddy, S. R., X. Qin, G. M. Weinstock, M. Hook, B. E. Murray. 2000. *Infect. Immun.* 68:5218-5224, the disclosure of which is incorporated herein by reference).

20 The resultant population of *Staphylococcus aureus*, *Enterococcus faecalis*, or other Gram-positive colonies can then be evaluated to determine whether the target sequence is required for proliferation by PCR amplification of the affected target sequence. If the targeted gene is not required for proliferation, then PCR analysis will show that roughly equal numbers of colonies have retained either the wild-type or the
25 mutant allele. If the targeted gene is required for proliferation, then only wild-type alleles will be recovered in the PCR analysis.

The method of cross-over PCR is used to generate the mutant allele by amplification of sequences flanking but not including the coding region of the gene of interest, such that overlap between the resulting two PCR amplification products
30 allows them to hybridize. Further PCR amplification of this hybridization product

using primers representing the extreme 5' and 3' ends can produce an amplification product containing an in-frame deletion of the coding region but retaining substantial flanking sequences.

For *Staphylococcus aureus*, this amplification product is subcloned into the suicide vector pSA3182 (Xia, M., *et al.* 1999 *Plasmid* 42:144-149, the disclosure of which is incorporated herein by reference in its entirety) which is host-dependent for autonomous replication. This vector includes a *tetC* tetracycline-resistance marker and the origin of replication of the well-known *Staphylococcus aureus* plasmid pT181 (Mojumdar, M and Kahn, S.A., Characterisation of the Tetracycline Resistance Gene of Plasmid pT181, *J. Bacteriol.* 170: 5522 (1988), the disclosure of which is incorporated herein by reference in its entirety). The vector lacks the *repC* gene which is required for autonomous replication of the vector at the pT181 origin. However autonomous replication of this vector can occur in a *Staphylococcus aureus* host strain such as SA3528, which expresses *repC* in trans. Once the amplified truncated target gene sequence is cloned and propagated in the pSA3182 vector, it can then be introduced into a *repC* minus strain such as RN4220 (Kreiswirth, B.N. *et al.*, The Toxic Shock Syndrome Exotoxin Structural Gene is Not Detectably Transmitted by a Prophage, *Nature* 305:709-712 (1983), the disclosure of which is incorporated herein by reference in its entirety) by electroporation with selection for tetracycline resistance. In this strain, the vector must integrate by homologous recombination at the targeted gene in the chromosome to impart drug resistance. This results in a inserted truncated copy of the allele, followed by pSA3182 vector sequence, and finally an intact and functional allele of the targeted gene.

Once a *tetR Staphylococcus aureus* strain is isolated from the above technique and shown to include truncated and wild-type alleles of the targeted gene as described above, a second plasmid, pSA7592 (Xia, M., *et al.* 1999 *Plasmid* 42:144-149, the disclosure of which is incorporated herein by reference in its entirety) is introduced into the strain by electroporation. This gene includes an erythromycin resistance gene and a *repC* gene that is expressed at high levels. Expression of *repC* in these transformants is toxic due to interference of normal chromosomal replication at the

integrated pT181 origin of replication. This counterselects for strains that have removed the vector sequence by homologous recombination, resulting in either of two outcomes: The counterselected cells either possess a wild-type allele of the targeted gene or a gene in which the wild-type allele has been replaced by the engineered in-frame deletion of the truncated allele.

PCR amplification can be used to test either outcome among the resulting erythromycin resistant, tet sensitive transformant colonies. If the targeted gene is not required for cellular replication, then PCR evidence for both wild-type and mutant alleles will be found among the population of resultant transformants. However, if the targeted gene is required for cellular replication, then only the wild-type form of the gene will be evident among the resulting transformants.

Similarly, for *Enterococcus faecalis* the PCR products containing the mutant allele of the target sequence may be introduced into an appropriate knockout vector and cells in which the wild type target has been disrupted are selected using the appropriate methodology.

The above methods have the advantage that insertion of an in-frame deletion mutation is far less likely to cause downstream polar effects on genes in the same operon as the targeted gene. However, it will be appreciated that other methods for disrupting *Staphylococcus aureus* or *Enterococcus faecalis*, which are familiar to those skilled in the art may also be used. Each gene in the operon may be disrupted using the methodology above to determine whether it is required for proliferation.

The example below describes approaches to express genes involved in cell proliferation.

25

EXAMPLE 16

Expression of the Proteins Encoded by Genes Identified as Required for *Staphylococcus aureus* or *Enterococcus faecalis* Proliferation

The following is provided as one exemplary method to express the proliferation-required proteins encoded by sequences identified by the methods described herein using expression systems designed either for *E. coli*, *Staphylococcus aureus*, or

30

Enterococcus faecalis. First, the initiation and termination codons for the gene are identified. If desired, methods for improving translation or expression of the protein are well known in the art. For example, if the nucleic acid encoding the polypeptide to be expressed lacks a methionine codon to serve as the initiation site, a strong Shine-Dalgarno sequence, or a stop codon, these sequences can be added. Similarly, if the identified nucleic acid sequence lacks a transcription termination signal, this sequence can be added to the construct by, for example, splicing out such a sequence from an appropriate donor sequence. In addition, the coding sequence may be operably linked to a strong promoter or an inducible promoter, such as the fusion promoters described herein, if desired. The identified nucleic acid sequence or portion thereof encoding the polypeptide to be expressed is obtained by, for example, PCR from the bacterial expression vector or genome using oligonucleotide primers complementary to the identified nucleic acid sequence or portion thereof and containing restriction endonuclease sequences appropriate for inserting the coding sequences into the vector such that the coding sequences can be expressed from the vector's promoter. Alternatively, other conventional cloning techniques may be used to place the coding sequence under the control of the promoter. In some embodiments, a termination signal may be located downstream of the coding sequence such that transcription of the coding sequence ends at an appropriate position.

Several expression vector systems for protein expression in *E. coli* are well known and available to someone knowledgeable in the art. The coding sequence may be inserted into any of these vectors and placed under the control of the promoter. The expression vector may then be transformed into DH5 α or some other *E. coli* strain suitable for the over expression of proteins.

Alternatively, an expression vector encoding a protein required for proliferation of *Staphylococcus aureus* or *Enterococcus faecalis* may be introduced into *Staphylococcus aureus* or *Enterococcus faecalis*. In some embodiments, gene encoding the protein required for proliferation may be operably linked to one of the fusion promoters described herein. Electroporation protocols for introducing nucleic acids into *Staphylococcus aureus* and *Enterococcus faecalis* are well known in the art.

For example, the electroporation protocol described in *Staphylococcus aureus* transformation protocols described in J.C.Lee "Electroporation of Staphylococci" from *Methods in Molecular Biology* vol 47: Electroporation Protocols for Microorganisms Edited by : J.A. Nickoloff Humana Press Inc., Totowa, NJ. pp209-216, the disclosure
5 of which is incorporated herein by reference in its entirety, may be used. Positive transformants are selected after growing the transformed cells on plates containing an antibiotic to which the vector confers resistance. For example, *Staphylococcus aureus* may be transformed with an expression vector in which the coding sequence is operably linked to the fusion promoters described herein, such as fusion promoters comprising the
10 T5 promoter operably linked to a *xyl* operator such that expression of the encoded protein is inducible with xylose. Such an example vector is pXyl-T5 p15a. In another example, *Enterococcus faecalis* may be transformed with an expression vector in which the coding sequence is operably linked to the fusion promoters described herein, such as fusion promoters comprising the PL promoter operably linked to a *xyl* operator
15 such that expression of the encoded protein is inducible with xylose. Such an example vector is pEPEF21.

In one embodiment, the protein is expressed and maintained in the cytoplasm as the native sequence. In an alternate embodiment, the expressed protein can be modified to include a protein tag that allows for differential cellular targeting, such as
20 to the periplasmic space of Gram-negative or Gram-positive expression hosts or to the exterior of the cell (i.e., into the culture medium). In some embodiments, the osmotic shock cell lysis method described in Chapter 16 of *Current Protocols in Molecular Biology*, Vol. 2, (Ausubel, *et al.*, Eds.) John Wiley & Sons, Inc. (1997) may be used to liberate the polypeptide from the cell. In still another embodiment, such a protein tag
25 could also facilitate purification of the protein from either fractionated cells or from the culture medium by affinity chromatography. Each of these procedures can be used to express a proliferation-required protein.

Expressed proteins, whether in the culture medium or liberated from the periplasmic space or the cytoplasm, are then purified or enriched from the supernatant
30 using conventional techniques such as ammonium sulfate precipitation, standard

chromatography, immunoprecipitation, immunochromatography, size exclusion chromatography, ion exchange chromatography, and HPLC. Alternatively, the polypeptide may be secreted from the host cell in a sufficiently enriched or pure state in the supernatant or growth media of the host cell to permit it to be used for its intended purpose without further enrichment. The purity of the protein product obtained can be assessed using techniques such as SDS PAGE, which is a protein resolving technique well known to those skilled in the art. Coomassie, silver staining or staining with an antibody are typical methods used to visualize the protein of interest.

Antibodies capable of specifically recognizing the protein of interest can be generated using synthetic peptides using methods well known in the art. See, *Antibodies: A Laboratory Manual*, (Harlow and Lane, Eds.) Cold Spring Harbor Laboratory (1988). For example, 15-mer peptides having a sequence encoded by the appropriate identified gene sequence of interest or portion thereof can be chemically synthesized. The synthetic peptides are injected into mice to generate antibodies to the polypeptide encoded by the identified nucleic acid sequence of interest or portion thereof. Alternatively, samples of the protein expressed from the expression vectors discussed above can be purified and subjected to amino acid sequencing analysis to confirm the identity of the recombinantly expressed protein and subsequently used to raise antibodies.

The protein encoded by the identified nucleic acid sequence of interest or portion thereof can be purified using standard immunochromatography techniques. In such procedures, a solution containing the desired protein, such as the culture medium or a cell extract, is applied to a column having antibodies against the secreted protein attached to the chromatography matrix. The desired protein is allowed to bind the immunochromatography column. Thereafter, the column is washed to remove non-specifically bound proteins. The specifically-bound secreted protein is then released from the column and recovered using standard techniques. These procedures are well known in the art.

In an alternative protein purification scheme, the identified nucleic acid sequence of interest or portion thereof can be incorporated into expression vectors designed for

use in purification schemes employing chimeric polypeptides. In such strategies the coding sequence of the identified nucleic acid sequence of interest or portion thereof is inserted in-frame with the gene encoding the other half of the chimera. The other half of the chimera can be maltose binding protein (MBP) or a nickel binding polypeptide encoding sequence. A chromatography matrix having antibody to MBP or nickel attached thereto is then used to purify the chimeric protein. Protease cleavage sites can be engineered between the MBP gene or the nickel binding polypeptide and the identified expected gene of interest, or portion thereof. Thus, the two polypeptides of the chimera can be separated from one another by protease digestion.

One useful expression vector for generating maltose binding protein fusion proteins is pMAL (New England Biolabs), which encodes the *malE* gene. In the pMal protein fusion system, the cloned gene is inserted into a pMal vector downstream from the *malE* gene. This results in the expression of an MBP-fusion protein. The fusion protein is purified by affinity chromatography. These techniques as described are well known to those skilled in the art of molecular biology. An Example describing in detail the generation of monoclonal and polyclonal antibodies appears below.

EXAMPLE 17

Production of an Antibody to an isolated *Staphylococcus aureus* or *Enterococcus faecalis* Protein

Substantially pure protein or polypeptide is isolated from the transformed cells as described in Example 16. The concentration of protein in the final preparation is adjusted, for example, by concentration on a 10,000 molecular weight cut off AMICON filter device (Millipore, Bedford, MA), to the level of a few micrograms/ml.

Monoclonal or polyclonal antibody to the protein can then be prepared as follows:

Monoclonal antibody to epitopes of any of the peptides identified and isolated as described can be prepared from murine hybridomas according to the classical method of Kohler, G. and Milstein, C., *Nature* 256:495 (1975) or any of the well-known derivative methods thereof. Briefly, a mouse is repetitively inoculated with a few micrograms of the selected protein or peptides derived therefrom over a period of a few weeks. The

mouse is then sacrificed, and the antibody-producing cells of the spleen isolated. The spleen cells are fused by means of polyethylene glycol with mouse myeloma cells, and the excess unfused cells destroyed by growth of the system on selective media comprising aminopterin (HAT media). The successfully-fused cells are diluted and
5 aliquots of the dilution placed in wells of a microtiter plate where growth of the culture is continued. Antibody-producing clones are identified by detection of antibody in the supernatant fluid of the wells by immunoassay procedures, such as ELISA, as described by Engvall, E., "Enzyme immunoassay ELISA and EMIT," *Meth. Enzymol.* 70:419 (1980), and derivative methods thereof. Selected positive clones can be expanded and
10 their monoclonal antibody product harvested for use. Detailed procedures for monoclonal antibody production are described in Davis, L. *et al. Basic Methods in Molecular Biology* Elsevier, New York. Section 21-2.

Polyclonal antiserum containing antibodies to heterogeneous epitopes of a single protein or a peptide can be prepared by immunizing suitable animals with the expressed
15 protein or peptides derived therefrom described above, which can be unmodified or modified to enhance immunogenicity. Effective polyclonal antibody production is affected by many factors related both to the antigen and the host species. For example, small molecules tend to be less immunogenic than larger molecules and can require the use of carriers and adjuvant. Also, host animals vary in response to site of inoculations
20 and dose, with both inadequate or excessive doses of antigen resulting in low titer antisera. Small doses (ng level) of antigen administered at multiple intradermal sites appears to be most reliable. An effective immunization protocol for rabbits can be found in Vaitukaitis, J. *et al. J. Clin. Endocrinol. Metab.* 33:988-991 (1971).

Booster injections can be given at regular intervals, and antiserum harvested
25 when antibody titer thereof, as determined semi-quantitatively, for example, by double immunodiffusion in agar against known concentrations of the antigen, begins to fall. See, for example, Ouchterlony, O. *et al.*, Chap. 19 in: *Handbook of Experimental Immunology* D. Wier (ed) Blackwell (1973). Plateau concentration of antibody is usually in the range of 0.1 to 0.2 mg/ml of serum (about 12 :M). Affinity of the antisera
30 for the antigen is determined by preparing competitive binding curves, as described, for

example, by Fisher, D., Chap. 42 in: *Manual of Clinical Immunology*, 2d Ed. (Rose and Friedman, Eds.) Amer. Soc. For Microbiol., Washington, D.C. (1980).

Antibody preparations prepared according to either protocol are useful in quantitative immunoassays which determine concentrations of antigen-bearing substances in biological samples; they are also used semi-quantitatively or qualitatively to identify the presence of antigen in a biological sample. The antibodies can also be used in therapeutic compositions for killing bacterial cells expressing the protein. The next section describes how to make and use an *in vivo* system for performing *in vivo* analysis on bacterial strains containing regulatable essential genes and for screening compounds that inhibit bacterial proliferation.

The example below describes the an *in vivo* system for analyzing the growth characteristics of *S. aureus* and *Enterococcus faecalis* that contain regulatable essential genes.

15

EXAMPLE 18

In Vivo Models for Analyzing the Regulation of Essential Genes and Screening Therapeutic Compounds

Whether a given gene is essential for a bacterium growing in nature (*in vivo*) or whether a gene is required for infection of a host organism can be determined by constructing a conditional allele of that gene which is suitable for animal testing. Expression of an antisense RNA molecule complementary to a candidate gene can, by decreasing synthesis of the candidate gene product, constitute a conditional allele that is suitable for animal testing. Candidate genes essential *in vivo* or required for virulence can be tested by preparing bacterial strains using the EGI, antisense, promoter replacement, operator insertion, and random promoter insertion techniques. Preferably, the constructs used to prepare these bacterial strains contain an expression vector under control of a fusion promoter described herein. Animals infected by these bacterial strains can be used to evaluate the ability to modulate bacterial proliferation *in vivo* by adding or decreasing the concentration of inducer and can be used to screen compounds *in vivo* to validate their therapeutic properties.

Stable transformation of *Staphylococcus aureus* or *Enterococcus faecalis* with constructs prepared as described herein can be accomplished by electroporation and selection for clones resistant to 15 µg/ml chloramphenicol in the case of *Staphylococcus aureus* or 20 µg/ml erythromycin in the case of *Enterococcus faecalis*.

- 5 Preferably, whether each construct is essential is verified *in vitro* using induction in medium containing various amounts of inducer or an analog thereof, prior to introducing the cells into animals.

For the thigh infection model, animals are rendered neutropenic chemically prior to the start of the experiment. By one approach, the inoculum will consist of
10 10⁵ to 10⁶ cells of a normally virulent Gram-positive organism expressing antisense to an essential gene such as that for *gyrB* under control of a fusion promoter described herein. These cells are injected to one thigh of a suitable animal (e.g., mice). Most microorganisms attain a logarithmic growth 2 hours after inoculation. Treated animals receive therapy from 2 hours post inoculation up to 24 hours. A typical
15 efficacy study consists of 2 control groups and 5 treatment groups, each being treated with different doses of inducer. With 15 animals per group, 105 mice are used per study. The control sets are given intraperitoneal injections of saline, which will not induce expression of the fusion promoter. The experimental animals are given the inducer (e.g. xylose, tetracycline, IPTG, maltose, absence of tryptophan, or
20 temperature change) in intraperitoneal injections to induce expression of the antisense promoter. Alternatively, the promoter can be induced by intravenous infusion of inducer at sub-toxic levels.

The end point used to follow the infection process is viable bacterial counts per thigh. The controls in which the antisense RNA is not induced will fail to impede
25 growth of the Gram-positive organism and thus a logarithmic increase in viable bacteria will occur. The Gram-positive cells recovered from the site of infection should be viable until antisense expression is subsequently induced. This will demonstrate that the plasmid is still patent. However, in animals receiving the xylose injections, expression of the antisense RNA will occur, the essential gene or gene
30 product will be compromised, and the Gram-positive cells infecting the mice will not

multiply. Accordingly, fewer viable cells will be recovered from the site of infection in the experimental animals. The Gram-positive cells from the induced mice will be recovered, if still present, and assayed as above to determine if the promoter and gene are still present and functional.

5 The sections below describe the use of fusion promoters in methods to discover
compounds that inhibit bacterial proliferation.

EXAMPLE 19

Screening Chemical Libraries in Cells Sensitized with Antisense RNA Transcribed from Fusion Promoters

Having isolated and expressed bacterial proteins shown to be required for bacterial proliferation, the present invention further contemplates the use of these expressed target proteins in assays to screen libraries of compounds for potential drug candidates. The generation of chemical libraries is well known in the art. For example, 15 combinatorial chemistry can be used to generate a library of compounds to be screened in the assays described herein. A combinatorial chemical library is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis by combining a number of chemical "building block" reagents. For example, a linear combinatorial chemical library such as a polypeptide library is formed by 20 combining amino acids in every possible combination to yield peptides of a given length. Millions of chemical compounds theoretically can be synthesized through such combinatorial mixings of chemical building blocks. For example, one commentator observed that the systematic, combinatorial mixing of 100 interchangeable chemical building blocks results in the theoretical synthesis of 100 million tetrameric compounds or 10 billion pentameric compounds. (Gallop *et al.*, 1994, "Applications of 25 Combinatorial Technologies to Drug Discovery, Background and Peptide Combinatorial Libraries" *Journal of Medicinal Chemistry*, 37: 1233-1250). Other chemical libraries known to those in the art may also be used, including natural product libraries.

Once generated, combinatorial libraries can be screened for compounds that
30 possess desirable biological properties. For example, compounds which may be useful

as drugs or to develop drugs would likely have the ability to bind to the target protein identified, expressed and purified as discussed above. Further, if the identified target protein is an enzyme, candidate compounds would likely interfere with the enzymatic properties of the target protein. For example, the enzymatic function of a target protein
5 may be to serve as a protease, nuclease, phosphatase, dehydrogenase, transporter protein, transcriptional enzyme, and any other type of enzyme known or unknown. Thus, the present invention contemplates using the protein products described above to screen combinatorial chemical libraries.

In one example, the target protein is a serine protease and the substrate of the
10 enzyme is known. The present example is directed towards the analysis of libraries of compounds to identify compounds that function as inhibitors of the target enzyme. First, a library of small molecules is generated using methods of combinatorial library formation well known in the art. U.S. Patent Nos. 5,463,564 and 5,574, 656, to Agraftotis, *et al.*, entitled "System and Method of Automatically Generating Chemical
15 Compounds with Desired Properties," the disclosures of which are incorporated herein by reference in their entirety, are two such teachings. Then the library compounds are screened to identify those compounds that possess desired structural and functional properties. U.S. Patent No. 5,684,711, the disclosure of which is incorporated herein by reference in its entirety, also discusses a method for screening libraries.

20 To illustrate the screening process, the target polypeptide and chemical compounds of the library are combined with one another and permitted to interact with one another. A labeled substrate is added to the incubation. The label on the substrate is such that a detectable signal is emitted from metabolized substrate molecules. The emission of this signal permits one to measure the effect of the combinatorial library
25 compounds on the enzymatic activity of target enzymes by comparing it to the signal emitted in the absence of combinatorial library compounds. The characteristics of each library compound are encoded so that compounds demonstrating activity against the enzyme can be analyzed and features common to the various compounds identified can be isolated and combined into future iterations of libraries.

Once a library of compounds is screened, subsequent libraries are generated using those chemical building blocks that possess the features shown in the first round of screen to have activity against the target enzyme. Using this method, subsequent iterations of candidate compounds will possess more and more of those structural and functional features required to inhibit the function of the target enzyme, until a group of enzyme inhibitors with high specificity for the enzyme can be found. These compounds can then be further tested for their safety and efficacy as antibiotics for use in mammals.

It will be readily appreciated that this particular screening methodology is exemplary only. Other methods are well known to those skilled in the art. For example, a wide variety of screening techniques are known for a large number of naturally-occurring targets when the biochemical function of the target protein is known. For example, some techniques involve the generation and use of small peptides to probe and analyze target proteins both biochemically and genetically in order to identify and develop drug leads. Such techniques include the methods described in PCT publications No. WO9935494, WO9819162, WO9954728, the disclosures of which are incorporated herein by reference in their entireties.

Current cell-based assays used to identify or to characterize compounds for drug discovery and development frequently depend on detecting the ability of a test compound to modulate the activity of a target molecule located within a cell or located on the surface of a cell. Most often such target molecules are proteins such as enzymes, receptors and the like. However, target molecules may also include other molecules such as DNAs, lipids, carbohydrates and RNAs including messenger RNAs, ribosomal RNAs, tRNAs and the like. A number of highly sensitive cell-based assay methods are available to those of skill in the art to detect binding and interaction of test compounds with specific target molecules. However, these methods are generally not highly effective when the test compound binds to or otherwise interacts with its target molecule with moderate or low affinity. In addition, the target molecule may not be readily accessible to a test compound in solution, such as when the target molecule is located inside the cell or within a cellular compartment such as the periplasm of a bacterial cell. Thus, current cell-based assay methods are limited in

that they are not effective in identifying or characterizing compounds that interact with their targets with moderate to low affinity or compounds that interact with targets that are not readily accessible.

5 The cell-based assay methods of the present invention have substantial advantages over current cell-based assays. These advantages derive from the use of sensitized cells in which the level or activity of at least one proliferation-required gene product (the target molecule) has been specifically reduced to the point where the presence or absence of its function becomes a rate-determining step for cellular proliferation. In the cell-based assays of the present invention, the level or activity of
10 a proliferation-required gene product is reduced by transcribing an antisense nucleic acid complementary to at least a portion of the nucleic acid encoding the gene product from one of the fusion promoters described herein. Such sensitized cells become much more sensitive to compounds that are active against the affected target molecule. Thus, cell-based assays of the present invention are capable of detecting compounds
15 exhibiting low or moderate potency against the target molecule of interest because such compounds are substantially more potent on sensitized cells than on non-sensitized cells. The effect may be such that a test compound may be two to several times more potent, at least 10 times more potent, at least 20 times more potent, at least 50 times more potent, at least 100 times more potent, at least 1000 times more potent,
20 or even more than 1000 times more potent when tested on the sensitized cells as compared to the non-sensitized cells.

Due in part to the increased appearance of antibiotic resistance in pathogenic microorganisms and to the significant side-effects associated with some currently used antibiotics, novel antibiotics acting at new targets are highly sought after in the art.
25 Yet, another limitation in the current art related to cell-based assays is the problem of repeatedly identifying hits against the same kinds of target molecules in the same limited set of biological pathways. This may occur when compounds acting at such new targets are discarded, ignored or fail to be detected because compounds acting at the "old" targets are encountered more frequently and are more potent than
30 compounds acting at the new targets. As a result, the majority of antibiotics in use

currently interact with a relatively small number of target molecules within an even more limited set of biological pathways.

The use of sensitized cells comprising the fusion promoters of the current invention provides a solution to the above problem in two ways. First, desired
5 compounds acting at a target of interest, whether a new target or a previously known but poorly exploited target, can now be detected above the "noise" of compounds acting at the "old" targets due to the specific and substantial increase in potency of such desired compounds when tested on the sensitized cells of the current invention. Second, the methods used to sensitize cells to compounds acting at a target of interest
10 may also sensitize these cells to compounds acting at other target molecules within the same biological pathway. For example, expression of an antisense molecule to a gene encoding a ribosomal protein is expected to sensitize the cell to compounds acting at that ribosomal protein and may also sensitize the cells to compounds acting at any of the ribosomal components (proteins or rRNA) or even to compounds acting at any
15 target which is part of the protein synthesis pathway. Thus an important advantage of the present invention is the ability to reveal new targets and pathways that were previously not readily accessible to drug discovery methods.

Sensitized cells of the present invention are prepared by reducing the activity or level of a target molecule. The target molecule may be a gene product, such as an
20 RNA or polypeptide produced from the proliferation-required nucleic acids described herein. Alternatively, the target may be a gene product such as an RNA or polypeptide which is produced from a sequence within the same operon as the proliferation-required nucleic acids described herein. In addition, the target may be an RNA or polypeptide in the same biological pathway as the proliferation-required nucleic acids
25 described herein. Such biological pathways include, but are not limited to, enzymatic, biochemical and metabolic pathways as well as pathways involved in the production of cellular structures such the cell wall.

Current methods employed in the arts of medicinal and combinatorial chemistries are able to make use of structure-activity relationship information derived
30 from testing compounds in various biological assays including direct binding assays

and cell-based assays. Occasionally compounds are directly identified in such assays that are sufficiently potent to be developed as drugs. More often, initial hit compounds exhibit moderate or low potency. Once a hit compound is identified with low or moderate potency, directed libraries of compounds are synthesized and tested
5 in order to identify more potent leads. Generally these directed libraries are combinatorial chemical libraries consisting of compounds with structures related to the hit compound but containing systematic variations including additions, subtractions and substitutions of various structural features. When tested for activity against the target molecule, structural features are identified that either alone or in
10 combination with other features enhance or reduce activity. This information is used to design subsequent directed libraries containing compounds with enhanced activity against the target molecule. After one or several iterations of this process, compounds with substantially increased activity against the target molecule are identified and may be further developed as drugs. This process is facilitated by use of the sensitized cells
15 of the present invention since compounds acting at the selected targets exhibit increased potency in such cell-based assays, thus; more compounds can now be characterized providing more useful information than would be obtained otherwise.

Thus, it is now possible using cell-based assays of the present invention to identify or characterize compounds that previously would not have been readily
20 identified or characterized including compounds that act at targets that previously were not readily exploited using cell-based assays. The process of evolving potent drug leads from initial hit compounds is also substantially improved by the cell-based assays of the present invention because, for the same number of test compounds, more structure-function relationship information is likely to be revealed.

25 The method of sensitizing a cell entails selecting a suitable gene or operon. A suitable gene or operon is one whose expression is required for the proliferation of the cell to be sensitized. The next step is to introduce into the cells to be sensitized, an antisense RNA capable of hybridizing to the suitable gene or operon or to the RNA encoded by the suitable gene or operon. Introduction of the antisense RNA can be in
30 the form of an expression vector in which antisense RNA is produced under the

control of a fusion promoter as described herein. The amount of antisense RNA produced is limited by varying the inducer concentration to which the cell is exposed and thereby varying the activity of the promoter driving transcription of the antisense RNA. Thus, cells are sensitized by exposing them to an inducer concentration that
5 results in a sub-lethal level of antisense RNA expression.

In one embodiment of the cell-based assays, an antisense nucleic acid that is complementary to a proliferation-required gene from a Gram-positive organism is used to inhibit the production of a proliferation-required protein. Expression vectors which contain an antisense nucleic acid complementary to identified genes required
10 for proliferation operably linked to a fusion promoter of the present invention are used to limit the concentration of a proliferation-required protein without severely inhibiting growth. To achieve that goal, a growth inhibition dose curve of inducer is calculated by plotting various doses of inducer against the corresponding growth inhibition caused by the antisense expression. From this curve, various percentages of
15 antisense induced growth inhibition, from 1 to 100% can be determined. If the fusion promoter contained in the expression vector contains a *xylO* linked to one of the promoters of SEQ ID NO.: 36 - 45, transcription in *Staphylococcus aureus* and *Enterococcus faecalis* will be regulatable by the *xyl* repressor and expression from the promoter can be induced with xylose. Similarly, IPTG, xylose, tetracycline, maltose,
20 absence of *trp* and temperature inducible promoters may be used. For example, the highest concentration of the inducer that does not reduce the growth rate significantly can be estimated from the curve. Cellular proliferation can be monitored by growth medium turbidity via OD measurements. In another example, the concentration of inducer that reduces growth by 25% can be predicted from the curve. In still another
25 example, a concentration of inducer that reduces growth by 50% can be calculated. Additional parameters such as colony forming units (cfu) can be used to measure cellular viability.

Cells to be assayed are exposed to the above-determined concentrations of inducer. The presence of the inducer at this sub-lethal concentration reduces the
30 amount of the proliferation required gene product to the lowest amount in the cell that

will support growth. Cells grown in the presence of this concentration of inducer are therefore specifically more sensitive to inhibitors of the proliferation-required protein or RNA of interest or to inhibitors of proteins or RNAs in the same biological pathway as the proliferation-required protein or RNA of interest but not to inhibitors of
5 unrelated proteins or RNAs.

Cells pretreated with sub-inhibitory concentrations of inducer and thus containing a reduced amount of proliferation-required target gene product are then used to screen for compounds that reduce cell growth. The sub-lethal concentration of inducer may be any concentration consistent with the intended use of the assay to
10 identify candidate compounds to which the cells are more sensitive. For example, the sub-lethal concentration of the inducer may be such that growth inhibition is at least about 5%, at least about 8%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60% at least about 75%, or more. Cells which are pre-sensitized using the preceding method are more sensitive to
15 inhibitors of the target protein because these cells contain less target protein to inhibit than wild-type cells.

It will be appreciated that the cell-based assays described above may be implemented in *Staphylococcus aureus* and *Enterococcus faecalis* using proliferation-inhibiting antisense sequences under the control of a fusion promoter described herein. It will
20 also be appreciated that the above cell-based assays employing the fusion promoters described herein can be implemented in other Gram-positive organisms, including but not limited to, *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria*
25 *monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species.

In another embodiment of the cell-based assays of the present invention, the
30 level or activity of a proliferation required gene product is reduced using a mutation,

such as a temperature sensitive mutation, in the proliferation-required sequence and an antisense nucleic acid complementary to the proliferation-required sequence which is transcribed from one of the fusion promoters described herein. Growing the cells at an intermediate temperature between the permissive and restrictive temperatures of the temperature sensitive mutant where the mutation is in a proliferation-required gene produces cells with reduced activity of the proliferation-required gene product. The antisense RNA complementary to the proliferation-required sequence further reduces the activity of the proliferation required gene product. Drugs that may not have been found using either the temperature sensitive mutation or the antisense nucleic acid alone may be identified by determining whether cells in which expression of the antisense nucleic acid has been induced and which are grown at a temperature between the permissive temperature and the restrictive temperature are substantially more sensitive to a test compound than cells in which expression of the antisense nucleic acid has not been induced and which are grown at a permissive temperature. Also drugs found previously from either the antisense nucleic acid alone or the temperature sensitive mutation alone may have a different sensitivity profile when used in cells combining the two approaches, and that sensitivity profile may indicate a more specific action of the drug in inhibiting one or more activities of the gene product.

Temperature sensitive mutations may be located at different sites within the gene and correspond to different domains of the protein. For example, the *dnaB* gene of *Escherichia coli* encodes the replication fork DNA helicase. DnaB has several domains, including domains for oligomerization, ATP hydrolysis, DNA binding, interaction with primase, interaction with DnaC, and interaction with DnaA [(Biswas, E.E. and Biswas, S.B. 1999, "Mechanism and DnaB helicase of *Escherichia coli*: structural domains involved in ATP hydrolysis, DNA binding, and oligomerization" *Biochem.* 38: 10919-10928; Hiasa, H. and Marians, K.J. 1999, "Initiation of bidirectional replication at the chromosomal origin is directed by the interaction between helicase and primase" *J. Biol. Chem.* 274: 27244-27248; San Martin, C., et al. 1998, "Three-dimensional reconstructions from cryoelectron microscopy images

reveal an intimate complex between helicase DnaB and its loading partner DnaC”
Structure 6:501-9; Sutton, M.D. *et al.* 1998. “*Escherichia coli* DnaA protein. The N-
terminal domain and loading of DnaB helicase at the *E. coli* chromosomal origin” *J.*
Biol. Chem. 273: 34255-62.), the disclosures of which are incorporated herein by
5 reference in their entireties]. Temperature sensitive mutations in different domains of
DnaB confer different phenotypes at the restrictive temperature, which include either
an abrupt stop or slow stop in DNA replication with or without DNA breakdown
(Wechsler, J.A. and Gross, J.D. 1971. “*Escherichia coli* mutants temperature-sensitive
for DNA synthesis” *Mol. Gen. Genetics* 113: 273-284, the disclosure of which is
10 incorporated herein by reference in its entirety) and termination of growth or cell
death. Combining the use of temperature sensitive mutations in the *dnaB* gene that
cause cell death at the restrictive temperature with an antisense to the *dnaB* gene could
lead to the discovery of very specific and effective inhibitors of one or a subset of
activities exhibited by DnaB.

15 It will be appreciated that the above method may be performed with any
mutation which reduces but does not eliminate the activity or level of the gene product
which is required for proliferation.

When screening for antimicrobial agents against a gene product required for
proliferation, growth inhibition of cells containing a limiting amount of that
20 proliferation-required gene product can be assayed. Growth inhibition can be
measured by directly comparing the amount of growth, measured by the optical
density of the growth medium, between an experimental sample and a control sample.
Alternative methods for assaying cell proliferation include measuring the activity of
any of a variety of reporter genes, various enzymatic activity assays, and other
25 methods well known in the art.

It will be appreciated that the above method may be performed in solid phase,
liquid phase or a combination of the two. For example, cells grown on nutrient agar
containing the inducer of the antisense construct may be exposed to compounds
spotted onto the agar surface. A compound’s effect may be judged from the diameter
30 of the resulting killing zone, the area around the compound application point in which

cells do not grow. Multiple compounds may be transferred to agar plates and simultaneously tested using automated and semi-automated equipment including but not restricted to multi-channel pipettes (for example the Beckman Multimek) and multi-channel spotters (for example the Genomic Solutions Flexys). In this way
5 multiple plates and thousands to millions of compounds may be tested per day.

The compounds may also be tested entirely in liquid phase using microtiter plates as described below. Liquid phase screening may be performed in microtiter plates containing 96, 384, 1536 or more wells per microtiter plate to screen multiple plates and thousands to millions of compounds per day. Automated and semi-
10 automated equipment may be used for addition of reagents (for example cells and compounds) and determination of cell density.

The examples below describe cell-based assays that exploit the fusion promoters described herein to sensitize cells and allow for the discovery of a diverse class of antibiotics. Example 20 describes experiments performed in *E. coli*, which
15 verify the effectiveness of the cell-based approach described above.

EXAMPLE 20

Cell-Based Assay Using Antisense RNA Complementary to Genes Encoding Ribosomal Proteins

20 The effectiveness of the above cell-based assay was validated using constructs expressing antisense RNA to the proliferation required *E. coli* genes *rplL*, *rplJ*, and *rplW* encoding ribosomal proteins L7/L12, L10 and L23 respectively. These proteins are part of the protein synthesis apparatus of the cell and as such are required for proliferation. These constructs were used to test the effect of antisense expression on
25 cell sensitivity to antibiotics known to bind to the ribosome and thereby inhibit protein synthesis. Constructs expressing antisense RNA to several other genes (*elaD*, *visC*, *yohH*, and *atpE/B*), the products of which are not involved in protein synthesis were used for comparison.

First, pLex5BA (Krause *et al.* 1997. *J. Mol. Biol.* 274: 365, the disclosure of
30 which is incorporated herein by reference in its entirety) expression vectors containing

antisense constructs to either *rplW* or to *elaD* were introduced into separate *E. coli* cell populations. Vector introduction is a technique well known to those of ordinary skill in the art. The expression vectors of this example contain IPTG inducible promoters that drive the expression of the antisense RNA in the presence of the inducer.

5 However, those skilled in the art will appreciate that other inducible promoters may also be used. Suitable expression vectors are also well known in the art. The *E. coli* antisense clones to genes encoding ribosomal proteins L7/L12, L10 and L23 were used to test the effect of antisense expression on cell sensitivity to the antibiotics known to bind to these proteins. Expression vectors containing antisense to either the

10 genes encoding L7/L12 and L10 or L23 were introduced into separate *E. coli* cell populations.

The cell populations were exposed to a range of IPTG concentrations in liquid medium to obtain the growth inhibitory dose curve for each clone. First, seed cultures were grown to a particular turbidity that is measured by the optical density (OD) of the

15 growth solution. The OD of the solution is directly related to the number of bacterial cells contained therein. Subsequently, sixteen 200 ul liquid medium cultures were grown in a 96 well microtiter plate at 37° C with a range of IPTG concentrations in duplicate two-fold serial dilutions from 1600 uM to 12.5 uM (final concentration). Additionally, control cells were grown in duplicate without IPTG. These cultures

20 were started from equal amounts of cells derived from the same initial seed culture of a clone of interest. The cells were grown for up to 15 hours and the extent of growth was determined by measuring the optical density of the cultures at 600 nm. When the control culture reached mid-log phase the percent growth (relative to the control culture) for each of the IPTG containing cultures was plotted against the log

25 concentrations of IPTG to produce a growth inhibitory dose response curve for the IPTG. The concentration of IPTG that inhibits cell growth to 50% (IC₅₀) as compared to the 0 mM IPTG control (0% growth inhibition) was then calculated from the curve. Under these conditions, an amount of antisense RNA was produced that reduced the expression levels of *rplW* and *elaD* to a degree such that growth was inhibited by

30 50%.

Alternative methods of measuring growth are also contemplated. Examples of these methods include measurements of proteins, the expression of which is engineered into the cells being tested and can readily be measured. Examples of such proteins include green fluorescent protein (GFP) and various enzymes.

5 Cells were pretreated with the selected concentration of IPTG and then used to test the sensitivity of cell populations to tetracycline, erythromycin and other protein synthesis inhibitors. FIG. 12 is an IPTG dose response curve in *E. coli* transformed with an IPTG-inducible plasmid containing either an antisense clone to the *E. coli* *rplW* gene (AS-*rplW*) which encodes ribosomal protein L23 which is required for
10 protein synthesis and essential for cell proliferation, or an antisense clone to the *elaD* (AS-*elaD*) gene which is not known to be involved in protein synthesis and which is also essential for proliferation.

 An example of a tetracycline dose response curve is shown in FIGs. 13A and 13B for the *rplW* and *elaD* genes, respectively. Cells were grown to log phase and
15 then diluted into media alone or media containing IPTG at concentrations which give 20% and 50% growth inhibition as determined by IPTG dose response curves. After 2.5 hours, the cells were diluted to a final OD₆₀₀ of 0.002 into 96 well plates containing (1) +/- IPTG at the same concentrations used for the 2.5 hour pre-incubation; and (2) serial two-fold dilutions of tetracycline such that the final
20 concentrations of tetracycline range from 1 µg/ml to 15.6 ng/ml and 0 µg/ml. The 96 well plates were incubated at 37°C and the OD₆₀₀ was read by a plate reader every 5 minutes for up to 15 hours. For each IPTG concentration and the no IPTG control, tetracycline dose response curves were determined when the control (absence of tetracycline) reached 0.1 OD₆₀₀. To compare tetracycline sensitivity with and without
25 IPTG, tetracycline IC_{50s} were determined from the dose response curves (FIGs. 13A and 13B). Cells with reduced levels of L23 (AS-*rplW*) showed increased sensitivity to tetracycline (FIG. 13A) as compared to cells with reduced levels of the *elaD* gene product (AS-*elaD*) (FIG. 13B). FIG. 14 shows a summary bar chart in which the ratios of tetracycline IC_{50s} determined in the presence of IPTG which gives 50%
30 growth inhibition versus tetracycline IC_{50s} determined without IPTG (fold increase in

tetracycline sensitivity) were plotted. Cells with reduced levels of either L7/L12 (encoded by genes *rplL*, *rplJ*) or L23 (encoded by the *rplW* gene) showed increased sensitivity to tetracycline (FIG. 14). Cells expressing antisense to genes not known to be involved in protein synthesis (*AS-atpB/E*, *AS-visC*, *AS-elaD*, *AS-yohH*) did not
5 show the same increased sensitivity to tetracycline, validating the specificity of this assay (FIG. 14).

In addition to the above, it has been observed in initial experiments that clones expressing antisense RNA to genes involved in protein synthesis (including genes encoding ribosomal proteins L7/L12 & L10, L7/L12 alone, L22, and L18, as well as
10 genes encoding rRNA and Elongation Factor G) have increased sensitivity to the macrolide, erythromycin, whereas clones expressing antisense to the non-protein synthesis genes *elaD*, *atpB/E* and *visC* do not. Furthermore, the clone expressing antisense to *rplL* and *rplJ* does not show increased sensitivity to nalidixic acid and ofloxacin, antibiotics which do not inhibit protein synthesis.

The results with the ribosomal protein genes *rplL*, *rplJ*, and *rplW* as well as the
15 initial results using various other antisense clones and antibiotics show that limiting the concentration of an antibiotic target makes cells more sensitive to the antimicrobial agents that specifically interact with that protein. The results also show that these cells are sensitized to antimicrobial agents that inhibit the overall function in
20 which the protein target is involved but are not sensitized to antimicrobial agents that inhibit other functions.

It will be appreciated that the cell-based assays described above may be implemented in *Staphylococcus aureus* and *Enterococcus faecalis* using proliferation-inhibiting antisense sequences under the control of a fusion promoter described herein. It will
25 also be appreciated that the above cell-based assays employing the fusion promoters described herein can be implemented in other Gram-positive organisms, including but not limited to, *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia*
30

asteroides, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species.

The cell-based assay described above may also be used to identify the
5 biological pathway in which a proliferation-required nucleic acid or its gene product lies. In such methods, cells transcribing a sub-lethal level of antisense to a target proliferation-required nucleic acid from the fusion promoters described herein and control cells in which expression of the antisense has not been induced are contacted with a panel of antibiotics known to act in various pathways. If the antibiotic acts in
10 the pathway in which the target proliferation-required nucleic acid or its gene product lies, cells in which expression of the antisense has been induced will be more sensitive to the antibiotic than cells in which expression of the antisense has not been induced.

As a control, the results of the assay may be confirmed by contacting a panel of cells expressing antisense nucleic acids to many different proliferation-required genes
15 including the target proliferation-required gene. If the antibiotic is acting specifically, heightened sensitivity to the antibiotic will be observed only in the cells expressing antisense to a target proliferation-required gene (or cells expressing antisense to other proliferation-required genes in the same pathway as the target proliferation-required gene) but will not be observed generally in all cells expressing antisense to
20 proliferation-required genes.

It will be appreciated that the cell-based assays described above may be implemented in *Staphylococcus aureus* and *Enterococcus faecalis* using proliferation-inhibiting antisense sequences under the control of a fusion promoter described herein. It will also be appreciated that the above cell-based assays employing the fusion
25 promoters described herein can be implemented in other Gram-positive organisms, including but not limited to, *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia*
30 *asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus*

xyloxis, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species.

Similarly, the above method may be used to determine the pathway on which a test compound, such as a test antibiotic acts. A panel of cells, each of which transcribes antisense to a proliferation-required nucleic acid in a known pathway from one of the fusion promoters described herein, is contacted with a compound for which it is desired to determine the pathway on which it acts. The sensitivity of the panel of cells to the test compound is determined in cells in which expression of the antisense has been induced and in control cells in which expression of the antisense has not been induced. If the test compound acts on the pathway on which an antisense nucleic acid acts, cells in which expression of the antisense has been induced will be more sensitive to the compound than cells in which expression of the antisense has not been induced. In addition, control cells in which expression of antisense to proliferation-required genes in other pathways has been induced will not exhibit heightened sensitivity to the compound. In this way, the pathway on which the test compound acts may be determined.

It will be appreciated that the cell-based assays described above may be implemented in *Staphylococcus aureus* and *Enterococcus faecalis* using proliferation-inhibiting antisense sequences under the control of a fusion promoter described herein. It will also be appreciated that the above cell-based assays employing the fusion promoters described herein can be implemented in other Gram-positive organisms, including but not limited to, *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xyloxis*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species.

The Example below provides one method for performing such assays.

30

EXAMPLE 21**Identification of the Pathway in which a Proliferation-Required****Gene Lies on the Pathway on which an Antibiotic Acts****A. Preparation of Bacterial Stocks for Assay**

5 To provide a consistent source of cells to screen, frozen stocks of host bacteria containing the desired antisense construct are prepared using standard microbiological techniques. For example, a single clone of the microorganism can be isolated by streaking out a sample of the original stock onto an agar plate containing nutrients for cell growth and an antibiotic for which the antisense construct contains a gene which
10 confers resistance. After overnight growth an isolated colony is picked from the plate with a sterile needle and transferred to an appropriate liquid growth media containing the antibiotic required for maintenance of the plasmid. The cells are incubated at 30°C to 37°C with vigorous shaking for 4 to 6 hours to yield a culture in exponential growth. Sterile glycerol is added to 15% (volume to volume) and 100µL to 500 µL
15 aliquots are distributed into sterile cryotubes, snap frozen in liquid nitrogen, and stored at -80°C for future assays.

B. Growth of Bacteria for Use in the Assay

 A day prior to an assay, a stock vial is removed from the freezer, rapidly thawed (37°C water bath) and a loop of culture is streaked out on an agar plate
20 containing nutrients for cell growth and an antibiotic to which the antisense construct confers resistance. After overnight growth at 37°C, ten randomly chosen, isolated colonies are transferred from the plate (sterile inoculum loop) to a sterile tube containing 5 mL of appropriate medium containing the antibiotic to which the antisense vector confers resistance. After vigorous mixing to form a homogeneous cell
25 suspension, the optical density of the suspension is measured at 600 nm (OD₆₀₀) and if necessary an aliquot of the suspension is diluted into a second tube of 5 mL, sterile, appropriate medium plus antibiotic to achieve an OD₆₀₀ ≤ 0.02 absorbance units. The culture is then incubated at 37° C for 1-2 hrs with shaking until the OD₆₀₀ reaches OD 0.2 – 0.3. At this point the cells are ready to be used in the assay.

C. Selection of Media to be Used in Assay

Two-fold dilution series of the inducer are generated in culture media containing the appropriate antibiotic for maintenance of the antisense construct. Several media are tested side by side and three to four wells are used to evaluate the effects of the inducer at each concentration in each media. For example, LB broth, TBD broth, Muller-Hinton, or other media may be tested with the inducer xylose at the following concentrations, 5 mM, 10 mM, 20 mM, 40 mM, 80 mM, 120 mM and 160 mM. Equal volumes of test media-inducer and cells are added to the wells of a 384 well microtiter plate and mixed. The cells are prepared as described above and diluted 1:100 in the appropriate media containing the test antibiotic immediately prior to addition to the microtiter plate wells. For a control, cells are also added to several wells of each media that do not contain inducer, for example 0 mM xylose. Cell growth is monitored continuously by incubation at 37°C in a microtiter plate reader monitoring the OD₆₀₀ of the wells over an 18-hour period. The percent inhibition of growth produced by each concentration of inducer is calculated by comparing the rates of logarithmic growth against that exhibited by cells growing in media without inducer. The medium yielding greatest sensitivity to inducer is selected for use in the assays described below.

D. Measurement of Test Antibiotic Sensitivity in the Absence of Antisense Construct Induction

Two-fold dilution series of antibiotics of known mechanism of action are generated in the culture media selected for further assay development that has been supplemented with the antibiotic used to maintain the construct. A panel of test antibiotics known to act on different pathways is tested side by side with three to four wells being used to evaluate the effect of a test antibiotic on cell growth at each concentration. Equal volumes of test antibiotic and cells are added to the wells of a 384 well microtiter plate and mixed. Cells are prepared as described above using the media selected for assay development supplemented with the antibiotic required to maintain the antisense construct and are diluted 1:100 in identical media immediately prior to addition to the microtiter plate wells. For a control, cells are also added to several wells that lack antibiotic, but contain the solvent used to dissolve the

antibiotics. Cell growth is monitored continuously by incubation at 37°C in a microtiter plate reader monitoring the OD₆₀₀ of the wells over an 18-hour period. The percent inhibition of growth produced by each concentration of antibiotic is calculated by comparing the rates of logarithmic growth against that exhibited by cells growing
5 in media without antibiotic. A plot of percent inhibition against log[antibiotic concentration] allows extrapolation of an IC₅₀ value for each antibiotic.

E. Measurement of Test Antibiotic Sensitivity in the Presence of Antisense Construct Inducer

The culture media selected for use in the assay is supplemented with inducer at
10 concentrations shown to inhibit cell growth by 50% and 80% as described above, as well as the antibiotic used to maintain the construct. Two fold dilution series of the panel of test antibiotics used above are generated in each of these media. Several antibiotics are tested side by side in each medium with three to four wells being used to evaluate the effects of an antibiotic on cell growth at each concentration. Equal
15 volumes of test antibiotic and cells are added to the wells of a 384 well microtiter plate and mixed. Cells are prepared as described above using the media selected for use in the assay supplemented with the antibiotic required to maintain the antisense construct. The cells are diluted 1:100 into two 50 mL aliquots of identical media containing concentrations of inducer that have been shown to inhibit cell growth by
20 50% and 80 % respectively and incubated at 37°C with shaking for 2.5 hours. Immediately prior to addition to the microtiter plate wells, the cultures are adjusted to an appropriate OD₆₀₀ (typically 0.002) by dilution into warm (37°C) sterile media supplemented with identical concentrations of the inducer and antibiotic used to maintain the antisense construct. For a control, cells are also added to several wells
25 that contain solvent used to dissolve test antibiotics but which contain no antibiotic. Cell growth is monitored continuously by incubation at 37°C in a microtiter plate reader monitoring the OD₆₀₀ of the wells over an 18-hour period. The percent inhibition of growth produced by each concentration of antibiotic is calculated by comparing the rates of logarithmic growth against that exhibited by cells growing in
30 media without antibiotic. A plot of percent inhibition against log[antibiotic

concentration] allows extrapolation of an IC_{50} value for each antibiotic.

F. Determining the Specificity of the Test Antibiotics

5 A comparison of the IC_{50} s generated by antibiotics of known mechanism of action under antisense induced and non-induced conditions allows the pathway in which a proliferation-required nucleic acid lies to be identified. If cells expressing an antisense nucleic acid complementary to a proliferation-required gene are selectively sensitive to an antibiotic acting via a particular pathway, then the gene against which the antisense acts is involved in the pathway on which the antibiotic acts.

G. Identification of Pathway in which a Test Antibiotic Acts

10 As discussed above, the cell-based assay may also be used to determine the pathway against which a test antibiotic acts. In such an analysis, the pathways against which each member of a panel of antisense nucleic acids acts are identified as described above. A panel of cells, each containing an inducible vector which transcribes an antisense nucleic acid complementary to a gene in a known proliferation-required pathway from one of the fusion promoters described herein, is
15 contacted with a test antibiotic for which it is desired to determine the pathway on which it acts under inducing and non-inducing conditions. If heightened sensitivity is observed in induced cells expressing antisense complementary to a gene in a particular pathway but not in induced cells expressing antisense complementary to genes in other
20 pathways, then the test antibiotic acts against the pathway for which heightened sensitivity was observed.

One skilled in the art will appreciate that further optimization of the assay conditions, such as the concentration of inducer used to induce antisense expression and/or the growth conditions used for the assay (for example incubation temperature
25 and media components) may further increase the selectivity and/or magnitude of the antibiotic sensitization exhibited.

The following example confirms the effectiveness of the methods described above.

30

EXAMPLE 22

Identification of the Biological Pathway in Which a Proliferation-Required Gene Lies

The effectiveness of the above assays was validated using proliferation-required genes from *E. coli* which were identified using procedures similar to those described above. Antibiotics of various chemical classes and modes of action were purchased from Sigma Chemicals (St. Louis, MO). Stock solutions were prepared by dissolving each antibiotic in an appropriate aqueous solution based on information provided by the manufacturer. The final working solution of each antibiotic contained no more than 0.2% (w/v) of any organic solvent. To determine their potency against a bacterial strain engineered for expression of an antisense complementary to a proliferation-required 50S ribosomal protein, each antibiotic was serially diluted two or three fold in growth medium supplemented with the appropriate antibiotic for maintenance of the anti-sense construct. At least ten dilutions were prepared for each antibiotic. 25 μ L aliquots of each dilution were transferred to discrete wells of a 384-well microplate (the assay plate) using a multi-channel pipette. Quadruplicate wells were used for each dilution of an antibiotic under each treatment condition (plus and minus inducer). Each assay plate contained twenty wells for cell growth controls (growth media replacing antibiotic), ten wells for each treatment (plus and minus inducer, in this example IPTG). Assay plates were usually divided into the two treatments: half the plate containing induced cells and an appropriate concentrations of inducer (in this example IPTG) to maintain the state of induction, the other half containing non-induced cells in the absence of IPTG.

Cells for the assay were prepared as follows. Bacterial cells containing a construct, from which expression of antisense nucleic acid complementary to *rplL* and *rplJ*, which encode proliferation-required 50S ribosomal subunit proteins, is inducible in the presence of IPTG, were grown into exponential growth (OD_{600} 0.2 to 0.3) and then diluted 1:100 into fresh media containing either 400 μ M or 0 μ M inducer (IPTG). These cultures were incubated at 37° C for 2.5 hr. After a 2.5 hr incubation, induced and non-induced cells were respectively diluted into an assay medium at a final OD_{600} value of 0.0004. The medium contained an appropriate concentration of the antibiotic for the maintenance of the antisense construct. In addition, the medium used to dilute

induced cells was supplemented with 800 μ M IPTG so that addition to the assay plate would result in a final IPTG concentration of 400 μ M. Induced and non-induced cell suspensions were dispensed (25 μ l/well) into the appropriate wells of the assay plate as discussed previously. The plate was then loaded into a plate reader, incubated at
5 constant temperature, and cell growth was monitored in each well by the measurement of light scattering at 595 nm. Growth was monitored every 5 minutes until the cell culture attained a stationary growth phase. For each concentration of antibiotic, a percentage inhibition of growth was calculated at the time point corresponding to mid-exponential growth for the associated control wells (no antibiotic, plus or minus
10 IPTG). For each antibiotic and condition (plus or minus IPTG), a plot of percent inhibition versus log of antibiotic concentration was generated and the IC_{50} determined. A comparison of the IC_{50} for each antibiotic in the presence and absence of IPTG revealed whether induction of the antisense construct sensitized the cell to the mechanism of action exhibited by the antibiotic. Cells which exhibited a significant
15 (standard statistical analysis) numerical decrease in the IC_{50} value in the presence of inducer were considered to have an increased sensitivity to the test antibiotic.

The above results demonstrate that induction of an antisense RNA to genes encoding 50S ribosomal subunit proteins results in a selective and highly significant sensitization of cells to antibiotics that inhibit ribosomal function and protein
20 synthesis. The above results further demonstrate that induction of an antisense construct to an essential gene sensitizes a microorganism to compounds that interfere with that gene product's biological role. This sensitization is restricted to compounds that interfere with pathways associated with the targeted gene and its product. It will be appreciated that the cell-based assays described above may be implemented in
25 *Staphylococcus aureus* and *Enterococcus faecalis* using proliferation-inhibiting antisense sequences under the control of a fusion promoter described herein..

The example below describes an analysis performed in *Staphylococcus aureus*.

EXAMPLE 23

Identification of the Biological Pathway in which a Gene Required for Proliferation of
Staphylococcus aureus Lies

Sensitized cells were also used to identify the pathway in which a gene required for proliferation of *Staphylococcus aureus* lies. Antibiotics of various
5 chemical classes and modes of action were purchased from chemical suppliers, for example Sigma Chemicals (St. Louis, MO). Stock solutions were prepared by dissolving each antibiotic in an appropriate aqueous solution based on information provided by the manufacturer. The final working solution of each antibiotic contained no more than 0.2% (w/v) of any organic solvent.

10 To determine its potency against a bacterial strain containing an antisense nucleic acid complementary to the sequence encoding the Beta (β) subunit of gyrase (which is required for proliferation) under the control of the Xyl-T5 fusion promoter, each antibiotic was serially diluted two or three fold in growth medium supplemented with the appropriate antibiotic for maintenance of the anti-sense construct. At least
15 ten dilutions were prepared for each antibiotic.

Aliquots (25 μ L) of each dilution were transferred to discrete wells of a 384-well microplate (the assay plate) using a multi-channel pipette. Quadruplicate wells were used for each dilution of an antibiotic under each treatment condition (plus and minus inducer). Each assay plate retained twenty wells for cell growth controls
20 (growth media replacing antibiotic), ten wells for each treatment (plus and minus inducer, in this example xylose). Half the assay plate contained induced cells (in this example *Staphylococcus aureus* cells) and appropriate concentrations of inducer (in this example xylose) to maintain the state of induction while the other half of the assay plate contained non-induced cells in the absence of inducer.

25 Cells of a bacterial clone containing a construct in which expression of antisense complementary to the sequence encoding the β subunit of gyrase under the control of a xylose inducible promoter were grown into exponential growth (OD₆₀₀ 0.2 to 0.3) and then diluted 1:100 into fresh media containing either 12 mM or 0 mM inducer (xylose). These cultures were incubated at 37° C for 2.5 hr. The presence of
30 inducer (xylose) in the medium initiates and maintains production of antisense RNA

from the antisense construct, in this case. After a 2.5 hr incubation, induced and non-induced cells were respectively diluted into an assay medium containing an appropriate concentration of the antibiotic for the maintenance of the anti-sense construct. In addition, medium used to dilute induced cells was supplemented with 24 mM xylose so that addition to the assay plate would result in a final xylose concentration of 12 mM. The cells were diluted to a final OD₆₀₀ value of 0.0004.

Induced and non-induced cell suspensions were dispensed (25 μ l/well) into the appropriate wells of the assay plate as discussed previously. The plate was then loaded into a plate reader and incubated at constant temperature while cell growth was monitored in each well by the measurement of light scattering at 595 nm. Growth was monitored every 5 minutes until the cell culture attained a stationary growth phase. For each concentration of antibiotic, a percentage inhibition of growth was calculated at the time point corresponding to mid-exponential growth for the associated control wells (no antibiotic, plus or minus xylose). For each antibiotic and condition (plus or minus xylose), plots of percent inhibition versus Log of antibiotic concentration were generated and IC₅₀s determined.

A comparison of each antibiotic's IC₅₀ in the presence and absence of inducer (in this example xylose) reveals whether induction of the antisense construct sensitized the cell to the antibiotic's mechanism of action. If the antibiotic acts against the β subunit of gyrase, the IC₅₀ of induced cells will be significantly lower than the IC₅₀ of uninduced cells.

FIG. 15 lists the antibiotics tested, their targets, and their fold increase in potency between induced cells and uninduced cells. As illustrated in FIG. 15, the potency of cefotaxime, ceftazidime, fusidic acid, lincomycin, tobramycin, trimethoprim and vancomycin, each of which act on targets other than the β subunit of gyrase, was not significantly different in induced cells as compared to uninduced cells. However, the potency of novobiocin, which acts against the β subunit of gyrase, was significantly different between induced cells and uninduced cells.

Thus, induction of an antisense nucleic acid complementary to the sequence encoding the β subunit of gyrase results in a selective and significant sensitization of

Staphylococcus aureus cells to an antibiotic which inhibits the activity of this protein. Furthermore, the results demonstrate that induction of an antisense construct to an essential gene sensitizes a microorganism to compounds that interfere with that gene product's biological role. This sensitization is apparently restricted to compounds that
5 interfere with the targeted gene and its product.

It will be appreciated that the cell-based assays described above may be implemented in *Enterococcus faecalis* using proliferation-inhibiting antisense sequences under the control of a fusion promoter described herein. It will also be appreciated that the above cell-based assays employing the fusion promoters described herein can be
10 implemented in other Gram-positive organisms, including but not limited to, *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*,
15 *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species.

Assays utilizing antisense constructs to essential genes can be used to identify compounds that interfere with the activity of those gene products. Such assays could
20 be used to identify drug leads, for example antibiotics.

Assays utilizing antisense constructs to essential genes can be used to identify compounds that specifically interfere with the activity of multiple targets in a pathway. Such constructs can be used to simultaneously screen a sample against multiple targets in one pathway in one reaction (Combinatorial HTS).

25 Furthermore, as discussed above, panels of antisense construct-containing cells may be used to characterize the point of intervention of any compound affecting an essential biological pathway including antibiotics with no known mechanism of action.

The following example provides the a method for determining the pathway on
30 which an antibiotic acts.

EXAMPLE 24**Cell-based assays using Promoter Replacement or Operator Insertion**

5 A gene encoding a gene product required for cellular proliferation is placed
under the control of an inducible promoter, such as one of the inducible fusion
promoters described herein, using the promoter replacement or operator insertion
methods described above. Sensitized cells are generated by growing cells in which the
gene encoding the gene product required for proliferation is under control of the
inducible fusion promoter in medium containing varying concentrations of inducer
10 which are less than the inducer concentration in which the promoter has maximal
activity. The sensitized cells are contacted with candidate compounds to be tested for
the ability to inhibit proliferation. In some embodiments, the candidate compounds
may be compounds in a library of compounds generated using combinatorial
chemistry techniques or compounds in a natural products library. The extent of
15 proliferation of the sensitized cells is compared to the extent of proliferation of control
cells in which the activity or level of the gene product is higher than in the sensitized
cells. For example, the control cells may be cells in which the inducible fusion
promoter is fully active. A compound which inhibits the growth of the sensitized cells
to a substantially greater extent than the control cells is identified. This compound
20 may be used as a drug or further optimized to generate a more potent compound.
It will be appreciated that the cell-based assays described above may be implemented
in *Staphylococcus aureus* and *Enterococcus faecalis* using the fusion promoters or
operators described herein. It will also be appreciated that the above cell-based assays
employing the fusion promoters or operators described herein can be implemented in
25 other Gram-positive organisms, including but not limited to, *Bacillus anthracis*,
Clostridium botulinum, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium*
teteni, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*,
Lactococcus lactis, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium*
tuberculosis, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus*

epidermidis, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species.

The example below describes an approach to determine if an antisense nucleic acid known to inhibit cell proliferation of one organism can successfully inhibit cell proliferation of a second organism.

EXAMPLE 25

Identification of Proliferation-Required Genes in Using Antisense Nucleic Acids from Heterologous Organisms

Once an antisense nucleic acid which inhibits the proliferation of a Gram-positive organism has been identified using the fusion promoters described herein, the antisense nucleic acid may be used to identify proliferation-required genes in heterologous organisms. In such methods, the antisense nucleic acid is transcribed in the heterologous organism and its ability to inhibit proliferation of the heterologous organism is determined.

In such embodiments, the antisense nucleic acids are inserted into expression vectors functional in the organisms in which the antisense nucleic acids are evaluated. Those skilled in the art will appreciate that a negative result in a heterologous microorganism does not mean that that microorganism is missing that gene nor does it mean that the gene is unessential. However, a positive result means that the heterologous microorganism contains a homologous gene which is required for proliferation of that microorganism. The homologous gene may be obtained using the methods described herein. Those cells that are inhibited by antisense may be used in cell-based assays as described herein for the identification and characterization of compounds in order to develop antibiotics effective in these microorganisms. Those skilled in the art will appreciate that an antisense molecule that works in the microorganism from which it was obtained will not always work in a heterologous microorganism.

The antisense nucleic acids that inhibit the growth of *Staphylococcus aureus* or *Enterococcus faecalis*, for example, may also be evaluated for their ability to inhibit

the growth of microorganisms other than *Staphylococcus aureus* or *Enterococcus faecalis*. In particular, the ability of the antisense nucleic acid to inhibit the growth of *Aspergillus fumigatus*, *Bacillus anthracis*, *Burkholderia cepacia*, *Campylobacter jejuni*, *Candida albicans*, *Chlamydia pneumoniae*, *Chlamydia trachomatis*,
 5 *Clostridium botulinum*, *Cryptococcus neoformans*, *Enterobacter cloacae*, *Enterococcus faecalis*, *Escherichia coli*, *Haemophilus influenzae*, *Helicobacter pylori*, *Klebsiella pneumoniae*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Neisseria gonorrhoeae*, *Nocardia asteroides*, *Pseudomonas aeruginosa*, *Salmonella choleraesuis*, *Salmonella paratyphi*, *Salmonella typhi*, *Salmonella typhimurium*,
 10 *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Moxarella catarrhalis*, *Shigella boydii*, *Shigella dysenteriae*, *Shigella flexneri*, *Shigella sonnei*, *Pseudomonas aeruginosa*, *Staphylococcus epidermidis*, *Streptococcus pneumoniae*, *Treponema pallidum*, *Yersinia pestis*, or any species falling within the genera of any of the above species may be evaluated.

15 In such methods, expression vectors in which the expression of an antisense nucleic acid that inhibits the growth of *Staphylococcus aureus* or *Enterococcus faecalis* is under the control of an inducible promoter, such as a fusion promoter described herein, are introduced into the microorganisms in which they are to be evaluated. In some embodiments, the antisense nucleic acids may be evaluated in
 20 microorganisms which are closely related to *Staphylococcus aureus* or *Enterococcus faecalis*. The ability of these antisense nucleic acids to inhibit the growth of the related microorganisms in the presence of the inducer is then measured.

In one example, thirty-nine antisense nucleic acids which inhibited the growth of *Staphylococcus aureus* were identified using methods such as those described
 25 herein and were inserted into an expression vector such that their expression was under the control of a xylose-inducible Xyl-T5 promoter. A vector with a reporter gene under control of the Xyl-T5 promoter was used to show that expression from the Xyl-T5 promoter in *Staphylococcus epidermidis* was comparable to that in *Staphylococcus aureus*.

30 The vectors were introduced into *Staphylococcus epidermidis* by

electroporation as follows: *Staphylococcus epidermidis* was grown in liquid culture to mid-log phase and then harvested by centrifugation. The cell pellet was resuspended in 1/3 culture volume of ice-cold EP buffer (0.625 M sucrose, 1 mM MgCl₂, pH=4.0), and then harvested again by centrifugation. The cell pellet was then resuspended with
 5 1/40 volume EP buffer and allowed to incubate on ice for 1 hour. The cells were then frozen for storage at -80°C. For electroporation, 50 µl of thawed electrocompetent cells were combined with 0.5 µg plasmid DNA and then subjected to an electrical pulse of 10 kV/cm, 25 uFarads, 200 ohm using a BioRad gene pulser electroporation device. The cells were immediately resuspended with 200 µl outgrowth medium and
 10 incubated for 2 hours prior to plating on solid growth medium with drug selection to maintain the plasmid vector. Colonies resulting from overnight growth of these platings were selected, cultured in liquid medium with drug selection, and then subjected to dilution plating analysis as described above to test growth sensitivity in the presence of the inducer xylose.

15 The results are shown in Table I below. The first column indicates the Molecule Number of the *Staphylococcus aureus* antisense nucleic acid which was introduced into *Staphylococcus epidermidis*. The second column indicates whether the antisense nucleic acid inhibited the growth of *Staphylococcus epidermidis*, with a "+" indicating that growth was inhibited. Of the 39 *Staphylococcus aureus* antisense
 20 nucleic acids evaluated, 20 inhibited the growth of *Staphylococcus epidermidis*.

TABLE I
Sensitivity of Other Microorganisms to Antisense Nucleic Acids That Inhibit
Proliferation of *Staphylococcus aureus*

Mol. No.	<i>S. epidermidis</i>
SaXA005	+
SaXA007	+
SaXA008	+
SaXA009	+
SaXA010	+

SaXA011	-
SaXA012	-
SaXA013	-
SaXA015	+
SaXA017	-
SaXA022	+
SaXA023	-
SaXA024	-
SaXA025	+
SaXA026	+
SaXA027	-
SaXA027b	-
SaXA02c	-
SaXA028	-
SaXA029	+
SaXA030	+
SaXA032	+
SaXA033	+
SaXA034	-
SaXA035	+
SaXA037	+
SaXA039	-
SaXA042	-
SaXA043	-
SaXA044	-
SaXA045	+
SaXA051	+
SaXA053	-
SaXA056b	-

SaXA059a	+
SaXA060	-
SaXA061	+
SaXA062	+
SaXA063	-
SaXA065	-

It will be appreciated above methods for identifying heterologous proliferation-required genes may be implemented in *Enterococcus faecalis* using the fusion promoters or operators described herein. It will be appreciated that the above methods

5 for identifying heterologous proliferation-required genes using the fusion promoters described herein can be implemented in other Gram-positive organisms, including but not limited to, *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria*

10 *monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* or any species falling within the genera of any of the above species.

Those of skill in the art will appreciate that the methods disclosed herein

15 provide a relatively easy, rapid, and economical way of identifying essential genes in an organism and identifying compounds that alter the expression levels of the identified essential genes.

One skilled in the art will appreciate that these methods and devices are and may be adapted to carry out the objects and obtain the ends and advantages mentioned,

20 as well as those inherent therein. The methods, procedures, and devices described herein are presently representative of preferred embodiments and are exemplary and are not intended as limitations on the scope of the invention. Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention and are defined by the scope of the disclosure.

It will be apparent to one skilled in the art that varying substitutions and modifications may be made to the invention disclosed herein without departing from the scope and spirit of the invention.

Those skilled in the art recognize that the aspects and embodiments of the invention set forth herein may be practiced separate from each other or in conjunction with each other. Therefore, combinations of separate embodiments are within the scope of the invention as disclosed herein.

All patents and publications mentioned in the specification are indicative of the levels of those skilled in the art to which the invention pertains. All patents and publications are herein incorporated by reference to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference.

The invention illustratively described herein suitably may be practiced in the absence of any element or elements, limitation or limitations which is not specifically disclosed herein. Thus, for example, in each instance herein any of the terms “comprising”, “consisting essentially of” and “consisting of” may be replaced with either of the other two terms. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention that in the use of such terms and expressions indicates the exclusion of equivalents of the features shown and described or portions thereof. It is recognized that various modifications are possible within the scope of the invention disclosed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification and variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention as defined by the disclosure.

In addition, where features or aspects of the invention are described in terms of Markush groups, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid comprising a fusion promoter said fusion promoter comprising at least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at least one operator
5 selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *lacIO*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter.
2. The isolated fusion promoter of Claim 1, wherein said at least one promoter is selected from the group consisting of SEQ ID NOs.: 36-45.
- 10 3. The isolated nucleic acid of Claim 1, wherein said at least one operator is *xylO*.
4. The isolated nucleic acid of Claim 3, wherein said at least one promoter is T5.
5. The isolated nucleic acid of Claim 3, further comprising a second
15 operator.
6. The isolated nucleic acid of Claim 5, wherein said second operator is *lacO*.
7. The isolated nucleic acid of Claim 1, wherein said fusion promoter is responsive to an inducer.
- 20 8. The isolated nucleic acid of Claim 7, wherein said inducer is xylose.
9. The isolated nucleic acid of Claim 1, wherein said fusion promoter is titratable.
10. The isolated nucleic acid of Claim 1, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*,
25 *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylosis*, *Streptococcus pneumoniae*, *Streptococcus*
30 *mutans* and any species falling within the genera of any of the above species.

11. The isolated nucleic acid of Claim 1, wherein said at least one gram-positive organism is *Staphylococcus aureus*.

12. The isolated nucleic acid of Claim 1, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

5 13. An isolated nucleic acid comprising a fusion promoter said fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at
10 least one operator represses transcription from said fusion promoter.

14. The isolated nucleic acid of Claim 13, wherein said at least one operator is *xylO*.

15. The isolated nucleic acid of Claim 14, wherein said at least one promoter is T5.

15 16. The isolated nucleic acid of Claim 14, further comprising a second operator.

17. The isolated nucleic acid of Claim 16, wherein said second operator is *lacO*.

18. The isolated nucleic acid of Claim 13, wherein said fusion promoter is
20 responsive to an inducer.

19. The isolated nucleic acid of Claim 18, wherein said inducer is xylose.

20. The isolated nucleic acid of Claim 13, wherein said fusion promoter is titratable.

21. The isolated nucleic acid of Claim 13, wherein said at least one gram-
25 positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus*

epidermidis, *Staphylococcus xylosis*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

22. The isolated nucleic acid of Claim 13, wherein said at least one gram-positive organism is *Staphylococcus aureus*.

5 23. The isolated nucleic acid of Claim 13, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

24. An isolated fusion promoter comprising one of SEQ ID NO.: 26-35.

25. A vector comprising the isolated nucleic acid of Claim 1.

26. A vector comprising the isolated nucleic acid of Claim 13.

10 27. A vector comprising the isolated nucleic acid of Claim 24.

28. The vector of Claim 25 further comprising at least one replicon selected from the group consisting of p15a, pC194 and pCT1138.

29. The vector of Claim 25 further comprising a reporter gene operably linked to said fusion promoter.

15 30. The vector of Claim 29, wherein said reporter gene is *lacL-lacM*.

31. The vector of Claim 30, wherein *lacL-lacM* is derived from *Leuconostoc mesenteroides*.

32. The vector of Claim 25, wherein said at least one operator is *xylO*.

33. The vector of Claim 32, wherein said at least one promoter is T5.

20 34. The vector of Claim 32, further comprising a second operator.

35. The vector of Claim 34, wherein said second operator is *lacO*.

36. The vector of Claim 25, wherein said fusion promoter is responsive to an inducer.

37. The vector of Claim 36, wherein said inducer is xylose.

25 38. The vector of Claim 25, wherein said fusion promoter is titratable.

39. The vector of Claim 25, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*,
30 *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium*

tuberculosis, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

5 40. The vector of Claim 25, wherein said at least one gram-positive organism is *Staphylococcus aureus*.

41. The vector of Claim 25, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

42. The vector of Claim 25, further comprising a random fragment of a microbial genome operably linked to said fusion promoter.

10 43. The vector of Claim 25, further comprising a nucleic acid that encodes a peptide, wherein said nucleic acid is operably linked to said fusion promoter.

44. The vector of Claim 25, further comprising a nucleic acid that is complementary to a portion of a microbial genome, wherein said nucleic acid is operably linked to said fusion promoter.

15 45. The vector of Claim 25, further comprising a nucleic acid that encodes a molecule that inhibits the proliferation of microbe, wherein said nucleic acid is operably linked to said fusion promoter.

46. A host cell comprising the nucleic acid of Claim 1.

47. A host cell comprising the nucleic acid of Claim 13

20 48. A method for identifying genes involved in cellular proliferation said method comprising the steps of:

(a) introducing into cells of a cell population a construct comprising an inducible fusion promoter operably linked to a nucleic acid, said fusion promoter comprising at least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *lacIO*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;

25

(b) inducing transcription of said nucleic acid from said inducible fusion promoter;

(c) identifying the cells in said cell population whose proliferation is reduced in response to the induction of transcription of said nucleic acid; and

5 (d) identifying the gene from a cell identified in step (c) to which at least a portion of said nucleic acid is complementary.

49. A method for identifying genes involved in cellular proliferation said method comprising the steps of:

10 (a) introducing into cells of a cell population a construct comprising an inducible fusion promoter operably linked to a nucleic acid, said fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *lacIO*, wherein said at least one operator is positioned such that binding of at
15 least one repressor to said at least one operator represses transcription from said fusion promoter;

(b) inducing transcription of said nucleic acid from said inducible fusion promoter;

20 (c) identifying the cells in said cell population whose proliferation is reduced in response to the induction of transcription of said nucleic acid; and

(d) identifying the gene from a cell identified in step (c) to which at least a portion of said nucleic acid is complementary.

50. The method of Claim 48, wherein said nucleic acid is random fragment of a microbial genome.

25 51. The method of Claim 48, wherein said nucleic acid encodes an aptamer.

52. The method of Claim 48, wherein said nucleic acid encodes a transcript that is complementary to a portion of a microbial genome.

30 53. The method of Claim 48, wherein said nucleic acid encodes a molecule that inhibits the proliferation of a microbe.

54. The method of Claim 48, wherein said at least one operator is *xylO*.
55. The method of Claim 54, wherein said fusion promoter is T5.
56. The method of Claim 54, wherein said fusion promoter further comprises a second operator.
- 5 57. The method of Claim 56, wherein said second operator is *lacO*.
58. The method of Claim 54, wherein said inducer is xylose.
59. The method of Claim 54, wherein said fusion promoter is titratable.
60. The method of Claim 48, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*,
10 *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.
- 15 61. The method of Claim 48, wherein said at least one gram-positive organism is *Staphylococcus aureus*.
62. The method of Claim 48, wherein said at least one gram-positive organism is *Enterococcus faecalis*.
- 20 63. A method for identifying genes involved in cellular proliferation said method comprising the steps of:
- (a) introducing into the genome of a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid, said fusion promoter comprising at least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at
25 least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *lacIO*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;

(b) comparing the proliferation of said cell cultured in the presence of a first concentration of an inducer that induces transcription from said fusion promoter with the proliferation of said cell cultured in the presence of a concentration of said inducer that is less than said first concentration, wherein a difference in proliferation indicates that said fusion promoter modulates a gene that is required for proliferation; and

(c) identifying the gene that is modulated by said fusion promoter.

64. A method for identifying genes involved in cellular proliferation said method comprising the steps of:

(a) introducing into the genome of a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid, said fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter being linked to at least one operator selected from the group consisting of *xyiO*, *tetO*, *trpO*, *malO* and *lacIO*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;

(b) comparing the proliferation of said cell cultured in the presence of a first concentration of an inducer that induces transcription from said fusion promoter with the proliferation of said cell cultured in the presence of a concentration of said inducer that is less than said first concentration, wherein a difference in proliferation indicates that said fusion promoter modulates a gene that is required for proliferation; and

(c) identifying the gene that is modulated by said fusion promoter.

65. The method of Claim 63, wherein said nucleic acid is random fragment of a microbial genome.

66. The method of Claim 63, wherein said nucleic acid encodes an aptamer.

67. The method of Claim 63, wherein said nucleic acid encodes a transcript that is complementary to a portion of a microbial genome.

68. The method of Claim 63, wherein said nucleic acid encodes a molecule that inhibits the proliferation of a microbe.

69. The method of Claim 63, wherein said at least one operator is *xylO*.

70. The method of Claim 69, wherein said at least one promoter is T5.

5 71. The method of Claim 69, wherein said fusion promoter further comprises a second operator.

72. The method of Claim 71 wherein said second operator is *lacO*.

73. The method of Claim 63, wherein said inducer is xylose.

74. The method of Claim 63, wherein said fusion promoter is titratable.

10 75. The method of Claim 63, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

76. The method of Claim 63, wherein said at least one gram-positive organism is *Staphylococcus aureus*.

20 77. The method of Claim 63, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

78. A method for identifying a compound which reduces the activity or level of a gene product required for proliferation of a cell said method comprising:

25 (a) introducing into a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid that is complementary to at least a portion of a proliferation-required gene, said fusion promoter comprising at least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *λc1O*,
30 wherein said at least one operator is positioned such that binding of at least one

repressor to said at least one operator represses transcription from said fusion promoter;

(b) sensitizing said cell by inducing transcription from said fusion promoter;

5 (c) contacting said sensitized cell with a compound; and

(d) determining the degree to which said compound inhibits proliferation of said sensitized cell relative to a cell which has not been sensitized.

79. A method for identifying a compound which reduces the activity or
10 level of a gene product required for proliferation of a cell said method comprising:

(a) introducing into a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid that is complementary to at least a portion of a proliferation-required gene, said fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59,
15 P1P2 and PL, said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *lacO*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;

(b) sensitizing said cell by inducing transcription from said fusion
20 promoter;

(c) contacting said sensitized cell with a compound; and

(d) determining the degree to which said compound inhibits proliferation of said sensitized cell relative to a cell which has not been sensitized.

25 80. The method of Claim 78, wherein said at least one operator is *xylO*.

81. The method of Claim 80, wherein said at least one promoter is T5.

82. The method of Claim 80, wherein said fusion promoter further comprises a second operator.

83. The method of Claim 82, wherein said second operator is *lacO*.

30 84. The method of Claim 78, wherein said inducer is xylose.

85. The method of Claim 78, wherein said fusion promoter is titratable.

86. The method of Claim 78, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*,
5 *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

10 87. The method of Claim 78, wherein said at least one gram-positive organism is *Staphylococcus aureus*.

88. The method of Claim 78, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

89. A compound identified using the method of Claim 78.

15 90. A method for inhibiting the activity or expression of a gene in an operon required for proliferation said method comprising:

(a) introducing into a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid that is complementary to at least a portion of a proliferation-required operon, said fusion promoter comprising at
20 least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion
25 promoter; and

(b) inducing transcription from said fusion promoter.

91. A method for inhibiting the activity or expression of a gene in an operon required for proliferation said method comprising:

(a) introducing into a cell a construct comprising an inducible fusion
30 promoter operably linked to a nucleic acid that is complementary to at least a

portion of a proliferation-required operon, said fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *lacO*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter; and

(b) inducing transcription from said fusion promoter.

92. The method of Claim 90, wherein said at least one operator is *xylO*.

93. The method of Claim 92, wherein said at least one promoter is T5.

94. The method of Claim 92, wherein said fusion promoter further comprises a second operator.

95. The method of Claim 94, wherein said second operator is *lacO*.

96. The method of Claim 90, wherein said inducer is xylose.

97. The method of Claim 90, wherein said fusion promoter is titratable.

98. The method of Claim 90, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

99. The method of Claim 90, wherein said at least one gram-positive organism is *Staphylococcus aureus*.

100. The method of Claim 90, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

101. A method of manufacturing an antibiotic comprising the steps of:

(a) introducing into a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid that is complementary to at least a portion of a proliferation-required gene, said fusion promoter comprising at

least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;

(b) sensitizing said cell by inducing transcription from said fusion promoter;

(c) contacting said sensitized cell with a compound;

(d) identifying a compound which substantially inhibits the proliferation of said sensitized cell relative to a cell which has not been sensitized; and

(e) manufacturing the compound so identified.

102. A method of manufacturing an antibiotic comprising the steps of:

(a) introducing into a cell a construct comprising an inducible fusion promoter operably linked to a nucleic acid that is complementary to at least a portion of a proliferation-required gene, said fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter;

(b) sensitizing said cell by inducing transcription from said fusion promoter;

(c) contacting said sensitized cell with a compound;

(d) identifying a compound which substantially inhibits the proliferation of said sensitized cell relative to a cell which has not been sensitized; and

(e) manufacturing the compound so identified.

103. The method of Claim 101, wherein said at least one operator is *xylO*.

104. The method of Claim 103, wherein said at least one promoter is T5.

105. The method of Claim 103, wherein said fusion promoter further comprises a second operator.

106. The method of Claim 105, wherein said second operator is *lacO*.

5 107. The method of Claim 101, wherein said inducer is xylose.

108. The method of Claim 101, wherein said fusion promoter is titratable.

109. The method of Claim 101, wherein said at least one gram-positive organism is selected from the group consisting of *Bacillus anthracis*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*,
10 *Corynebacterium diphtheriae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Lactococcus lactis*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Nocardia asteroides*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus xylois*, *Streptococcus pneumoniae*, *Streptococcus mutans* and any species falling within the genera of any of the above species.

15 110. The method of Claim 101, wherein said at least one gram-positive organism is *Staphylococcus aureus*.

111. The method of Claim 101, wherein said at least one gram-positive organism is *Enterococcus faecalis*.

112. A method for identifying a gene which is required for proliferation of a
20 prokaryotic cell said method comprising the steps of:

(a) replacing the native promoter of a gene in the chromosome of a prokaryotic cell having an enhanced frequency of homologous recombination with a regulatable fusion promoter comprising at least one promoter that is modified to have altered activity in at least one gram-positive organism said
25 promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO*, *lacO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter; and

(b) identifying cells in which the extent of proliferation of said cell
30 when said fusion promoter is active at a first level is substantially different

than the extent of proliferation of said cell when said fusion promoter is active at a second level, said first level being greater than said second level.

113. A method for identifying a gene which is required for proliferation of a prokaryotic cell said method comprising the steps of:

5 (a) replacing the native promoter of a gene in the chromosome of a prokaryotic cell having an enhanced frequency of homologous recombination with a regulatable fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter being linked to at least one operator selected from the group consisting of
10 *xylO*, *tetO*, *trpO*, *malO*, *lacO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter; and

(b) identifying cells in which the extent of proliferation of said cell when said fusion promoter is active at a first level is substantially different
15 than the extent of proliferation of said cell when said fusion promoter is active at a second level, said first level being greater than said second level.

114. The method of Claim 112, wherein said fusion promoter is an inducible promoter.

115. The method of Claim 112, wherein the step of replacing said native
20 promoter comprises introducing a linear nucleic acid comprising a 5' portion homologous to a first portion of said native promoter, a 3' portion homologous to a second portion of said native promoter and said fusion promoter disposed between said 5' portion and said 3' portion into said cell such that homologous recombination occurs between said 5' portion and said first portion of said native promoter and
25 between said 3' portion and said second portion of said native promoter.

116. The method of Claim 115, wherein said linear nucleic acid is double stranded.

117. The method of Claim 115, wherein said linear nucleic acid is single stranded.

118. A method for identifying a compound which inhibits the proliferation of a prokaryotic cell said method comprising the steps of:

5 (a) replacing the native promoter of a gene in the chromosome of a prokaryotic cell having an enhanced frequency of homologous recombination with a regulatable fusion promoter comprising at least one promoter that is modified to have altered activity in at least one gram-positive organism said promoter being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO*, *lacO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least
10 one operator represses transcription from said fusion promoter; and

(b) comparing the extent of proliferation of a first sample of said cell in the presence of said compound to the extent of proliferation of a second sample of said cell in the presence of said compound, wherein said first sample of said cell has a reduced activity of said fusion promoter activity relative the
15 activity of said fusion promoter in said second sample of said cell and wherein said compound inhibits the proliferation of said cell if the extent of proliferation of said first sample of said cell is substantially less than the extent of proliferation of said second sample of said cell.

119. A method for identifying a compound which inhibits the proliferation
20 of a prokaryotic cell said method comprising the steps of:

(a) replacing the native promoter of a gene in the chromosome of a prokaryotic cell having an enhanced frequency of homologous recombination with a regulatable fusion promoter comprising at least one promoter selected from the group consisting of T5, CP25, P32, P59, P1P2 and PL, said promoter
25 being linked to at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO*, *lacO* and *λc1O*, wherein said at least one operator is positioned such that binding of at least one repressor to said at least one operator represses transcription from said fusion promoter; and

(b) comparing the extent of proliferation of a first sample of said cell in
30 the presence of said compound to the extent of proliferation of a second

sample of said cell in the presence of said compound, wherein said first sample of said cell has a reduced activity of said fusion promoter activity relative the activity of said fusion promoter in said second sample of said cell and wherein said compound inhibits the proliferation of said cell if the extent of proliferation of said first sample of said cell is substantially less than the extent of proliferation of said second sample of said cell..

120. The method of Claim 118, wherein the step of replacing said native promoter comprises introducing a linear nucleic acid comprising a 5' portion homologous to a first portion of said native promoter, a 3' portion homologous to a second portion of said native promoter and said fusion promoter disposed between said 5' portion and said 3' portion into said cell such that homologous recombination occurs between said 5' portion and said first portion of said native promoter and between said 3' portion and said second portion of said native promoter.

121. The method of Claim 120, wherein said linear nucleic acid is double stranded.

122. The method of Claim 120, wherein said linear nucleic acid is single stranded.

123. A method for identifying a gene which is required for proliferation of a prokaryotic cell said method comprising the steps of:

(a) introducing at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO*, *lacO* and *λc1O* into a prokaryotic cell having an enhanced frequency of homologous recombination such that said at least one operator regulates transcription of a target nucleic acid in the chromosome of said cell; and

(b) identifying cells in which the extent of proliferation of said cell when said target nucleic acid is transcribed at a first level is substantially different than the extent of proliferation of said cell when said target nucleic acid is transcribed at a second level, said first level being greater than said second level.

124. The method of Claim 123, wherein the step of introducing said at least one operator comprises introducing a linear nucleic acid comprising a 5' portion homologous to a first portion of the chromosome of said cell, a 3' portion homologous to a second portion of the chromosome of said cell and said at least one operator
5 disposed between said 5' portion and said 3' portion into said cell such that homologous recombination occurs between the 5' portion and said first portion of the chromosome of said cell and between the 3' portion and said second portion of the chromosome of said cell.

125. The method of Claim 124, wherein said linear nucleic acid is double
10 stranded.

126. The method of Claim 124, wherein said linear nucleic acid is single stranded.

127. A method of identifying a compound which inhibits the proliferation of a prokaryotic cell said method comprising the steps of:

15 (a) obtaining a prokaryotic cell in which transcription of a nucleic acid required for proliferation of said cell is regulated by at least one operator which has been introduced into the chromosome of said cell said at least one operator selected from the group consisting of *xylO*, *tetO*, *trpO*, *malO*, *lacO* and *λcI O*;

20 (b) sensitizing said cell by growing said cell under conditions in which the level of transcription of said gene is lower than that of a wild type cell;

(c) contacting said sensitized cell with said compound; and

(d) determining the degree to which said compound inhibits the growth of said sensitized cell relative to an unsensitized cell.

25 128. The method of Claim 127, wherein said cell has an enhanced frequency of homologous recombination.

129. The method of Claim 128, wherein said linear nucleic acid is single stranded.

30 130. The method of Claim 128, wherein said linear nucleic acid is double stranded.

131. A method of identifying a nucleic acid sequence having promoter activity in *Enterococcus faecalis* said method comprising the steps of:

5 (a) inserting a candidate nucleic acid into a vector comprising *lacL-lacM* reporter genes such that said candidate nucleic acid is upstream of the *lacL-lacM* reporter genes;

(b) introducing said vector comprising said candidate nucleic acid into *Enterococcus faecalis*;

10 (c) detecting expression of the *lacL-lacM* reporter genes, wherein expression of said *lacL-lacM* reporter genes indicates that said candidate nucleic acid sequence has promoter activity.

132. The method of Claim 131, wherein said detecting step comprises measuring β -galactosidase activity.

133. The method of Claim 131, wherein said candidate nucleic acid is a promoter modified to increase activity in a gram-positive organism.

15 134. The method of Claim 131, wherein said *lacL-lacM* reporter gene is derived from *Leuconostoc mesenteroides*.

135. The method of Claim 131, wherein said vector is pEPEF1.

20

1/18

GACCGTCATAAAAAATTTATTTGCTTTCAGGAAAATTTTCTGTATAATAGATTC
1 2 3 4

AAGTTAGTTTGTATTATTAATAACCAACTAAAATGTAGAATTCCGAGCTCGAG
5 6

CCGGTACCCGGGGATCCTCTAGAGTCGACCTGCAGCCAAGCTTGGGCTTTTCA
6

GCCTGATACAGATTAAATCAGAACGCAGAAGCGGTCTGATAAAACAGAATTG

CCTGGCGGCAGTAGCGCGGTGGTCCCACCTGACCCCATGCCGAACTCAGAAGT

GAAACGCCGTAGCGCCGATGGTAGTGTGGGGTCTCCCCATGCGAGAGTAGGGA

ACTGCCAGGCATCAAATAAAACGAAAGGCTCAGTCGAAAGACTGGGCCTTTTCG
7A

TTTTATCTGTTGTTTGTCTGGTGAACGCTCTCCTGAGTAGGACAAATCCGCCGGG

AGCGGATTTGAACGTTGCGAAGCAACGGCCCGGAGGGTGGCGGGCAGGACGC

CCGCCATAAACTGCCAGGCATCAAATTAAGCAGAAGGCCATCCTGACGGATGG
7B

CCTTTTTGCGTTTCTACAAACTC

FIG. 1

2/18

ATTAAATATAAATAACATTCAAATGAATATTAAATAATTACAGCAGTCTGAG
1
TTATAAAATAGATATCTCGGACCGACATTAAATTCTTGACAGGGAGAGATAG
a 2 3
GTTTGATAGAAATATAATAGTTGTCTCGAGAGTTAGTTTGTTTATTAAATTAAC
4 b 5
CAACTGGACGA

FIG. 2

3/18

 SEQUENCE HOMOLOGOUS TO TARGET

 HOMOLOGOUS CHROMOSOMAL SEQUENCE

P = PROMOTER

DRUG RESISTANCE = GENE ENCODING GENE PRODUCT WHICH
CONFERS DRUG RESISTANCE

P_T = PROMOTER OF TARGET GENE

T = TRANSCRIPTIONAL TERMINATOR

P_R = REGULATABLE PROMOTER




FIG. 3A

4/18

 = SEQUENCE HOMOLOGOUS TO TARGET

 = CHROMOSOMAL TARGET

P_R = REGULATABLE TARGET

P_T PROMOTER OF TARGET GENE




FIG. 3B

5/18




FIG. 4A

6/18

 = SEQUENCE HOMOLOGOUS TO TARGET

 = CHROMOSOMAL TARGET

O = OPERATOR

P_T = CHROMOSOMAL PROMOTER OF TARGET GENE




FIG. 4B





FIG. 5

8/18






FIG. 7

10/18




FIG. 8




FIG. 9

12/18

Xylose Titrations With P59 Promoter in
Enterococcus faecalis




FIG. 10

13/18




FIG. 11




FIG. 12




FIG. 13A

16/18




FIG. 13B




FIG. 14

The Selective Sensitization to an Antibiotic Inhibiting Gyrase B Subunit Activity Following the Induction of an Antisense Construct to the B Subunit of Gyrase.




FIG. 15

SEQUENCE LISTING

<110> ELITRA PHARMACEUTICALS, INC.

HASELBECK, Robert

WALL, Daniel

GROSS, Molly

<120> BACTERIAL PROMOTERS AND METHODS OF USE

<130> ELITRA.010VPC

<150> US 60/259,434

<151> 2000-12-27

<150> US 09/948,993

<151> 2001-09-06

<160> 68

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 556

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequence map showing the Xyl-T5 fusion promoter
and associated DNA sequences

<400> 1

```

gaccgtcata aaaaatttat ttgctttcag gaaaattttt ctgtataata gattcaagtt 60
agtttggtta ttaaattaac caactaaaat gtagaattcc gagctcgagc cggtagcccg 120
ggatcctcta gagtcgacct gcagccaagc ttgggctttt cagcctgata cagattaaat 180
cagaacgcag aagcgggtctg ataaaacaga atttgccctgg cggcagtagc gcggtgggtcc 240
cacctgacct catgccgaac tcagaagtga aacgccgtag cggcgatggg agtgtggggg 300
ctcccatgag gagagtaggg aactgccagg catcaaataa aacgaaaggc tcagtcgaaa 360
gactgggcct ttggttttat ctgttggttg tcggtgaacg ctctcctgag taggacaaat 420
ccgcccggag cggatttgaa cgttgccaag caacggcccg gaggggtggc ggcaggacgc 480
ccgcataaaa ctgccaggca tcaaattaag cagaaggcca tcctgacgga tggccttttt 540
gcgtttctac aaactc                                     556

```

<210> 2

<211> 168

<212> DNA

<213> Artificial Sequence

<220>

<223> CP25 fusion promoter

<400> 2

```

attaatata aataacattc aaatgaatat taaataatta cagcagtcct agttataaaa 60
tagatatctc ggaccgacat taaattcttg acaggagag ataggtttga tagaatataa 120
tagttgtctc gagagttagt ttgtttatta aattaaccaa ctggacga          168

```

<210> 3

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer

<400> 3

tggcatttcta cgactataac

20

<210> 4

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer

<400> 4

gcgaatttcta catttttagtt ggttaattt

29

<210> 5

<211> 94

<212> DNA

<213> Escherichia coli

<220>

<223> Oligonucleotide Xyl-T5

<400> 5

gaccgtcata aaaaatttat ttgctttcag gaaaattttt ctgtataata gattcaagtt 60
agtttgttta ttaaattaac caactaaaat gtag 94

<210> 6

<211> 95

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide Xyl-T5 complement

<400> 6

aattctacat tttagttggt taatttaata aacaaactaa cttgaatcta ttatacagaa 60
aaattttcct gaaagcaaat aaatttttta tgacg 95

<210> 7

<211> 10929

<212> DNA

<213> Artificial Sequence

<220>

<223> Vector pEP25

<400> 7

aagcttgcct gcaggtcgac gataaccaca taacagtcac aaaaccactc ctttttaaca 60
aacttttatca caagaaatat tttggcattc tacgactata acttaaaattt atatttttta 120
ctttataata tataattgat tatagaataa tggtgtcat atcgtttgcc aacatctagt 180
actcaaatta cactatgtta cacttggtta tattaaccga acttcccctg tccaaattag 240
ataagaggta ataataaatg gaaaataatt ttatagtaaa tgaaaatgag aagcgtgtat 300
taaaacaaat tttcaataac agcaatattt cacgaacaca aatatcgaag aatttagaac 360
ttaataaagc tactatttct aacattctga acaacttaaa acacaagagt ttagttaatg 420
aagtaggaga aggtaatagt actaaaagt gtaggacgaa gcctatttta ctcgaaatta 480
acaaaaata tggctactat atttctatgg atttaacata tgattccgtt gaattaatgt 540
acaactactt tgatgctact atattaaagc aagattccta cgaattaaat gataaaaatg 600

taagcagtat	attacaaatt	ttaaaatcta	atataaacgt	ctcagaaaaa	tatgatacgt	660
tatatgggtt	acttgggtata	tctatatcca	tacacgggtat	cggttgacgat	gagcaaaaaca	720
taatcaatct	tccttttcat	aaaaatgaga	aacgcacatt	taccgatgaa	ttaaagtcac	780
tcacaaatgt	tcctgtcgtt	atagaaaatg	aagcaaattt	atcagcgcta	tatgaaaaaa	840
gtttatatat	taattcaaac	ataaataatt	tgattacttt	aagtattcac	aaggggtatag	900
gcgctggcat	cataataaat	aaaaaacttt	atcgtggctc	aaatggagag	gctggagaga	960
taggtaagac	attgggtttg	gaatctataa	ataacaatga	caacaaatat	tataaaatcg	1020
aagatatatg	ctcccaagac	gctttaatac	agaaaaataa	taatagggtg	ggcgctcacat	1080
tgacgtttac	agaactaatc	caatattaca	acgaaggaaa	ttcaattgtt	gctcatgaaa	1140
ttaacaatt	tattaataaa	atgacagttc	tgattcataa	tttgaatata	caatttaacc	1200
cagacgctat	ttatattaac	tgctctttaa	ttaatgaatt	accaaattat	ttaaatgaaa	1260
ttaaagagca	attctcctgt	ttttctcaag	gcagtcacat	tcaattacat	ttaactacta	1320
atgtaaaaca	agctacttta	ttgggtggca	ctttagcaat	aatgcaaaaa	acattaaata	1380
taaataacat	tcaaataaat	attaaataat	tacagcagtc	tgagttataa	aatagatatc	1440
tcggaccgct	ttggcagttt	attcttgaca	tgtagtgagg	gggctgggtat	aatcacatag	1500
ctcgagagtt	agtttgttta	ttaaattaac	caactggatc	cggaggaaca	gctatgacca	1560
tgattacgga	ttcactggcc	gtcgttttac	aacgtcgtga	ctgggaaaaac	cctggcggtta	1620
cccaacttaa	tcgccttgca	gcacatcccc	ctttcgccag	ctggcgtaat	agcgaagagg	1680
cccgaccga	tcgcccctcc	caacagttgc	gcagcctgaa	tggcgaatgg	cgctttgcct	1740
ggtttccggc	accagaagcg	gtgccggaaa	gctggctgga	gtgcgatctt	cctgaggccg	1800
atactgtcgt	cgccccctca	aactggcaga	tgacaggtta	cgatgcgccc	atctacacca	1860
acgtgacctt	ccccattacg	gtcaatccgc	cgtttgttcc	cacggagaat	ccgacgggtt	1920
gttactcgct	cacttttaac	gttgatgaaa	cttggtctaca	ggaaggccag	acgcgaatta	1980
tttttgatgg	cgtaactcgc	gcgtttcatc	tggtgtgcaa	cgggcgctgg	gtcggttacg	2040
gccaggacag	tcgtttgccc	tctgaatttg	acctgagcgc	atttttacgc	gccggagaaa	2100
accgcctcgc	ggtgatgggt	ctgcgctgga	gtgacggcag	ttatctggaa	gatcaggata	2160
tggtggcgat	gagcggcatt	ttccgtgacg	tctcgttgct	gcataaaacc	actacacaaa	2220
tcagcgattt	ccatggttgc	actcgtctta	atgatgattt	cagccgcgct	gtactggagg	2280
ctgaagttca	gatgtgcggc	gagttgcgtg	actacctacg	ggtaacagtt	tctttatggc	2340
aggtgaaac	gcaggtcgcc	agcggcaccg	cgcttttcgg	cggtgaaatt	atcgatgagc	2400
gtggtggtta	tgccgatcgc	gtcacactac	gtctgaacgt	cgaaaaaccg	aaactgtgga	2460
gcgcgaaat	cccgaatctc	tatcgtgcgg	tggttgaact	gcacaccgcc	gacggcacgc	2520
tgattgaagc	agaagcctgc	gatgtcgggt	tccgcgaggt	gcgattgaa	aatggtctgc	2580
tgctgctgaa	cggaagccg	ttgctgatcc	gaggcggtta	ccgtcacgag	catcatcctc	2640
tgcatgggtca	ggtcatggat	gagcagacga	tggtgcagga	tatcctgctg	atgaagcaga	2700
acaactttta	cgccgtgcgc	tgcttcgcat	atccgaacca	tccgctgtgg	tacacgctgt	2760
gcgaccgcta	cggcctgtat	gtggtggatg	aagccaatat	tgaaccacac	ggcatgggtc	2820
caatgaatcg	tctgaccgat	gatccgcgct	ggctaccggc	gatgagcgaa	cgcgtaacgc	2880
gaatggtgca	gcgcgatcgt	aatcacccga	gtgtgatcat	ctggtcgtcg	gggaatgaat	2940
caggccacgg	cgctaatac	gacgcgctgt	atcgctggat	caaatctgtc	gatccttccc	3000
gcccggtgca	gtatgaaggc	ggcggagccg	acaccacggc	caccgatatt	atttgcccga	3060
tgtagcgcgc	cgtggatgaa	gaccagccct	tcccggctgt	gccgaaatgg	tccatcaaaa	3120
aatggtcttc	gctacctgga	gagacgcgcc	cgctgatcct	ttgcgaatac	gcccacgcga	3180
tggttaaacg	tcttgccggt	ttcgctaaat	actggcaggc	gtttcgtcag	tatccccgtt	3240
tacagggcgg	cttcgtctcg	gactgggtgg	atcagtcgct	gattaaatat	gatgaaaacg	3300
gcaaccgctg	gtcggcttac	ggcggtgatt	ttggcgatac	gccgaacgat	cgccagttct	3360
gtatgaacgg	tctggtcttt	gccgaccgca	cgccgcaccc	agcgtgacg	gaagcaaaac	3420
accagcagca	gtttttccag	ttccgtttat	ccgggcaaac	catcgaaagt	accagcgaat	3480
acctgttccg	tcatagcgat	aacgagctcc	tgactgggat	ggtggcgctg	gatggtaagc	3540
cgctggcaag	cggtgaagtg	cctctggatg	tcgctccaca	aggtaaacag	ttgattgaac	3600
tgcttgaact	accgcagccg	gagagcgccg	ggcaactctg	gctcacagta	cgcgtagtgc	3660
aaccgaacgc	gaccgatggg	tcagaagccg	ggcacatcag	cgctggcag	cagtggcgct	3720
tggcggaaaa	cctcagtggt	acgctcccg	ccgcgtccca	cgccatcccg	catctgacca	3780
ccagcgaaat	ggatttttgc	atcgagctgg	gtaataagcg	ttggcaattt	aaccgccagt	3840
caggctttct	ttcacagatg	tggtttggcg	ataaaaaaca	actgctgacg	ccgctgcgcg	3900
atcagttcac	ccgtgcaccg	ctggataacg	acattggcgt	aagtgaagcg	acccgcattg	3960
accctaacgc	ctgggtcgaa	cgctgggaag	cgccgggcca	ttaccaggcc	gaagcagcgt	4020
tggtgcagtg	cacggcagat	acacttgctg	atgcggtgct	gattacgacc	gctcacgcgt	4080
ggcagcatca	ggggaaaacc	ttatttatca	gccggaaaac	ctaccggatt	gatggtagt	4140
gtcaaatggc	gattaccgtt	gatgttgaag	tggcgagcga	tacaccgat	ccggcgcgga	4200
ttggcctgaa	ctgccagctg	gcgcaggtag	cagagcgggt	aaactggctc	ggattagggc	4260

```

cgcaagaaaa ctatcccgcg cgccttactg ccgcctgttt tgaccgctgg gatctgccat 4320
tgtcagacat gtataccccg tacgtcttcc cgagcgaaaa cggctctgcg tgcgggacgc 4380
gcgaattgaa ttatggccca caccagtggc gcggcgactt ccagttcaac atcagccgct 4440
acagtcaaca gcaactgatg gaaaccagcc atcgccatct gctgcacgcg gaagaaggca 4500
catggctgaa tatcgacggt ttccatatgg ggatttgggtg cgacgactcc tggagcccgt 4560
cagtatcggc ggaattccag ctgagcgccg gtcgctacca ttaccagtgt gtctgggtgc 4620
aaaaataata atgtcgaggg taccgagctc gaattccttt tagtccagcc gagcgcttag 4680
tggaatttg taccctttat cgatacaaat tccccgtagg cgctagggac ctcttttagct 4740
ccttggaagc tgtcagtagt atacctaata atttatctac attcccttaa gtaacgtgta 4800
actttccaaa ttacaaaaag cgactcatag aattatttcc tcccgtaaaa taatagataa 4860
ctattaaaaa tagacaatac ttgctcataa gtaacggtac ttaaattgtt tactttggcg 4920
tgtttcattg cttgatgaaa ctgattttta gtaaacagtt gacgatattc tcgattgacc 4980
cattttgaaa caaagtacgt atatagcttc caatatttat ctggaacatc tgtggtagtg 5040
cgggtaagtt ttattaagac actgtttact tttggtttag gatgaaagca ttccgctggc 5100
agcttaagca attgctgaat cgagacttga gtgtgcaaga gcaaccctag tgttcgggtg 5160
atatccaagg tacgcttgta gaatccttct tcaacaatca gatagatgtc agacgcagtg 5220
ctttcaaaaa ccactttttt aataatttgt gtgcttaaat ggtaaggaat actcccaaca 5280
attttatacc tctgtttgtt aggggaattga aactgtagaa tatcttgggtg aattaaagt 5340
acacgagtat tcagttttta tttttctgac gataagttga atagatgact gtctaattca 5400
atagacgtta cctgtttact tatttttagcc agtttcgctg ttaaagccc tttacctgtt 5460
ccaatttcgt aaacgggtatc ggtttccttt aaattcaatt gttttattat ttggttgagt 5520
actttttcac tcgttaaaaa gttttgagaa tattttata tttgttcat gtaatcactc 5580
cttcttaatt acaatttttt agcatctaat ttaacttcaa ttctattat acaaaatttt 5640
aagatactgc actatcaaca cactcttaag tttgcttcta agtcttattt ccataacttc 5700
ttttacgttt ccgccattct ttgctgtttc gatttttatg atatggtgca agtcagcacg 5760
aacacgaacc gtcttatatc ccattatata ttttttgca ctgattgggtg tatcatttcg 5820
tttttctttt tgtgcgccta aatttcccac aatcactcac ttctttctat ttcttcttat 5880
tcttatttta tcatcaacaa tcacaaatca cttgtgattt gtgataagtg atttgtgatt 5940
aatatataaa agccctcttt aaagggtctt tatgtttatt ttgagaaaga tataaaatca 6000
atatatccct tttctccgat ttttacaacg gcattgtagg actttctatc ttctgttttg 6060
attcctttta ccagggtttc ttttccctct agtaattctt ttacatttgt tttggtagtg 6120
tttttctttc taaaatgttc agctaaagta aacttacatt caggataatt tgaacaacca 6180
taaaacgatt tttttaatac aatattgttg ccacacttag gacattttcc tacaataact 6240
ttttctgctt ctttttcttt ctgttccctg taatcagaaa aatttagttt ttctatatcg 6300
ttaggtagag cttccagtaa atgaacaatg aattttttga ttttcgtaat aaagttctct 6360
tgattgcctt ctcttttacc gatttttttt aaatacgttt cccatttagc cgtcatttca 6420
gcactcgta aaaggtgctg actttcaact gcctggcaca ataattttcc tttttcagtt 6480
acaacaagct ctgatttcac tttttgcatt ctacaaactg cataactcat atgtaaatcg 6540
ctccttttta ggtggcacia atgtgaggca ttttcgctct ttccggcgag gctagttacc 6600
cttaagttat tggtagact ggttttaagc gcaaaaaaag ttgctttttc gtacctatta 6660
atgtatcggt agaaaaacga ctgtaaaaaa tagctcggc attatctcat attataaaag 6720
ccagtcatta ggcctatctg acaattcctg aatagagttc ataaacaatc ctgcatgata 6780
accatcacia acagaatgat gtacctgtaa agatagcggg aaatatattg aattaccttt 6840
attaatgaat tttctgctg taataatggg tagaaggtaa ttactattat tattgatatt 6900
taagttaaac ccagtaaatg aagtcctatg aataatagaa agagaaaaag cattttcagg 6960
tataggtgtt ttgggaaaca atttccccga accattatat ttctctacat cagaaaggta 7020
taaatcataa aactctttga agtcattctt tacaggagtc caaataccag agaattgttt 7080
agatacacca tcaaaaattg tataaagtgg ctctaactta tccaataaac ctaactctcc 7140
gtcgctattg taaccagttc taaaagctgt atttgagttt atcaccttg tcactaagaa 7200
aataaatgca gggtaaaatt tatatccttc ttgttttatg tttcgggtata aaacactaat 7260
atcaatttct gtgggtatac taaaagtcgt ttgggtggtc aaataatgat taaatatctc 7320
ttttctcttc caattgtcta aatcaatttt attaaagttc atttgatatg cctcctaaat 7380
ttttatctaa agtgaattta ggaggcttac ttgtctgctt tcttcattag aatcaatcct 7440
tttttaaaag tcaatattat tgtaacataa atatatattt taaaaatata ccactttatc 7500
caattttcgt ttgttgaact aatgggtgct ttagttgaag aataaaaagac cacattaaaa 7560
aatgtggtct tttgtgtttt tttaaaggat ttgagcgtag cgaaaaatcc ttttctttct 7620
tatcttgata ataagggtaa ctattgccgg gatagactgt aacattctca cgcataaaat 7680
cccctttcat tttctaattg aaatctatta ccttattatt aattcaattc gctcataaatt 7740
aatccttttt cttattacgc aaaatggccc gatttaagca caccctttat tccgttaatg 7800
cgccatgaca gccatgataa ttactaatac taggagaagt taataaatat gtaaccaaca 7860
tgattaacaa ttattagagg tcatcgttca aaatggtatg cgttttgaca catccactat 7920

```

```

atatccgtgt cgttctgtcc actcctgaat cccattccag aaattctcta gcgattccag 7980
aagtttctca gagtgcgaaa gttgaccaga cattacgaac tggcacagat ggtcataacc 8040
tgaaggaaga tctgattgct taactgcttc agttaagacc gaagcgctcg tcgtataaca 8100
gatgcgatga tgcagaccaa tcaacatggc acctgccatt gctacctgta cagtcaagga 8160
tggtagaaat gttgtcggtc cttgcacacg aatattacgc catttgccctg catattcaaa 8220
cagctcttct acgataaggg cacaatcgc atcgtggaac gtttgggctt ctaccgattt 8280
agcagtttga tacactttct ctaagtatcc acctgaatca taaatcggca aaatagagaa 8340
aaattgacca tgtgtaagcg gccaatctga ttccacctga gatgcataat ctagtagaat 8400
ctcttcgcta tcaaaattca cttccacctt ccactcaccg gttgtccatt catggctgaa 8460
ctctgcttcc tctgttgaca tgacacacat catctcaata tccgaatagg gcccatcagt 8520
ctgacgacca agagagccat aaacaccaat agccttaaca tcatccccat atttatccaa 8580
tattcgttcc ttaatttcat gaacaatctt cattctttct tctctagtca ttattattgg 8640
tccattcact attctcattc ctttttcaga taattttaga tttgcttttc taaataagaa 8700
tatttgaga gcaccgttct tattcagcta ttaataactc gtcttcctaa gcatccttca 8760
atccttttaa taacaattat agcatctaata cttcaacaaa ctggcccggt tgttgaaacta 8820
ctcctttaata aaataatttt tccgttccca attccacatt gcaataatag aaaaatccatc 8880
ttcatcggtt ttttcgtcat catctgtatg aatcaaatcg cttcttcttg tgtcatcaag 8940
gtttaatttt ttatgtattt cttttaacaa accaccatag gagattaacc ttttacggtg 9000
taaaccttcc tccaaatcag acaaacgttt caaattcttt tcttcatcat cggtcataaa 9060
atccgtatcc tttacaggat attttgcagt ttctgcaatt gccgattgta tatccgattt 9120
atatttattt ttcggtcgat tttttattaa aacgtctcaa aatcgtttct gagacgtttt 9180
agcgtttatt tcgtttagtt atcggcataa tcgttaaaac aggcgttatc gtagcgtaaa 9240
agccttgag cgttagcgtg ctttgacgag aagatgttgt ctglttagatt atgaaagccg 9300
atgactgaat gaaataataa gcgcagcgtc cttctatttc gggttgagga ggctcaagg 9360
agtttgagg aatgaaattc cctcatgggt ttgattttaa aaattgcttg caattttgcc 9420
gagcgttagc gctggaat ttttgaaaaa aatttggaaat ttggaaaaaa atggggggga 9480
aagggaagcg attttgcttc cgtactacga cccccatta agtgccgagt gccaattttt 9540
gtgccaaaaa cgctctatcc caactggctc aagggtttga ggggtttttc aatcgccaac 9600
gaatcgccaa cgttttcgcc aacgtttttt ataaatctat atttaagtag ctttatttgt 9660
gtttttatga ttacaaagtga atacactaat tttataaaat tatttgattg gagtttttta 9720
aatggttgatt tcagaatcga aaaaaagagt tatgatttct ctgacaaaag agcaagataa 9780
aaaattaaca gatatggcga aacaaaaagg tttttcaaaa tctgcgggtt gcgggcgtta 9840
gctatagaag aatatgcaag aaaggaatca gaacaaaaaa aataagcgaa agctcgcggt 9900
tttagaagga tacgagtttt cgctacttgt ttttgataag gtaattatat catggctatt 9960
aaaaatacta aagctagaaa ttttgattt ttattatct ctgactcaat tcctaatgat 10020
tggaagaaaa aattagagag tttgggcgta tctatggctg tcagtccttt acacgatatg 10080
gacgaaaaaa aagataaaga tacatggaat agtagtgatg ttatacgaaa tggaaagcac 10140
tataaaaaac cacactatca cgttatatat attgcacgaa atcctgtaac aatagaaagc 10200
gttaggaaca agattaagcg aaaattgggg aatagttcag ttgctcatgt tgagatactt 10260
gattatatca aaggttcata tgaatatttg actcatgaat caaaggacgc tattgctaag 10320
aataaacata tatacgacaa aaaagatatt ttgaacatta atgattttga tattgaccgc 10380
tatataacac ttgatgaaag ccaaaaaaga gaattgaaga atttactttt agatatagtg 10440
gatgactata atttggtaaa taaaaaagat ttaatggctt ttattcgctt taggggagcg 10500
gagtttgga ttttaaatc gaatgatgta aaagatattg tttcaacaaa ctctagcgcc 10560
tttagattat ggtttgaggg caattatcag tgtggatata gagcaagtta tgcaaagggt 10620
cttgatgctg aaacggggga aataaaatga caaacaaaaga aaaagagtta tttgctgaaa 10680
atgaggaatt aaaaaaagaa attaaaggact taaaagagcg tattgaaaga tacagagaaa 10740
tggaagtga attaaagtaca acaatagatt tattgagagg agggattatt gaataaataa 10800
aagccccct gacgaaagtc gaagggggct tttatttttg tttgatgttg cgattaatag 10860
caatacgatt gcaataaaca aaatgatcga tgaaagcaat gtgagagaaa aatatgttcc 10920
taaaacaac 10929

```

<210> 8

<211> 12739

<212> DNA

<213> Artificial Sequence

<220>

<223> Vector pEPEF1

<400> 8

```

aagcttgccg gcaggcgacg ataaccacat aacagtcata aaaccactcc tttttaacaa 60
actttatcac aagaaatatt ttggcattct acgactataa cttaaattta ttttttttac 120
tttataatat ataattgatt atagaataat gttgctcata tcgtttgcca acatctagta 180
ctcaaattac actatgttac acttggtaat attaacgaa cttccctgt ccaaattaga 240
taagaggtaa taataaatgg aaaataatth ttagttaaag gaaaatgaga agcgtgtatt 300
aaaacaaatt ttcaataaca gcaatatttc acgaacacaa atatcgaaga atttagaact 360
taataaagct actatttcta acattctgaa caacttaaaa cacaagagtt tagttaatga 420
agtaggagaa ggtaatagta ctaaaagtgg tggacgaaag cctattttac tcgaaattaa 480
ccaaaaatat ggctactata tttctatgga tttaacatat gattccgttg aattaatgta 540
caactacttt gatgctacta tattaagca agattcctac gaattaaatg ataaaaatgt 600
aagcagtata ttacaaatth taaaatctaa tataaacgtc tcagaaaaat atgatacgtt 660
atatgggtta ctgggtatat ctatatccat acacgggtatc gttgacgatg agcaaaacat 720
aatcaatctt ccttttcata aaaatgagaa acgcacattt accgatgaat taaagtcatt 780
cacaaatggt cctgtcgtta tagaaaatga agcaaattha tcagcgtat atgaaaaaag 840
tttatatatt aattcaaaca taaataatth gattacttha agtattcaca aggggtatagg 900
cgctggcatc ataataaata aaaaacttha tcgtggctca aatggagagg ctggagagat 960
aggtagaca ttggttttgg aatctataaa taacaatgac aacaaatatt ataaaaatga 1020
agatatagtc tcccaagacg ctttaataca gaaaataaat aatagggttg gcgtcacatt 1080
gacgtttaca gaactaatcc aatattacaa cgaaggaaat tcaattgttg ctcatgaaat 1140
taaacaatth attataaaaa tgacagttct gattcataat ttgaatacac aatttaaccc 1200
agacgctatt tatattaact gtcctttaat taatgaatta ccaaatattt taaatgaaat 1260
taaagagcaa ttctcctggt tttctcaagg cagtcacatt caattacatt taactactaa 1320
tgtaaaacaa gctactttat tgggtggcac tttagcaata atgcaaaaaa cattaaatat 1380
aaataacatt caaatgaata ttaaataatt acagcagtc gagttataaa atagatatct 1440
cggaccgctt tggcagttta ttcttgacat gtagtgagg ggctgggata atcacatagt 1500
actgttctcg agagttagtt tgtttattaa attaaccaac tggatcccg gtctagatta 1560
gggtaaactt gaaaggatat tctcatgca agctaattct caatgggttag atgaccaga 1620
agtcttccgg gtcaaccaat tacctgcaca tagtgatcac cattattatc acgacacagc 1680
agaattcaaa acgggtagtc gcttcatcaa gagtctcaat ggcgcttggc gttttaactt 1740
cgccaagaca ccggctgaac gccaggtga tttttatcaa ccgatttcg atgcaaccga 1800
ctttgatagc attcaagttc ccggtcatat tgaactagcc ggctatggc aaattcaata 1860
cattaacacg ttatacccat gggaaggtaa aatttaccgt cggccaccgt ataccctcaa 1920
tcaagatcaa ttaacaccag gctatttcag cgacgctgcg gacaacaccg tcggctcgta 1980
cctcaaaacc ttgatctcg acgatgtttt taaagggcaa cgtattatca ttcagttcca 2040
aggggtagaa gaagccctgt acgtctggtt aaatggccat tttattggct actctgaaga 2100
tagtttcacc ccttcagaat ttgatttgac gccgtatatt caggaccaag gtaacgtttt 2160
agcggttcgg gtctacaaac acagtactgc tgcctttatt gaagaccaag atatgttccg 2220
tttctctggt attttccgtg acgtcaatat actggcggag cctgctagcc atattactga 2280
tttgacatc cgaccagttc caaatgcaa tctcaaaagt ggtgagctca acatcactac 2340
taaagtaacc ggccaaccag ccactttagc gctgaccgtt aaagaccatg acgggcgagt 2400
actgacgagt caaacgcaaa ccggtagtg cagtgtaac tttgatacta tgttattcga 2460
ccaactgcac ttgtggtcac cacaacgcc gtatctctat caattgacaa ttgaagttha 2520
cgatgctgat caccaactct tggagtcgt cccatcatcg tttgggttcc ggacggtcga 2580
gctgcgcgat gacaaagtca tttacgtcaa caataaacgg ttgggtgatc acgggggtta 2640
ccggcacgaa tggaaagccc acacgggtcg cgttatcagt atggctgata tgcgcgctga 2700
tatccaaacc atgttagcta acaatatcaa tgccgatcgg acctgccatt atcctgatca 2760
attaccttgg tatcaattat gtgacgagg cggtatctac ctaatggccg aaaccaacct 2820
cgaatcgac ggggtcatggc aaaagatgg ggctatcgag ccttcttaca atgttccctg 2880
cgataatcca cactggccag cagcgggtgat cgaccgggcc cgttcaaact acgaatggtt 2940
taaaaaccac ccatcaatca ttttttggc acttggaat gaatcgtatg ctggcgaaga 3000
tatcgcgggc atgcaggctt tttataaaga acacgatgat tcacgactcg tccactacga 3060
agggcttttc tacacaccag aattaaaaga tcgcatthct gatgttgaaa gtcggatgta 3120
cgaaaagccc caaaatattg tagcttactt ggaagataac ccaaccaaac ctttcctaaa 3180
ttgtgaatat atgcatgaca tggggaattc tctggcggt atgcaatcat ataagattt 3240
gatcgacaag tatccaatgt atcaagggtg ctttatttgg gactttattg atcaagcact 3300
cttcgttcat gacccaatca ccgaccaaga cgtgctccgg tatggcggtg atttcgacga 3360
acgccactcc gattatgcat tctccggtta cggttaatg tttgccgacc ggacaccaa 3420
accagcaatg caagaggtga aatattatta tggcttacac aaataatcaa ctacacgtta 3480
tttacggcga cgggagctta ggactacagg gggctaattt ccactacctc tttagctacg 3540

```

```

aacgtggggg acttgaatca ctgctgctca acgataaaga gtggtcttat cgtacaccca 3600
cgcccatgtt ttggcgggcg acaaccgata atgatacagg tagcggcttt tcagtcaa 3660
ccgcacagtg gtacgcggcc gataagttct caacttgctca agatatacgaa ttgacggttg 3720
acgaccaacc agtcacaccg ttaccaatcg cgccactcaa taacaaatac acggatcacg 3780
aaatcgccac gaaagtctct ctggcttacc acttcgttac cacgaccgtt cctagtacca 3840
tcgtcacagt gacttatacg gtgacagcag acggtcagat caataatcgcc acccattata 3900
gcggtcagtc tgatttgcca gagctaccg catttggtct gcggtttatc atgccaacta 3960
ccgcgaccgg cttcgactat accggtttgt ccggtgagac ttatcctgac cggctggctg 4020
gcgcaacaca cgggcaattc cacgttgaca gtctgccagt cacaccatac ttggtccac 4080
aagaatgcgg catgcacatg caaactgaac aagtgcagat aacgcgatca acaacacaaa 4140
ataacgctga ccacgacaac acaccattca gtttaacatt tagccaaacc gatgcacat 4200
tcgccttcag ctgccttccc tataccgctg ctgaactaga aaacgcaaca cacatggaag 4260
aattaccatt agcacggcga acggtcttat caatctacgg tgccgttcgt ggggtcgggtg 4320
gcattgacag ttggggaacg gacgtagaag ccccatatca tatctcgtc aatcaagaca 4380
ttgacttcag ctttaacatt catttctaaa atttcaaaag aatcgctccg 4440
gcagttatct gccagagcgt tcttttaata tgttttatgg ctgagcttta gtcctttgaa 4500
gtgggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4560
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4620
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4680
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4740
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4800
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4860
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4920
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4980
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 5040
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 5100
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 5160
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 5220
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 5280
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 5340
gggggggggg gggggggggg gatgcccttg agagccttca acccagtcag ctcttcccg 5400
tggtgcgggg gcatgactat cgtcgcgcga cttatgactg tcttctttat catgcaactc 5460
gtaggacagg tgccgcgagc gctctgggtc atttccggcg aggaccgctt tcgctggagc 5520
gcgacgatga tcggcctgtc gcttgcggta ttcggaatct tgacgcctt cgctcaagcc 5580
ttcgtcactg gtcccgccac caaacgtttc ggcgagaagc aggccattat cgccggcatg 5640
gcggccgacg cgttgggcta cgtcttgctg gcgttcgcga cgcgaggctg gatggccttc 5700
ccattatga ttcttctcgc ttccggcggc atcgggatgc ccggttgca ggcatgctg 5760
tccaggcagg tagatgacga ccatcagga gagcttcaag gatcgcctgc ggctcttacc 5820
agcctaactt cgatcactgg accgctgac gtacgcgcga tttatgcgc ctgcgcgagc 5880
acatggaacg ggttggcag gattgtaggc gccgcctat acctgtctg cctcccccg 5940
ttgcgtcgcg gtgcatggag ccgggccacc tcgacctgaa tggaaagccg cggcacctcg 6000
ctaacggatt caccactcca agaattggag ccaatcaatt cttcgggaga actgtgaatg 6060
cgcaaaccaa cccttggcag aacatatcca tcgcgtccgc catctccagc agccgcacgc 6120
ggcgcatctc gggcagcgtt gggctctggc cacgggtgcg catgatcgtg ctctgtcgt 6180
tgaggaccgc gtaggctgg cggggttgcc ttactgggta gcagaatgaa tcaccgatac 6240
gcgagcgaac gtgaagcgac tgctgctgca aaacgtctgc gacctgagca acaacatgaa 6300
tggtcttcgg tttccgtgtt tcgtaaagtc tggaaacgcg gaagtccctt acgtgctgct 6360
gaagttgccc gcaacagaga gtggaaccaa ccggtgatac cacgatacta tgactgagag 6420
tcaacgccat gagcgcctc atttcttatt ctgagttaca acagtccgca ccgctgtccg 6480
gtagctcctt ccggtgggcg cggggcatga ctatcgtcgc cgcacttatg actgtcttct 6540
ttatcatgca actcgtaggc caggtgcccg cagcgcctaa cagtcccccg gccacggggc 6600
ctggccacct accacgcgg aaacaagcgc cctgcacat tatgttccg atctgcatcg 6660
caggtatgctg ctggctaccc tgtggaacac ctacatctgt attaacgaag cgctaaccgt 6720
ttttatcagg ctctgggagg cagaataaat gatcatatcg tcaattatta cctccacggg 6780
gagagcctga gcaaaactgg ctcaggcatt tgagaagcac acggtcacac tgcttccggt 6840
agtcaataaa ccggtaaacc agcaatagac ataagcggct atttaacgac cctgccctga 6900
accgacgacc gggctgaatt tgcttccgaa tttctgccat tcatccgctt attatcactt 6960
attcaggcgt agcaccaggc gtttaagggc accaataact gccttaaaaa aattacgccc 7020
cgccctgcca ctcatcgag tactgttgta attcatgaag cattctgccc acatggaagc 7080
catcacagac ggcgatgata acctgaatcg ccagcggcat cagcaccttg tcgccttgcg 7140
tataatattt gcccatggtg aaaacggggg cgaagaagtt gtccatattg gccacgttta 7200

```

```

aatcaaaact ggtgaaactc acccagggat tggctgagac gaaaaacata ttctcaataa 7260
accctttagg gaaataggcc aggttttcac cgtaacacgc cacatcttgc gaatatatgt 7320
gtagaaactg ccggaatcg tcgtggtatt cactccagag cgatgaaaac gtttcagttt 7380
gctcatggaa aacggtgtaa caagggtgaa cactatccca tatcaccagc tcaccgtctt 7440
tcattgccat acgaattcag aggtttgatg actttgacct ttggttttaa atttttgaaa 7500
aaaaataaaa aaggcgaagc ctattatata tttatcatat atattttaat cttttgttct 7560
tttgctgtaa aaaaaaggca gtgttttcgc tagttataga aattaaacag tcacaaaaat 7620
cgatgtatac agtcacaaaa atcgattttt gtgactctat gcacgatgt atacgagtca 7680
caaaaatcga tgtacacagt acgacttttg tatttgtgta ctgtatatag tataataaaa 7740
gcatagagaa aactcactat gaaatgactt tctctatgct actactaaaa cacgcaaagg 7800
agcgtattta tactatgatt ataccagaaa aacaaaacaa acaaaaagcag gtgctgacct 7860
tgaatgaatt agaaaaacgc aaagttagtg agcataacgc ttttaattcaa agtgtggcta 7920
aaatgcaaaa aacagcttta aaaatgtttg agttagccgt gtcttgtatt gataccgaag 7980
aacccecaaa aaataatact gtttatcttt taaaatctga acttttttaa ttttttgaag 8040
tatcgtctag cagtaagcat agtcaattca aagaagcagt taattatatg caaaagcaag 8100
cctttttcaa tattaagct gataagaac taggtattga gtatgaaagc attgtgccta 8160
ttccttatgt aaaatggaac gattataacg atgaagtaac aatacgtttt gaccaagcca 8220
ttatgcccta tctgattgat ttaaaggcag aatttactca atataaaatt tccgaactac 8280
aaaaactcaa tagcaagtat tcgattattt tgtatcgttg gttatccatg aattataacc 8340
aatacgaaca ttacagcgtg aaagggggac ggagagttga acaagtggaa tcctaccgca 8400
accctcaat aaaagtgaag gaattgcgag aaataactga tacaataaat gaacatcaac 8460
atthtccca ttttgaact agagtattaa aaaaagcaat tgaagaaatc aacgctcaca 8520
cctcttttaa tgtaacctat gagaaaaaga aagcagggcg gagcattgac tctattgtct 8580
ttcatattga gaagaaacgc atggcagacg ataacagtt caagttggga gataaagact 8640
atcaagacga taaaaagcaa aaatcaagaa atgaagctga cttactaaaa caggcaatgg 8700
aaagtaataa tacaagactt ttatctgaaa atthtctaat tggcatgaat gatattatgg 8760
atacagccac aatggttagc ttacagaaga atgtttatcc gctttatgac gaactgaaag 8820
aattaagagg gctaaatggt gtcaaagacc acttgtctta tgttgccagc aaaaaagaag 8880
cctattctaa acgtaatgta gcgaaatata ttaaaaaagc cattgaacaa tatctgcca 8940
cggttaaaag gcaggactta aaccatgagt gaagacttaa aaacgataaa agagtggcg 9000
gacgagctag gactagctaa acagaccgtt caataccatt taaagttttt gccgaccgaa 9060
aatcgacaaa aaaacaatag tggttcgatc gtcctaaacc ttgaagaaca aaagtttatc 9120
aaaaaaagc taacaaaaat aaaaaagtt caaccgaccg aaaatcgaca agaaaaaatg 9180
aagaattaaag cgttaattag cttttgttag gaattccgga tgagcattca tcaggcgggc 9240
aagaatgtga ataaaggccg gataaaactt gtgcttattt ttctttacgg tctttaaaa 9300
ggccgtaata tccagctgaa cggctctggt ataggtacat tgagcaactg actgaaatgc 9360
ctcaaaatgt tctttacgat gccattggga tatatcaacg gtggtatatc cagtgattht 9420
tttctccatt ttagcttccct tagctcctga aaatctcgat aactcaaaaa atacgcccg 9480
tagtgatctt atthcattat ggtgaaagtt ggaacctctt acgtgccgat caacgtctca 9540
ttttcgccaa agtttgccc agggcttccc ggtatcaaca gggacaccag gatttattha 9600
ttctgcgaag tgacttccg tcacaggtat ttattcggcg caaagtgcgt cgggtgatgc 9660
tgccaaacta ctgatttagt gtatgatggt gtttttgagg tgcctcagtg gcttctgtht 9720
ctatcagctg tccctcctgt tcagctactg acggggtggt gcgtaacggc aaaagcaccg 9780
ccggacatca gcgctagcgg agtgtatact ggcttactat gttggcactg atgagggtgt 9840
cagtgaagtg cttcatgtgg caggagaaaa aaggctgcac cgggtgcgta gcagaatatg 9900
tgatacagga tatattccgc ttctcgcgc actgactcgc tacgctcggg cgttcgactg 9960
cggcgagcgg aaatggctta cgaacggggc ggagatttcc tggagatgc caggaaagata 10020
cttaacaggg aagtgaagg gcccggcaa agcgttttt ccataggctc cggccctctg 10080
acaagcatca cgaatctga cgtcaaatc agtgggtggc aaaccgaca ggactataaa 10140
gataccaggc gtttccccct ggcggtctcc tcgtgcgctc tctgttctt gcctttcggt 10200
ttaccggtgt cattccgctg ttatggccgc gtttgtctca ttccacgctt gacactcagt 10260
tccgggtagc cagttcgcgc caagctggac tgtatgcacg aacccccgt tcagtcggac 10320
cgctgcgcct tatccggtaa ctatcgtctt gagtccaacc cggaaaagaca tgcaaaaagca 10380
ccactggcag cagccactgg taattgattt agaggagtta gtcttgaagt catgcgccg 10440
ttaaggctaa actgaaagga caagtttttg tgactgcgct cctccaagcc agttacctc 10500
gttcaaaagc ttggtagctc agagaacct cgaaaaacgc cctgcaagg cggttttttc 10560
gttttcagag caagagatta cgcgcagacc aaaacgatct caagaagatc atcttattha 10620
tcagataaaa tatttctaga ttctcagtga atthtctct tcaaatgtag cacctgaagt 10680
cagcccata cgatataagt tgtaattctc atgtttgaca gcttatcatc gatataaatt 10740
ccccgtagc gctagggacc tctttagctt cttggaagct gtcagtagta tatctaata 10800
tttatctcca ttcccttag taacgtgtaa ctttccaaat ttaaaaaagc gactcataga 10860

```

```

attatttcct cccgttaaatt aatagataac tattaataat agacaatact tgctcataag 10920
taatggtaact taaattgttt actttggcgt gtttcattgc ttgatgaaac tgatttttag 10980
taaacagttg acgatattct cgattgaccc attttgaaac aaagtacgta tatagcttcc 11040
aatattttatc tggaacatct gtggtatggc gggtaagttt tattaagaca ctgtttactt 11100
ttggttttagg atgaaagcat tccgctggca gcttaagcaa ttgctgaatc gagacttgag 11160
tgtgcaagag caaccctagt gtccggtgaa tatccaaggt acgcttgtag aatccttctt 11220
caacaatcag atagatgtca gacgcacggc tttcaaaaac cactttttta ataatttggtg 11280
tgcttaaattg gtaaggaata ttcccaacaa ttttatacct ctgtttggtta ggggaattgaa 11340
actgtagaat atcttggtga attaaagtga cacgagtatt cagttttaat ttttctgacg 11400
ataagttgaa tagatgactg tctaattcaa tagacgttac ctgtttactt attttagcca 11460
gtttcgctcgt taaatgccct ttacctgttc caatttcgta aacgggtatcg gtttctttta 11520
aattcaattg ttttattatt tgggtgagta ctttttactt cgtaaaaaag ttttgagaat 11580
attttatatt ttgttctatg taatctctcc tgaagtgtt acatctataa acaaatacag 11640
aagttaaacg atttgtttgt aatttttagtt atctgtttta aaagtcataa gattagtcac 11700
tggttaggaat taatctaacg tatttattta tctgcgtaat cactgttttt agtctgtttc 11760
aaaacagtag atgttttatc tacattacgc atttgaata ccaacatgac gaatccctcc 11820
ttcttaatta caaattttta gcatctaatt taacttcaat tcctattata caaaatttta 11880
agataatgca ctatcaacac actcttaagt ttgcttctaa gtcttatttc cataacttct 11940
tttaacgtttc cgccattctt tgctgtttcg atttttatga tatggtgcaa gtcagcacga 12000
acacgaaccg tcttatctcc cattatatct tttttgcac tgattggtgt atcatttctg 12060
tttctttttt gtgcgcctaa atttcccaca atcactcact tctttctatt tcttcttatt 12120
cttattttat catcaacaat cacaatatc ttgtgatttg tgataagtga tttgtgataa 12180
gtgatttggtg attaatatat aaaagccctc tttaaagggc ttttatgttt attttgagaa 12240
agataataaaa tcaatatatc ccttttccc aatttttaca acggcattgt agggcttttt 12300
ctctttgttt ttgatttctt tttaccaggg tttcttttcc ctccagtaat tcttttacat 12360
tcgttttggt tagttttttc tttctaaaat gttcagctaa agtaaaactta cattcaggat 12420
aatttgaaca accataaaac gattttttta atacaatatt gttgccacac ttaggacatt 12480
ttcttacaat acttttttct gcttcttttt ctttctgttc ctggtaatca gaaaaattta 12540
gtttttctat atcgtttaggt acagcttcca gtaaatgaac aatgaatttt ttgatattcg 12600
taataaagtt ctcttgattg ccttctcttt taccgatttt ttttaataac gtttccatt 12660
tagccgtcat ttcagcactc gttaaaaggt gctgactttc aactgcctgg cacaataatt 12720
ttcctttttc agttacaac 12739

```

<210> 9
 <211> 57
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide P59

<400> 9
 gaccgacatt aaattcttga cagggagaga taggtttgat agaataataat agttgtc 57

<210> 10
 <211> 58
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide P59 complement

<400> 10
 tcgagacaac tatttatattc tatcaaacct atctctccct gtcaagaatt taatgtcg 58

<210> 11
 <211> 17
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Lac operator sequence

<400> 11

ttgttatccg ctcacaa

17

<210> 12

<211> 94

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide Xyl/Lac-T5

<400> 12

gaccgtcata aaaaatttat ttgctttgtg agcggataac aattataata gattcaagtt 60
agtttggtta ttaaattaac caactaaaat gtag 94

<210> 13

<211> 95

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide Xyl/Lac-T5 complement

<400> 13

aattctacat tttagttggt taatttaata aacaaactaa cttgaatcta ttataattgt 60
tatccgctca caaagcaaat aaatttttta tgacg 95

<210> 14

<211> 57

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide Xyl/Lac-P59

<400> 14

gaccgacatt aaattcttga cattgtgagc ggataacaat agaataataat agttgtc 57

<210> 15

<211> 58

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide Xyl/Lac P59 complement

<400> 15

tcgagacaac tattatattc tattgttatc cgctcacaat gtcaagaatt taatgtcg 58

<210> 16

<211> 6052

<212> DNA

<213> Artificial Sequence

<220>

<223> Vector pXyl-T5 p15a

<400> 16

```

tggcattcta cgactataac ttaaatttat attttttact ttataatata taattgatta 60
tagaataatg ttgctcatat cgtttgccaa catctagtag tcaaattaca ctatgttaca 120
cttggttaata ttaaccgaac ttcccctgtc caaattagat aagaggtaat aataaatgga 180
aaataatttt atagtaaag aaaatgagaa gcgtgtatta aaacaaattt tcaataacag 240
caatatttca cgaacacaaa tatcgaagaa tttagaactt aataaagcta ctattttctaa 300
cattctgaac aacttaaaac acaagagttt agttaatgaa gtaggagaag gtaatagtag 360
taaaagtggg ggacgaaagc ctattttact cgaaattaac caaaaatatg gctactatat 420
ttctatggat ttaacatatg attccgttga attaagtac aactactttg atgtactat 480
attaagcaa gattcctacg aattaaatga taaaaatgta agcagtatat taaaaatttt 540
aaaatctaataaaacgtct cagaaaaata tgatacgtta tatgggttac ttggtatata 600
tatatccata cacggtatcg ttgacgatga gcaaacata atcaatcttc cttttcataa 660
aatgagaaa cgacatttta ccgatgaatt aaagtcattc acaaatgttc ctgtcgttat 720
agaaaatgaa gcaaatttat cagcgctata tgaaaaagt ttatatatta attcaaact 780
aaataatttg attactttta gtattcaca gggtatagg gctggcatca taataaataa 840
aaaactttat ctgggctcaa atggagaggc tggagagata ggtaagacat tggttttgga 900
atctataaat acaaatgaca acaaatatta taaaatcgaa gatatatgct cccaagacgc 960
tttaatacag aaaaaaata ataggttggg cgtcacattg acgtttacag aactaatcca 1020
atattacaac gaaggaaatt caattgttgc tcatgaaatt aaacaattta ttaataaaat 1080
gacagtcttg attcataatt tgaatacaca atttaacca gacgctattt atattaactg 1140
tcctttaatt aatgaattac caaatatttt aaatgaaatt aaagagcaat tctcctgttt 1200
ttctcaaggc agtccaattc aattacattt aactactaat gtaaaacaag ctactttatt 1260
gggtggcact ttgcaataaa tgcaaaaaac attaaatata aataacattc aaatgaatat 1320
taataaatta cagcagctcg agttataaaa tagatatctc ggaccgcat aaaaaattta 1380
tttgctttca ggaaaatttt tctgtataat agattcaagt tagtttgttt attaaattaa 1440
ccaactaaaa ttagaattc gagctcggta cccggggatc ctctagagtc gacctgcagc 1500
caagcttggg cttttcagcc tgatacagat taaatcagaa cgcagaagcg gtctgataaa 1560
acagaatttg cctggcggca gtagcgcggg ggtcccacct gaccccatgc cgaactcaga 1620
agtgaacgc cgtagcgcgg atggtagtgt ggggtctccc catgagagag tagggaactg 1680
ccaggcatca aataaaactg aaggctcagt cgaagactg ggcctttcgt ttatctgtt 1740
gtttgtcgtt gaacgctctc ctgagtagga caaatccgcc gggagcggat ttgaacgttg 1800
cgaagcaacg gcccgagggg tggcgggcag gacgcccgcc ataaactgcc aggcatacaa 1860
ttaagcagaa ggccatcctg acggatggcc tttttgcgtt tctacaaact ctttttgttt 1920
atttttctaa atacattcaa atatgtatcc gctgatcccc atctatcga tgataagctg 1980
tcaaacatga gaattaaatc aatctaaagt atatatgagt aaacttggtc tgacagttac 2040
caatgcttaa tcagtggagg acctatctca gcgatctgtc tatttcgttc atccatagtt 2100
gcctgactcc cgtcgtgta gataactacg atacgggagg gcttaccatc tggccccagt 2160
gctgcaatga taacgcgaga cccacgctca ccggctccag atttatcagc aataaaccag 2220
ccagccggaa gggccgagcg cagaagtggg cctgcaactt tatccgcctc catccagttc 2280
attaattgtt gccgggaagc tagagtaagt agttcgccag ttaatagttt gcgcaacgtt 2340
gttgccattg ctacaggcat cgtggtgtca cgctcgtcgt ttggtatggc ttcattcagc 2400
tccggttccc aacgatcaag gcgagttaca tgatcccca tgttggtcaa aaaagcgggt 2460
agctccttcg gtccctcgat cgtgtcaga agtaagttgg ccgagtggt atcactcatg 2520
gttatggcag cactgcataa ttctcttact gtcatgccat ccgtaagatg cttttctgtg 2580
actggtgagt actcaacca gtcatctga gaatagtga tgcggcgacc gagttgctct 2640
tgccggcgct caacacggga taataccgcg ccacatagca gaactttaaa agtgctcatc 2700
attgaaaaac gctcttcggg gcgaaaactc tcaaggatct taccgctgtt gagatccagt 2760
tcgatgtaac ccactcgtgc acccaactga tcttcagcat cttttacttt caccagcgtt 2820
tctgggtgag caaaaaacag aaggcaaaat gccgcaaaa agggataaag ggcgacacgg 2880
aaatgttgaa tactcatact cttccttttt caatattatt gaagcattta tcagggttat 2940
tgtctcatga gcgatacat atttgaatgt atttagaaaa ataaacaaat aggggttccg 3000
cgcacatttc cccgaaaagt gccacctggc ggccgcataa gatgatcttc ttgagatcgt 3060
tttggctctgc gcgtaatctc ttgctctgaa aacgaaaaaa ccgccttgca gggcggtttt 3120
tcgaagggttc tctgagctac caactctttg aaccgaggta actggcttgg aggagcgag 3180
tcacaaaaac ttgtcctttc agtttagcct taaccggcgc atgacttcaa gactaactcc 3240
tctaaatcaa ttaccagtgg ctgctgccag tgggtctttt gcatgtcttt ccgggttggg 3300
ctcaagacga tagttaccgg ataaggcgca gcggtcggac tgaacggggg gttcgtgcat 3360
acagtccagc ttggagcgaa ctgcctacc ggaactgagt gtcaggcgtg gaatgagaca 3420
aacgcggcca taacagcgga atgacaccgg taaaccgaaa ggcaggaaca ggagagcgca 3480
cgagggagcc gccaggggaa acgcctggta tctttatagt cctgtcgggt ttcgccacca 3540
ctgatttgag cgtcagattt cgtgatgctt gtcagggggg cggagcctat ggaaaaacgg 3600

```

```

ctttgccgcg gccctctcac ttccctgtta agtatcttcc tggcatcttc caggaaatct 3660
ccgccccgtt cgtaagccat ttccgctcgc cgcagtcgaa cgaccgagcg tagcgagtca 3720
gtgagcgagc aagcgggaata tatcctgtat cacatatctt gctgacgcac cggcgagcc 3780
ttttttctcc tgccacatga agcacttcac tgacaccctc atcagtgcca acatagtaag 3840
ccagtgcggc cgcttctttc ctgcgttata cctgattctt gtggataacc gtattaccgc 3900
ctttgagtga gctgataccg ctgcgcgag ccgaacgacc gagcgagcg agtcagtga 3960
cgaggaagcg gaagagcgcc tgatgcggtt ttttctcctt acgcatctgt gcggtatttc 4020
acaccgcata agatcccctc gacctgcagg catgcaagct tctgtagggt tttaggcata 4080
aaactatatg atttaccctt aaatctttaa aatgccctt aaaattcaaa ataaaggcat 4140
ttaaaattta aatatttctt gtgataaagt ttgttaaaaa ggagtgggtt tatgactgtt 4200
atgtgggtat cgattatagg tatgtgggtt tgtattggaa tggcattttt tgctatcaag 4260
gttattaaaa ataaaaatta gaccacgat ttatgccgag aaaatttatt gtgcgttgag 4320
aagaaccctt aactaaactt gcagacgaat gtcggcatag cgtgagctat taagccgacc 4380
attcgacaag ttttgggatt gttaagggtt ccgaggctca acgtcaataa agcaattgga 4440
ataaagaagc gaaaaaggag aagtcggtt agaaaaagaa ggatatggat ctggagctgt 4500
aatataaaaa ctttcttcaa ctaacggggc aggttagtga cattagaaaa ccgactgtaa 4560
aaagtacagt cggcattatc tcatattata aaagccagtc attaggccta tctgacaatt 4620
cctgaataga gttcataaac aatcctgcat gataaccatc acaaacagaa tgatgtacct 4680
gtaaagatag cggtaaatat attgaattac ctttattaat gaattttcct gctgtaataa 4740
tggttagaag gtaattacta ttattattga tatttaagtt aaaccagta aatgaagtc 4800
atggaataat agaaagagaa aaagcatttt cagggtatagg tgttttgga aacaatttc 4860
ccgaaccatt atatttctct acatcagaaa ggtataaatc ataaaactct ttgaagtc 4920
tctttacagg agtccaaata ccagagaatg ttttagatac accatcaaaa attgtataaa 4980
gtggctctaa cttatcccaa taacctaaact ctccgtcgct attgtaacca gttctaaaag 5040
ctgtatttga gtttatcacc cttgtcacta agaaaaataa tgcagggtaa aatttatatc 5100
cttcttggtt tatgtttcgg tataaaacac taatatcaat ttctgtggtt atactaaaag 5160
tcgtttgttg gttcaaataa tgattaaata tctcttttct cttccaattg tctaaatcaa 5220
ttttattaaa gttcatttga tatgctcctt aaatttttat ctaaagtga tttaggaggc 5280
ttacttgtct gctttcttca ttagaatcaa tcttttttta aaagtcaata ttactgtaac 5340
ataaatatat attttaaaaa tatcccactt tatccaattt tcgtttgttg aactaatggg 5400
tgcttttagt gaagaataaa gaccacatta aaaaatgtgg tcttttgtgt ttttttaaag 5460
gatttgagcg tagcgaaaaa tcttttctt tcttatcttg ataataaggg taactattgc 5520
cggcgaggct agttaccctt aagttattgg tatgactggt tttaagcgca aaaaaagttg 5580
cttttctgta cctattaatg tatcgtttta aatgactagt aaaaaacata catagaaaag 5640
ggaaaaagca acttttttta ttgtcatagt ttgtgaaaac taagttgttt ttatgtgtta 5700
taacatggaa aggtatactg agaaaaaaca agaaatcaa gtatttcaga aatttattaa 5760
acgtcatatt ggagagaatc aaatggattt agttgaagat tgcaatacat ttctgtcttt 5820
tgtagctgat aaaactttag aaaaacagaa attatataaa gctaattctt gtaaaaatcg 5880
attttgtcct gtctgtgctt ggagaaaagc tagaaaagat gcattgggtt tatctttgat 5940
tgatgaatat attagcagc agagaaaaa ggagtttata tttttaactt tgactacacc 6000
taatgtaatg agtgatgaat tagaaaatga aataaaacgt tataataatt cttttagaaa 6060
acttataaag agaaaaaaag taggtagtgt tataaaggga tatgttcgta agttagagat 6120
tacatataat aaaaaaagag atgattataa tctcatctt catgtgttaa ttgcagtaaa 6180
taaactcgat ttcacagata aaagatatta tattagccaa caagaatggt tagatttatg 6240
gcgtgatgta acgggcattt cagaaataac acaagttcaa gttcaaaaaa taagacaaaa 6300
taataataaa gaattatatg aaatggctaa gtattctggt aaagatagtg attatttaat 6360
aaataaatca aaaagtcttt gatgcatttt ataaatcact taaaggtaaa cagggtattag 6420
tttattcagg attattttaa gaggctaaaa agaaattaaa aaatggggat ttagattact 6480
taaaagaaat tgatccaacc gtaatatatc tatcaaattt tttatatttg gaaacaaaaa 6540
gagtatttag ctagtgaact ttatgactta acagaacaag aaaaaagaga aattaatcac 6600
aaaatgatag acgaaatcga ggaagaacaa taacaaaata taagtgttaa cagtcgtctg 6660
caagtttagt taagggttct tctcaacgca caataaattt tctcggcata aatgcgtggt 6720
ctaattttta tttttaataa ctttgatagc aaaaaatgcc attccaatac aaaaccacat 6780
acctataatc gataaccaca taacagtcac aaaaccactc ctttttaaca aactttatca 6840
caagaatatat tt 6852

```

<210> 17
<211> 9437
<212> DNA

<213> Artificial Sequence

<220>

<223> Vector pEPEF3

<400> 17

```

caacccagtc agctccttcc ggtgggcgcg gggcatgact atcgtcgccg cacttatgac 60
tgtcttcttt atcatgcaac tcgtaggaca ggtgccggca gcgctctggg tcattttcgg 120
cgaggaccgc ttctcgctgga gcgcgacgat gatcgccctg tcgcttgccg tattcggaat 180
cttgacgccc ctgcctcaag ccttcgtcac tgggtccgcc accaaacgtt tcggcgagaa 240
gcaggccatt atcgccggca tggcgccga cgcgctgggc tacgtcttgc tggcgctcgc 300
gacgcgaggg tggatggcct tccccattat gattcttctc gcttccggcg gcatcgggat 360
gcccgcgttg caggccatgc tgtccaggca ggtagatgac gaccatcagg gacagcttca 420
aggatcgctc gcggtcttta ccagcctaac ttcgatcact ggaccgctga tcgtcacggc 480
gatttatggc gcctcggcga gcacatggaa cgggttgga tggattgtag gcgccgccct 540
ataccttgtc tgactccccg cgttgcgctc cggtgcattg agccggggcca cctcgacctg 600
aatggaagcc ggcggcacct cgctaaccga ttaccactc caagaattgg agccaatcaa 660
ttcttgccga gaactgtgaa tgcgcaaac aacccttggc agaacatata catcgctcc 720
gccatctcca gaagcgcac gcggcgcatc tcgggcagcg ttgggtcctg gccacgggtg 780
cgcatgatcg tgctcctgtc gttgaggacc cggctaggct ggccggggtt ccttactggt 840
tagcagaatg aatcacccgat acgcgagcga acgtgaagcg actgctgctg caaaacgtct 900
gcgacgttag caacaacatg aatggtcttc ggtttccgtg ttctgtaaa tctggaaacg 960
cggaagtccc ctacgtgctg ctgaagtgtc cggtcaacga gagtggaaac aaccgggtgat 1020
accacgatac tatgactgag agtcaacgcc atgagcggcc tcattttctta ttctgagtta 1080
caacagtcct caccgctgtc cggtagctcc ttccggtggg cgccggggcat gactatcgtc 1140
gccgcactta tgactgtctt ctttatcatg caactcgtag gacaggtgcc ggcagcggcc 1200
aacagtcccc cgccacaggg gcctgccacc ataccacgc cgaacaagc gccctgcacc 1260
attatgttcc gcatctgcat cgaggatgc tgctggctac cctgtggaac acctacatct 1320
gtattaacga agcgctaacc gtttttatca ggctctggga ggagaataa atgatcatat 1380
cgtaacttat tactcctcag gggagagcgt ggcgaactg gcctcaggca ttgagaagc 1440
acacggtcac actgcttccg gtagtcaata aaccggtaaa ccagcaatag acataagcgg 1500
ctatttaacg accctgccct gaaccgacga ccgggtcgaa ttgtcttctg aatttctgcc 1560
attcatccgc ttattatcac ttattcaggc gtagcaccag gcgtttaagg gcaccaataa 1620
ctgccttaaa aaaattacgc cccgccctgc cactcatcgc agtactgttg taattcatta 1680
agcattctgc cgacatggaa gccatcacag acggcatgat gaacctgaat cgccagcggc 1740
atcagcacct tgcgccttg cgtataatat ttgccatgg tgaacacggg gcggaagaag 1800
ttgtccatat tggccacgtt taaatcaaaa ttggtgaaac tcaccacagg attggctgag 1860
acgaaaaaca tattctcaat aaacccttta gggaaatagg ccaggttttc accgtaacac 1920
gccacatctt gcgaatatat gtgtagaaac tgccggaaat cgtcgtggta ttactccag 1980
agcgatgaaa acgtttcagt ttgctcatgg aaaaagggtg aacaagggtg aacactatcc 2040
catatcacca gtcaccgtc ttctattgcc atacgaattc agaggtttga tgactttgac 2100
ctttggtttt aaatttttga aaaaaataaa aaaaaggcga gcctattata tatttatcat 2160
atataattta atcttttgtt cttttgcgtg aaaaaaaagg cagtgttttc gtagttata 2220
gaaattaaac agtcacaaaa atcgatgat acagtcacaa aaatcgattt ttgtgactct 2280
atgcacgat gtatacgagt cacaaaaatc gatgtacaca gtacgacttt tgtatttgtg 2340
tactgtatat agtataataa aagcatagag aaaactcact atgaaatgac tttctctatg 2400
ctactactaa aacacgcaaa ggagcgtatt tatactatga ttataccaga aaaacaaaaa 2460
aaacaaaagc aggtgctgac cttgaatgaa ttagaaaaac gcaagtagt ggagcataac 2520
gctttaattc aaagtgtggc taaaatgcaa aaacagctt taaaatgtt tgagttagcc 2580
gtgtcttgta ttgataccga agaaccoccc aaaaataata ctgtttatct tttaaaatct 2640
gaacttttta aattttttga agtatcgtct agcagtaagc atagtcaatt caaagaagca 2700
gttaattata tgcaaaagca agcctttttc aatattaaag ctgataagaa actaggtatt 2760
gagtatgaaa gcattgtgcc tattccttat gtaaaatgga acgattataa cgatgaagta 2820
acaatacgtt ttgaccaagc cattatgccc tatctgattg atttaaaggc agaatttact 2880
caatataaaa ttccggaact acaaaaactc aatagcaagt attcgattat tttgtatcgt 2940
tggttatcca tgaattataa ccaatacgaa cattacagcg tgaaaggggg acggagagtt 3000
gaacaagtgg aatcctaccg caacccctca ataaaagtga aagaattgcg agaaataact 3060
gatacaataa atgaacatca acattttccc cattttgaaa ctagagtatt aaaaaagca 3120
attgaagaaa tcaacgctca cacctctttt aatgtaacct atgagaaaaa gaaagcaggg 3180
cggagcattg actatttgt cttcatatt gagagaagac gcatggcaga cgataacagt 3240
tacaagttgg gagataaaga ctatcaagac gataaaaagc aaaaatcaag aaatgaagct 3300

```

```

gacttactaa aacaggcaat ggaaagtaaa tatacaagac ttttatctga aaatttctta 3360
attggcatga atgatattat ggatacagcc acaatggtag gcttacagaa gaatgtttat 3420
ccgctttatg acgaactgaa agaattaaga gggctaaatg gtgtcaaga ccacttgtct 3480
tatgttgcca gcaaaaaaga agcctattct aaacgtaatg tagcgaata tcttaaaaaa 3540
gccattgaac aatatctgcc aacggttaaa aggcaggact taaaccatga gtgaagactt 3600
aaaaacgata aaagagttgg cggacgagct aggactagct aaacagaccg ttcaatacca 3660
tttaaagttt ttgccgaccg aaaatcgaca aaaaaacaat agtggttcga tcgtcctaaa 3720
ccttgaagaa caaaagttta tcaaaaaaag actaacaaaa ataaaaaag ttcaaccgac 3780
cgaaaaatcg caagaaaaaa tgaagaatta agcgtaata gtcttttggt aggaattccg 3840
gatgagcatt catcaggcgg gcaagaatgt gaataaaggc cggataaaac ttgtgcttat 3900
ttttctttac ggtctttaa aaggccgtaa tatccagctg aacggtctgg ttataggtac 3960
attgagcaac ttactgaaat gcctcaaaat gttctttacg atgccattgg gatatatcaa 4020
cgggtgtata tccagtgtat tttttctcca ttttagcttc cttagctcct gaaaatctcg 4080
ataactcaaa aaatacgcgc ggtagtgate ttatttcatt atggtgaaag ttggaacctc 4140
ttacgtgccc atcaacgtct cattttcgcc aaaagtggc ccagggttc ccggtatcaa 4200
cagggacacc aggatttatt tattctcgca agtgatcttc cgtcacagggt atttattcgg 4260
cgcaaaagtg gtcgggtgat gctgccaaact tactgattta gtgtatgatg gtgtttttga 4320
ggtgctccag tggcttctgt ttctatcagc tgtccctcct gttcagctac tgacgggggtg 4380
gtgcgtaacg gcaaaagcac cgccggacat cagcgctagc ggagtgtata ctggcttact 4440
atgttggcac tgatgaggtg gtcagtgaag tgcttcatgt ggcaggagaa aaaaggctgc 4500
accggtgcgt cagcagaata tgtgatacag gatataatcc gcttccctgc tcaactgactc 4560
gctacgctcg gtcgttcgac tgcggcgagc ggaaatggct tacgaacggg gcggagattt 4620
cctggaagat gccaggaaga tacttaacag ggaagtgaag gggccgcggc aaagccgttt 4680
ttccataggc tccgcccccc tgacaagcat cacgaaatct gacgctcaaa tcagtgtgtg 4740
cgaaaccoga caggactata aagataccag gcgtttcccc ctggcggtc cctcgtgcgc 4800
tctctgttcc ctgcctttcg gtttaccggg gtcattccgc tgttatggcc gcgtttgtct 4860
cattccacgc ctgacactca gttccgggta ggcagttcgc tccaagctgg actgtatgca 4920
cgaaccccc gttcagtcgc accgctgcgc cttatccggg aactatcgtc ttgagtccaa 4980
cccggaaaga catgcaaaag caccactggc agcagccact ggtaattgat ttagaggagt 5040
tagtcttgaa gtcatgcgcc ggtaaggct aaactgaaag gacaagtgtt ggtgactgcg 5100
ctcctccaag ccagttacct cggttcaaa agtttgtagc tcagagaacc ttcgaaaaac 5160
cgccctgcaa ggcggttttt tcgttttcag agcaagagat tacgcgcaga ccaaaacgat 5220
ctcaagaaga tcatcttatt aatcagataa aatatttcta gatttcagtg caatttatct 5280
cttcaaatgt ctgacactga gtcagccca tacgatataa gttgtaattc tcatgtttga 5340
cagcttatca tcgatacaaa tccccgtag gcgctaggga cctctttagc ttcttggaag 5400
ctgtcagtag tatacttaat aatttatctc cattcccttt agtaacgtgt aactttccaa 5460
atttaaaaaa gcgactcata gaattatttc ctcccgttaa ataatagata actattaaaa 5520
atagacaata cttgctcata agtaatggta cttaaattgt ttactttggc gtgtttcatt 5580
gcttgatgaa actgattttt agtaaacagt tgacgatatt ctcgattgac ccattttgaa 5640
acaaagtacg tatatagctt ccaatattta tctggaacat ctgtggtatg gcgggtaagt 5700
tttatgaaga cactgtttac ttttggttta gtagaaagc attccgctgg cagcttaagc 5760
aattgtcgaa tcgagacttg agtgagcaag agcaacccta gtgttcggtg aatatccaag 5820
gtacgcttgt agaatccttc ttcaacaatc agatagatgt cagacgcacg gctttcaaaa 5880
accacttttt taataatttg tgtgcttaaa tggtaaggaa tattcccaac aattttatac 5940
ctctgtttgt tagggaattg aaactgtaga atatcttggg gaattaaagt gacacgagta 6000
ttcagtttta atttttctga cgataagttg aatagatgac tgtctaattc aatagacgtt 6060
acctgtttac ttatttttagc cagtttcgtc gttaaatgcc ctttacctgt tccaatttcg 6120
taaaccgtat cggtttcttt taaattcaat tgttttatta tttggttgag tactttttca 6180
ctcgttaaaa agttttgaga atattttata tttttgttca tgtaattctc cctgaagtga 6240
ttacatctat aaacaaatac agaagttaaa cgatttggtt gtaatttttag ttatctgttt 6300
aaaaagtcac aagatttagc actggttaga attaatctaa cgtattttatt tatctgcgta 6360
atcactgttt ttagtctgtt tcaaaacagt agatgtttta tctacattac gcatttgga 6420
taccaacatg acgaatccct ccttcttaat taaaaattt tagcatctaa tttaacttca 6480
attcctatta taaaaattt taagataatg cactatcaac acactcttaa gtttgcttct 6540
aagtcttatt tccataactt cttttacgtt tccgccattc tttgctgttt cgatttttat 6600
gatattggtc aagtgcagac cgtcttatct cccattatat ctttttttgc 6660
actgattggg gtatcatctt gttttctttt ttgtgcgcct aaatttccca caatcactca 6720
cttctttcta tttcttctta ttcttatttt atcatcaaca atcacaaatc acttggtgatt 6780
tgtgataagt gatttgtgat aagtgatttg tgattaatat ataaaagccc tctttaagg 6840
gcttttatgt ttattttgag aaagatataa aatcaatata tcccttttcc ccaattttta 6900
caacggcatt gtagggcttt ttctctttgt ttttgatttc cttttaccag ggtttctttt 6960

```

```

ccctccagta attcttttac attcgttttg gttagttttt tctttctaaa atgttcagct 7020
aaagtaaact tacatttcagg ataatttgaa caaccataaa acgatttttt taatacaata 7080
ttgttgccac acttaggaca ttttcctaca atactttttt ctgcttcttt ttctttctgt 7140
tcctggtaat cagaaaaatt tagtttttct atactgtag gtacagcttc cagtaaatga 7200
acaatgaatt ttttgatatt cgtaataaag ttctcttgat tgccttctct tttaccgatt 7260
ttttttaaat acgtttccca tttagccgtc atttcagcac tcgttaaaag gtgctgactt 7320
tcaactgcct ggcacaataa ttttcctttt tcagttacaa caagcttgcc tgcaggtcga 7380
cgataaccac ataacagtca taaaaccact cctttttaac aaactttatc acaagaaata 7440
ttttggcatt ctacgactat aacttaaat tatatttttt actttataat atataattga 7500
ttatagaata atgttgctca tatcgtttgc caacatctag tactcaaatt acactatgtt 7560
acacttggtg atattaaccg aacttcccct gtccaaatta gataagaggt aataataaat 7620
ggaaaaaat tttatagtaa atgaaaatga gaagcgtgta ttaaaacaaa ttttcaataa 7680
cagcaatatt tcacgaacac aaatatcgaa gaatttagaa cttaataaag ctactatttc 7740
taacattctg aacaacttaa aacacaagag tttagttaat gaagtaggag aaggtaatag 7800
tactaaaagt ggtggcgaag agcctatttt actcgaaatt aacaaaaaat atggctacta 7860
tatttctatg gatttaacat atgattccgt tgaattaatg tacaactact ttgatgtac 7920
tatattaaag caagattcct acgaattaaa tgataaaaaat gtaagcagta tattacaaat 7980
tttaaaatct aatataaacg tctcagaaaa atatgatacg ttatatgggt tacttggtat 8040
atctatatcc atacacggtg tcgttgacga tgagcaaaac ataatcaatc ttcttttca 8100
taaaaatgag aaacgcacat ttaccgatga attaaagtca ttcacaaatg ttctgtcgt 8160
tatagaaaaat gaagcaaatt tatcagcgct atatgaaaaa agtttatata ttaattcaaa 8220
cataaataat ttgattactt taagtattca caaggtata ggcgctggca tcataataaa 8280
taaaaaactt tatctggct caaatggaga ggctggagag ataggttaaga cattggtttt 8340
ggaatctata aataacaatg acaacaaata ttataaaatc gaagatatat gctcccaaga 8400
cgctttaata cagaaaataa ataataggtt gggcgtcaca ttgacgttta cagaactaat 8460
ccaatattac aacgaaggaa attcaattgt tgctcatgaa attaaacaat ttattaataa 8520
aatgacagtt ctgattcata atttgaatac acaatttaac ccagacgcta tttatattaa 8580
ctgtccttta attaatgaat taccaaatat tttaaatgaa attaaagagc aattctcctg 8640
tttttctcaa ggcagtccaa ttcaattaca tttactact aatgtaaaac aagctacttt 8700
attgggtggc acttttagcaa taatgcaaaa aacatttaat ataaataaca ttcaaataa 8760
tattaaataa ttacagcagt ctgagttata aaatagatat ctcggaccgc tttggcagtt 8820
tattcttgac atgttagtga gggggctggt ataatcacat agtactgttc tcgagagtta 8880
gtttgtttat taaattaacc aactggatcc gaattcccgg ggcacgctg acctgcagcc 8940
aagcttgggc ttttcagcct gatacagatt aaatcagaac gcagaagcgg tctgataaaa 9000
cagaatttgc ctggcggcag tagcgcggtg gtcccacctg acccatgcc gaactcagaa 9060
gtgaaacgcc gtagcgccga tggtagtgtg ggggtctccc atgcbagagt agggaaactgc 9120
caggcatcaa ataaaacgaa aggtcagtc gaaagactgg gcctttcgtt ttatctgttg 9180
ttgtcgggtg aacgctctcc tgagttagac aaatccgccc ggagcgatt tgaacgttgc 9240
gaagcaacgg cccggagggt ggcgggcagg acgcccgcga taaactgccg ggcatacaat 9300
taagcagaag gccatcctga cggatggcct ttttgcggtt ctacaaactc tttttgttta 9360
tttttctaaa tacattcaaa tatgtatccg ctgatcccca tctatcgga tctcgaccg 9420
atgcccttga ggcctt 9437

```

<210> 18

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide PL10

<400> 18

gaccgtttcg tgaacttttt tgttgacaaa gataaaaaaca catgatataa ttaaatcac 59

<210> 19

<211> 60

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide PL10 complement

<400> 19

tcgagtgatt taattatattc atgtgttttt atctttgtca acaaaaaagt tcacgaaacg 60

<210> 20

<211> 64

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide P59*

<400> 20

gaccgaaaaa tgacagttta ttcttgacag ggagagatag gtttgatata atataatagt 60
tgtc 64

<210> 21

<211> 65

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide P59* complement

<400> 21

tcgagacaac tattatatta tatcaaact atctctccct gtcaagaata aactgtcatt 60
tttcg 65

<210> 22

<211> 94

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide Xyl-T5-DD

<400> 22

gaccgtcata aaaaatttat ttacatcag gaaaattttt ctgtatatta gattcaagtt 60
agtttgttta ttaaattaac caactaaat gtag 94

<210> 23

<211> 95

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide Xyl-T5-DD complement

<400> 23

aattctacat tttagttggt taatttaata aacaaactaa cttgaatcta atatacagaa 60
aaattttcct gatgtaaaat aaatttttta tgacg 95

<210> 24

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> pXyl-T5F primer

<400> 24

cagcagtctg agttataaaa tag

23

<210> 25

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> LexL primer

<400> 25

tgttttatca gaccgctt

18

<210> 26

<211> 80

<212> DNA

<213> Artificial Sequence

<220>

<223> Xyl-T5 fusion sequence

<400> 26

tcataaaaaa tttatttgct ttcaggaaaa tttttctgta taatagattc aagttagttt 60

gtttattaaa ttaaccaact

80

<210> 27

<211> 80

<212> DNA

<213> Artificial Sequence

<220>

<223> Xyl-T5-DD fusion sequence

<400> 27

tcataaaaaa tttatttttac atcaggaaaa tttttctgta tattagattc aagttagttt 60

gtttattaaa ttaaccaact

80

<210> 28

<211> 95

<212> DNA

<213> Artificial Sequence

<220>

<223> CP25 fusion sequence

<400> 28

ctttggcagt ttattcttga catgttagtg agggggctgg tataatcaca tagtactgtt 60

ctcgagagtt agtttggtta ttaaattaac caact

95

<210> 29

<211> 86

<212> DNA

<213> Artificial Sequence

<220>

<223> P32 fusion sequence

<400> 29

gtgagcttgg actagaaaa aacttcacaa aatgctatac taggtaggta actcgagagt 60
tagtttgttt attaaattaa ccaact 86

<210> 30
<211> 86
<212> DNA
<213> Artificial Sequence

<220>
<223> P59 fusion sequence

<400> 30
acattaaatt cttgacaggg agagataggt ttgatagaat ataatagttg tctcgagagt 60
tagtttgttt attaaattaa ccaact 86

<210> 31
<211> 93
<212> DNA
<213> Artificial Sequence

<220>
<223> P59* fusion sequence

<400> 31
aaaaatgaca gtttattctt gacagggaga gataggtttg atataatata atagttgtct 60
cgagagtttag tttgtttatt aaattaacca act 93

<210> 32
<211> 150
<212> DNA
<213> Artificial Sequence

<220>
<223> P1P2 fusion sequence

<400> 32
gtctagttat tcttgacaaa aaataatatt ttgatataat taaatagttg tcgtttgata 60
acacagttta ttcttgacaa aaaaatataa aagtgtataa tagaaaagta ctgttctcga 120
gagttagttt gtttattaaa ttaaccaact 150

<210> 33
<211> 242
<212> DNA
<213> Artificial Sequence

<220>
<223> P1P2* fusion sequence

<400> 33
aaacagccga ataaggctgt ttttcttttt tttatgtttt agaataagtg gtctagttta 60
ttcttgacaa aaaataatat ttgatataa ttaaataagtt gtcgtttgag acgactgact 120
ttcttattat tcatctaaaa tattattttg aaaagataac acagttattc ttgacaaaaa 180
aatataaaag tgtataatag aaaagtactc gagagttagt ttgtttatta aattaaccaa 240
ct 242

<210> 34
<211> 88
<212> DNA
<213> Artificial Sequence

<220>

<223> PL fusion sequence

<400> 34

tttcgtgaac ttttttggtg acaaagataa aaacacatga tataacttaa tcaactcgaga 60
gttagtttgt ttattaaatt aaccaact 88

<210> 35

<211> 88

<212> DNA

<213> Artificial Sequence

<220>

<223> PL10 fusion sequence

<400> 35

tttcgtgaac ttttttggtg acaaagataa aaacacatga tataattaaa tcaactcgaga 60
gttagtttgt ttattaaatt aaccaact 88

<210> 36

<211> 51

<212> DNA

<213> Artificial Sequence

<220>

<223> Xyl-T5 promoter sequence

<400> 36

tcataaaaaa tttatttgct ttcaggaaaa tttttctgta taatagattc a 51

<210> 37

<211> 51

<212> DNA

<213> Artificial Sequence

<220>

<223> Xyl-T5-DD promoter sequence

<400> 37

tcataaaaaa tttattttac atcaggaaaa tttttctgta tattagattc a 51

<210> 38

<211> 60

<212> DNA

<213> Artificial Sequence

<220>

<223> CP25 promoter sequence

<400> 38

ctttggcagt ttattcttga catggttagtg agggggctgg tataatcaca tagtactgtt 60

<210> 39

<211> 51

<212> DNA

<213> Artificial Sequence

<220>

<223> P32 promoter sequence

<400> 39

gtgagcttgg actagaaaaa aacttcacaa aatgctatac taggtaggta a 51

<210> 40
 <211> 51
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> P59 promoter sequence

<400> 40
 acattaaatt cttgacaggg agagataggt ttgatagaat ataatagttg t 51

<210> 41
 <211> 58
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> P59* promoter sequence

<400> 41
 aaaaatgaca gtttattctt gacagggaga gatagggttg atataatata atagttgt 58

<210> 42
 <211> 115
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> P1P2 promoter sequence

<400> 42
 gtctagttat tcttgacaaa aaataatatt ttgatataat taaatagttg tcgtttgata 60
 acacagttta ttcttgacaa aaaaatataa aagtgtataa tagaaaagta ctgtt 115

<210> 43
 <211> 207
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> P1P2* promoter sequence

<400> 43
 aaacagccga ataaggctgt ttttcttttt tttatgtttt agaataagtg gtctagttta 60
 ttcttgacaa aaaataatat ttgatataa tttaaagttt gtcgtttgag acgactgact 120
 ttcttattat tcatctaaaa tattattttt aaaagataac acagttattc ttgacaaaaa 180
 aatataaaaag tgtataatag aaaagta 207

<210> 44
 <211> 43
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PL promoter sequence

<400> 44
 ttttttggtg acaaagataa aaacacatga tataacttaaa tca 43

<210> 45
 <211> 53
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PL10 promoter sequence

<400> 45
 tttcgtgaac ttttttggtg acaaagataa aaacacatga tataattaaa tca 53

<210> 46
 <211> 9425
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Vector pEPEF9

<400> 46
 acccagtcag ctctctccgg tgggcgcggg gcatgactat cgtcgcgcga cttatgactg 60
 tcttctttat catgcaactc gtaggacagg tgccgcgagc gctctgggtc attttcggcg 120
 aggaccgctt tcgctggagc gcgacgatga tcggcctgtc gcttgcggta ttcggaatct 180
 tgcacgccct cgtcaagcc ttcgctcactg gtcccgccac caaacgtttc ggcgagaagc 240
 aggccattat cgcgcgcatg gcggccgacg cgtcgggcta cgtcttgctg gcgttcgcga 300
 cgcgaggctg gatggccttc cccattatga ttcttctcgc ttccgcgcgc atcgggatgc 360
 ccgcgttgca ggcctatgctg tccaggcagg tagatgacga ccatcaggga cagcttcaag 420
 gatcgtctgc ggctcttacc agcctaactt cgatcactgg accgctgac gtcacggcga 480
 tttatggcgc ctccgcgagc acatggaacg ggttggcatg gattgtaggc gccgccctat 540
 acctgtctct cctcccccgc ttgcgtcgcg gtgcatggag ccgggccacc tcgacctgaa 600
 tggaagccgg cggcacctcg ctaacggatt caccactcca agaattggag ccaatcaatt 660
 cttgcccaga actgtgaatg cgcaaaccac cccttggcag aacatatcca tcgcgtccgc 720
 catctccagc agccgcagc gcgcgatctc gggcagcgtt gggtcctggc cacgggtgcg 780
 catgatcgtg ctctgtcgtg tgaggaccgc gctaggctgg cgggggttgc ttactggtta 840
 gcagaatgaa tcaccgatac gcgagcgaac gtgaagcgac tgctgtctga aaacgtctgc 900
 gacctgagca acaacatgaa tggctctcgg ttcccggtt tcgtaaagtc tggaacgcg 960
 gaagtccctc acgtgtgct gaagtggcc gcaacagaga gtggaaccac ccggtgatac 1020
 cagcatacta tgactgagag tcaacgccat gagcggcctc atttcttatt ctgagttaca 1080
 acagtccgca ccgctgtccg gtagctcctt ccggtgggag cggggcatga ctatcgtcgc 1140
 cgcacttatg actgtcttct ttatcatgca actcgttaga cagggtgccg cagcgcccaa 1200
 cagtcccccgc gccacggggc ctgccaccat acccagccgc aaacaagcgc cctgcaccat 1260
 tatgttccgg atctgcatcg caggatgctg ctggttacc tgtggaacac ctacatctgt 1320
 attaacgaag cgctaaccgt ttttatcagg ctctgggagg cagaataaat gatcatatcg 1380
 tcaattatta cctccacggg gagagcctga gcaaaactgc ctcaggcatt tgagaagcac 1440
 acggtcacac tgcttccggt agtcaataaa ccggtaaaacc agcaatagac ataagcggct 1500
 atttaacgac cctgccctga accgacgacc gggctgaatt tgctttcgaa tttctgccat 1560
 tcatccgctt attatcactt attcaggcgt agcaccaggc gtttaagggc accaataact 1620
 gccttaaaaa aattacgccc cgccctgcc ctcatcgag tactgttgta attcattaag 1680
 cattctgcgc acatggaagc catcacagac ggcgatgatga acctgaatcg ccagcggcat 1740
 cagcaccttg tcgccttgcg tataatattt gccatggtg aaaacggggg cgaagaagtt 1800
 gtccatattg gccacgttta aatcaaaact ggtgaaactc acccagggat tggctgagac 1860
 gaaaaacata ttctcaataa accctttagg gaaataggcc aggttttcac cgtaaacgcg 1920
 cacatcttgc gaataatgt gtagaaactg ccggaatgc tcgtggtatt cactccagag 1980
 cgatgaaaac gtttcagttt gctcatggaa aacgggtgaa caagggtgaa cactatccca 2040
 tatcaccagc tcaccgtctt tcattgccat acgaattcag aggtttgatg actttgacct 2100
 ttggttttaa atttttgaaa aaaataaaaa aaggcgaagc ctattatata tttatcatat 2160
 atattttaat cttttgttct tttgcgtgaa aaaaaaggca gtgttttcgc tagttataga 2220
 aattaaacag tcacaaaaat cgatgtatag agtcacaaaa atcgattttt gtgactctat 2280
 gcctgatgt atacgagtc caaaaatcga tgtacacagt acgacttttg tatttgtgta 2340
 ctgtatatag tataataaaa gcatagagaa aactcactat gaaatgactt tctctatgct 2400
 actactaaaa cagcgaagg agcgtattta tactatgatt ataccagaaa aacaaaacaa 2460

```

acaaaagcag gtgctgacct tgaatgaatt agaaaaacgc aaagtagtgg agcataacgc 2520
ttaaattcaa agtgtggcta aaatgcaaaa aacagcttta aaaatgtttg agttagccgt 2580
gtcttgattt gataccgaag aacccccaaa aaataatact gtttatcttt taaaatctga 2640
actttttaaa ttttttgaag tatcgtctag cagtaagcat agtcaattca aagaagcagt 2700
taattatatg caaaagcaag cctttttcaa tattaaagct gataagaaac taggtattga 2760
gtatgaaagc attgtgccta ttccttatgt aaaatggaac gattataacg atgaagtaac 2820
aatagctttt gaccaagcca ttatgcccta tctgattgat ttaaaggcag aatttactca 2880
atataaaatt tccgaactac aaaaactcaa tagcaagtat tcgattattt tgtatcgttg 2940
gttatccatg aattataacc aatacgaaca ttacagcgtg aaagggggac ggagagttga 3000
acaagtggaa tcctaccgca acccctcaat aaaagtgaaa gaattgcgag aaataactga 3060
tacaataaat gaacatcaac attttcccca ttttgaaact agagtattaa aaaaagcaat 3120
tgaagaaatc aacgctcaca cctcttttaa tgtaacctat gagaaaaaga aagcagggcg 3180
gagcattgac tctattgtct ttcattatga gaagaaacgc atggcagacg ataacagtta 3240
caagttggga gataaagact atcaagacga taaaaagcaa aaatcaagaa atgaagctga 3300
cttactaaaa caggcaatgg aaagtaaata tacaagactt ttatctgaaa atttcttaat 3360
tggcatgaat gatattatgg atacagccac aatggtaggc ttacagaaga atgtttatcc 3420
gctttatgac gaactgaaag aattaagagg gctaaatggt gtcaaagacc acttgtctta 3480
tgttgccagc aaaaaagaag cctatttctaa acgtaatgta gcgaaatctt ttaaaaaagc 3540
cattgaaaca tatctgccaa cggttaaaag gcaggactta aacctgaggt gaagacttaa 3600
aaacgataaa agagttggcg gacgagctag gactagctaa acagaccgtt caataccatt 3660
taaagttttt gccgaccgaa aatcgacaaa aaaacaatag tggttcgatc gtcctaaacc 3720
ttgaagaaca aaagttttatc aaaaaaagac taacaaaaat aaaaaaagtt caaccgaccg 3780
aaaatcgaca agaaaaaatg aagaatttaag cgtaaatagt cttttgttag gaattccgga 3840
tgagcattca tcaggcgggc aagaatgtga ataaaggccg gataaaactt gtgcttattt 3900
ttcttttacgg tctttaaaaa ggccgtaata tccagctgaa cggctcggtt ataggtacat 3960
tgagcaactg actgaaatgc ctcaaatgt tctttacgat gccattggga tatatcaacg 4020
gtggtatatac cagtgaattt tttctccatt ttagcttcct tagctcctga aaatctcgat 4080
aactcaaaaa atacgcccgg tagtgatctt atttcattat ggtgaaagtt ggaacctctt 4140
acgtgccgat caacgtctca ttttcgcaa agtttgccc agggcttccc ggtatcaaca 4200
gggacaccag gatttattta ttctgcgaag tgatcttcctg tcacaggtat ttattcggcg 4260
caaagtgcgt cgggtgatgc tgccaactta ctgatttagt gtatgatggt gtttttgagg 4320
tgctccagtg gcttctgttt ctatcagctg tccctcctgt tcagctactg acgggggtgt 4380
gcgtaacggc aaaagcaccg ccggacatca gcgctagcgg agtgtatact ggcttactat 4440
gttggaactg actgaagggt cagtgaagtg cttcatgtgg caggagaaaa aaggctgcac 4500
cgggtgcgtca gcagaatatg tgatacagga tatattccgc ttctcgcgc actgactcgc 4560
tacgctcggg cgttcgactg cggcgagcgg aaatggctta cgaacggggc ggagatttcc 4620
tggaagatgc caggaagata cttaacaggg aagtgaaggg gccgcggcaa agccgttttt 4680
ccataggctc cgccccctg acaagcatca cgaaatctga cgctcaaatc agtgggtggc 4740
aaacccgaca ggactataaa gataccaggc gtttccccct ggcggtctcc tctgtcgtc 4800
tctgttccct gcctttccgt ttaccgggtg cattccgctg ttatggccgc gtttgtctca 4860
ttccagccct gacactcagt tccgggtagg cagttcgctc caagctggac tgtatgcacg 4920
aacccccctg tcagtcggac cgctgcgcct tatccggtaa ctatcgtctt gagtccaacc 4980
cggaaagaca tgcaaaagca ccactggcag cagccactgg taattgattt agaggagtta 5040
gtcttgaagt catgcgccg ttaaggctaa actgaaagga caagttttgg tgactgcgct 5100
cctccaagcc agttacctcg gttcaaagag ttggtagctc agagaacctt cgaaaaacgc 5160
ccctgcaagg cggttttttc gttttcagag caagagatta cgcgagacc aaaacgatct 5220
caagaagatc atcttattaa tcagataaaa tatttctaga tttcagtga atttatctct 5280
tcaaatgtag cactgaagt cagccccata cgataaagt tgtaattctc atgtttgaca 5340
gcttatcatc gatacaaatt cccgtaggc gctagggacc tctttagctt cttggaagct 5400
gtcagtagta tatctaataa tttatctcca ttccttttag taacgtgtaa ctttccaaat 5460
ttaaaaaagc gactcataga attatttctt cccgttaaata aatagataac tattaataat 5520
agacaatact tgctcataag taatgggtact taaattgttt actttggcgt gtttcattgc 5580
ttgatgaaac tgattttttg taacagttg acgatattct cgattgacct attttgaaac 5640
aaagtacgta tatagcttcc aatattttatc tggaacatct gtggtatggc gggtaagttt 5700
tattaagaca ctggttactt ttggttttagg atgaaagcat tccgctggca gcttaagcaa 5760
ttgctgaact gagacttgag tgtgcaagag caaccctagt gttcggtgaa tatccaaggt 5820
acgctgttag aatccttctt caacaatcag atagatgtca gacgcacggc tttcaaaaac 5880
cactttttta ataatttgtg tgcttaaatg gtaaggaata ttccaacaa ttttataacct 5940
ctgtttgtta gggaattgaa actgtagaat atcttggtga attaaagtga cacgagtatt 6000
cagttttaat ttttctgacg ataagttgaa tagatgactg tctaattcaa tagacgttac 6060
ctgtttactt attttagcca gtttcgtcgt taaatgccct ttacctgttc caatttcgta 6120

```

```

aacggtatcg gtttctttta aattcaattg ttttattatt tggttgagta ctttttcaact 6180
cgtaaaaaag ttttgagaat attttatatt tttgttcatg taatctctcc tgaagtgatt 6240
acatctataa acaaatacag aagttaaacg atttgtttgt aatttttagtt atctgtttta 6300
aaagtcataa gattagtcac tggtaggaat taatctaacg tatttattta tctgcgtaat 6360
cactgttttt agtctgtttc aaaacagtag atgtttttatc tacattacgc atttggaata 6420
ccaacatgac gaatccctcc ttcttaatta caaattttta gcatctaatt taacttcaat 6480
tcctattata caaaatttta agataatgca ctatcaacac actcttaagt ttgcttctaa 6540
gtcttatttc cataacttct tttacgtttc cgccattctt tgctgtttcg atttttatga 6600
tatggtgcaa gtcagcacga acacgaaccg tcttatctcc cattatatct tttttgcac 6660
tgattggtgt atcatttcgt ttttcttttt gtgcgcctaa atttcccaca atcactcact 6720
tctttctatt tcttcttatt cttattttat catcaacaat cacaaatcac ttgtgatttg 6780
tgataagtga tttgtgataa gtgatttggt attaatatat aaaagccctc tttaaagggc 6840
ttttatgttt attttgagaa agatataaaa tcaatatatc ccttttcccc aatttttaca 6900
acggcattgt agggcttttt ctctttgttt ttgatttcct tttaccaggg tttcttttcc 6960
ctccagtaat tcttttacct tcgttttggt tagttttttc tttctaaaaat gttcagctaa 7020
agtaaaactta cattcaggat aatttgaaca accataaaac gattttttta atacaatatt 7080
gttgccacac ttaggacatt ttctacaat acttttttct gcttcttttt ctttctgttc 7140
ctggtaatca gaaaaattta gtttttctat atcgtaggt acagcttcca gtaaataaac 7200
aatgaatttt ttgatattcg taataaagtt ctcttgattg ccttctcttt taccgatttt 7260
ttttaaatac gtttcccat tagccgtcat ttcagcactc gttaaaaggt gctgactttc 7320
aactgcctgg cacaataatt ttcttttttc agttacaaca agcttgctg caggtcgacg 7380
ataaccacat aacagtcata aaaccactcc tttttaacaa actttatcac aagaaatatt 7440
ttggcattct acgactataa cttaaaattta tttttttac tttataatat ataattgatt 7500
atagaataat gttgctcata tcgtttgcca acatctagta ctcaaattac actatgttac 7560
acttggtaat attaacgaa cttcccctgt ccaaattaga taagaggtaa taataaatgg 7620
aaaataattt tatagtaaat gaaaatgaga agcgtgtatt aaaacaaatt ttcaataaca 7680
gcaatatttc acgaacacaa atatcgaaga atttagaact taataaagct actatttcta 7740
acattctgaa caacttaaaa cacaagagtt tagttaatga agtaggagaa ggtaatagta 7800
ctaaaagtgg tggacgaaag cctattttac tcgaaattaa ccaaaaatat ggctactata 7860
tttctatgga tttaacatat atttaattta tttttttac caactacttt gatgctacta 7920
tattaaagca agattcctac gaattaaatg ataaaaatgt aagcagtata ttacaaattt 7980
taaaatctaa tataaacgtc tcagaaaaat atgatacgtt atatgggtta cttggtatat 8040
ctatatccat acacggtatc gttgacgatg agcaaaacat aatcaatctt ctttttcata 8100
aaaatgagaa acgcacattt accgatgaat taaagtcatt cacaaatgtt cctgtcgtta 8160
tagaaaatga agcaaattta tcagcgctat atgaaaaaag tttatatatt aattcaaaca 8220
taaaataatt gattacttta agtattcaca agggtagagg cgctggcatc ataataaata 8280
aaaaacttta tcgtggctca aatggagagg ctggagagat aggtaagaca ttggttttgg 8340
aatctataaa taacaatgac aacaaatatt ataaaatcga agatatatgc tccaagacg 8400
ctttaatata gaaaataaat aataggttg gcgtcacatt gacgtttaca gaactaatcc 8460
aatattacaa cgaaggaaat tcaattgttg ctcatgaaat taaacaattt attaatataa 8520
tgacagttct gattcataat ttgaatacac aatttaacc agacgctatt tatattaact 8580
gtcctttaat taatgaatta ccaaataatt taaatgaaat taaagagcaa ttctcctgtt 8640
tttctcaagg cagtccaatt caattacatt taactactaa tgtaaaacaa gctactttat 8700
tggttgccac tttagcaata atgcaaaaaa cattaaatat aaataacatt caaatgaata 8760
ttaaataatt acagcagtct gagttataaa atagatatct cggaaccgtg agcttggaat 8820
agaaaaaac ttcacaaaaa gctatactag gtgagtaact cgagagttag tttgtttatt 8880
aaattaacca actggacgaa ttcccgggga tccgtcgacc tgacgccaag cttgggcttt 8940
tcagcctgat acagattaaa tcagaacgca gaagcgtct gataaaacag aatttgctg 9000
gcgcagtag cgcggtgggc ccacctgacc catgccgaa ctcaagagtg aaacgccgta 9060
gcgcgatgg tagtgtgggg tctcccatg cgagagtagg gaactgccag gcatcaata 9120
aaacgaaagg ctacgtcgaa agactgggcc tttcgtttta tctgttgttt gtcggtgaac 9180
gctctcctga gtaggacaaa tccgccggga gcgatttga acgttgcgaa gcaacggccc 9240
ggagggtagc gggcaggacg cccgcataa actgccaggc atcaaattaa gcagaaggcc 9300
atcctgacgg atggcctttt tgcgtttcta caaactctt ttgtttatt ttctaaatac 9360
attcaaatat gtatccgctg atccccatcc tatcgatccc caccgatgc ccttgagagc 9420
cttca 9425

```

<210> 47

<211> 12733

<212> DNA

<213> Artificial Sequence

<220>

<223> Vector pEPEF14

<400> 47

```

aagcttgcct gcaggcgcac gataaccaca taacagtcac aaaaccactc ctttttaaca 60
aactttatca caagaaatat tttggcattc tacgactata acttaaattt atatttttta 120
ctttataata tataattgat tatagaataa tgttgctcat atcgtttgcc aacatctagt 180
actcaaatta cactatgtta cacttggtta tattaaccga acttcccctg tccaaattag 240
ataagaggta ataataaatg gaaaataatt ttatagtaaa tgaaaatgag aagcgtgtat 300
taaaacaaat tttcaataac agcaatatat cacgaacaca aatatcgaag aatttagaac 360
ttaataaagc tactatttct aacatttctga acaacttaaa acacaagagt ttagttaatg 420
aagtaggaga aggtaatagt actaaaagtg gtggacgaaa gcctatttta ctcgaaatta 480
acaaaaaata tggctactat atttctatgg atttaacata tgattccgtt gaattaatgt 540
acaactactt tgatgctact atattaaagc aagatttcta cgaattaaat gataaaaatg 600
taagcagtat attacaaatt ttaaaatcta atataaacgt ctcagaaaaa tatgatacgt 660
tatatgggtt acttgggtata tctatatcca tacacgggat cgttgacgat gagcaaaaaa 720
taatcaatct tccttttcat aaaaatgaga aacgcacatt taccgatgaa tlaaagtcac 780
tcacaaatgt tcctgtcgtt atagaaaatg aagcaaattt atcagcgcta tatgaaaaaa 840
gtttatatat taattcaaac ataaataatt tgattacttt aagtattcac aagggtatag 900
gcgctggcat cataataaat aaaaaacttt atcgctggctc aaatggagag gctggagaga 960
taggtaagac attgggtttg gaatctataa ataacaatga caacaaatat tataaaatcg 1020
aagatatatg ctcccagac gctttaatac agaaaataaa taataggttg ggcgtcacat 1080
tgacgtttac agaactaatc caatattaca acgaaggaaa ttcaattgtt gctcatgaaa 1140
ttaaacaatt tattaataaa atgacagttc tgattcataa tttgaataca caatttaacc 1200
cagacgctat ttatatatac tgtcctttta ttaatgaatt accaaatatt ttaaatgaaa 1260
ttaaagagca attctcctgt ttttctcaag gcagtcacat tcaattacat ttaactacta 1320
atgtaaaaaca agctacttta ttgggtggca ctttagcaat aatgcaaaaa acattaaata 1380
taaataacat tcaaatgaat attaaataat tacagcagtc tgagttataa aatagatatac 1440
tcggaccgac attaaattct tgacaggag agataggttt gatagaatat aatagttgtc 1500
tcgagagtta gtttgtttat taaattaac aactggatcc cgggtctaga ttagggtaac 1560
tttaacaagg gcaagctaat cttcaatggg tagatgaccc agaagtcctc 1620
cgggtcaacc aattacctgc acatagtgt caccattatt atcacgacac agcagaattc 1680
aaaacgggta gtcgcttcat caagagtcctc aatggcgctt ggcgttttta cttcgccaag 1740
acaccggctg aacgcccagt tgatttttat caaccgatt tcgatgcaac cgactttgat 1800
acgattcaag ttcccgttca tattgaacta gccggctatg gtcaaattca atacattaac 1860
acgtataacc catgggaagg taaaatttac cgtcgccac cgtataccct caatcaagat 1920
caattaacac caggcctatt cagcgacgct gcggacaaca ccgtcgctc gtacctcaa 1980
accttcgata tcgacgtatg ttttaaaggc acagctatta tcattcagtt ccaagggta 2040
gaagaagccc tgtacgtctg gttaaatggc cattttattg gctactctga agatagtttc 2100
accttctcag aatttgattt gacgcgctat attcaggacc aaggtaacgt tttagcggtt 2160
cgggtctaca aacacagtac tgcctccttt attgaagasc caagatatgt tccgtttctc 2220
tggtattttc cgtgacgtca atatactggc ggagcctgct agccatatta ctgatttggg 2280
catccgacca gttccaaatg ccaatctcaa aagtgggtgag ctcaacatca ctactaaagt 2340
aaccggcgaa ccagccactt tagcgtgac cgttaaagac catgacgggc gagtactgac 2400
gagtcaaacg caaacgggta gtggcagtg aacctttgat actatgttat tcgaccaact 2460
gcacttgtgg tcaccacaaa cgccgtatct ctatcaattg acaattgaag tttacgatgc 2520
tgatcaccaa ctcttggagc tcgtcccata tcagtttggg ttccggacgg tcgagctgcg 2580
cgatgacaaa gtcattttac tcaacaataa acggttgggt atcaacgggg ttaaccggca 2640
cgaatggaac gccacaccg gtcgcgttat cagtatggct gatatgcgcg ctgatatcca 2700
aaccatgtta gctaacaata tcaatgccga tcggacctgc cattatcctg atcaattacc 2760
ttggtatcaa ttatgtgacg aggccggtat ctacctaatg gccgaaacca acctcgaatc 2820
gcacgggtca tggcaaaaaga tgggggctat cgagccttct tacaatgttc ctggcgataa 2880
tccacatgga ccagcagcgg tgatcgaccg ggcccgttca aactacgaat ggtttaaaaa 2940
ccacccatca atcatttttt ggctcatttg caatgaatcg tatgctggcg aagatatcgc 3000
ggcgatgcag gctttttata aagaacacga tgattcacga ctcgtccact acgaaggcgt 3060
tttctacaca ccagaattaa aagatcgcat ttctgatgtt gaaagtcgga tgtacgaaaa 3120
gccccaaaat attgtagctt acttgggaaga taaccaacc aaacctttcc taaattgtga 3180
atatatgcat gacatgggga attctctggg cggtatgcaa tcatataatg atttgatcga 3240

```

```

caagtatcca atgtatcaag gtggctttat ttgggacttt attgatcaag cactcttcgt 3300
tcatgaccca atcaccgacc aagacgtgct ccggtatggc ggtgatttcg acgaacgcca 3360
ctccgattat gcattctccg gtaacggctt aatgtttgcc gaccggacac caaaaccagc 3420
aatgcaagag gtgaaatatt attatggctt acacaaataa tcaactacac gttattttacg 3480
gcgacgggag cttaggacta cagggggcta atttccacta cctcttttagc tacgaacgtg 3540
ggggacttga atcactcgtc gtcaacgata aagagtggct ctatcgtaca cccacgcccc 3600
tgttttggcg ggcgacaacc gataatgata acggtagcgg cttttcagtc aaatccgcac 3660
agtgggtacgc ggcgataag ttctcaactt gtcaagatat cgaattgacg gttgacgacc 3720
aaccagtcac acggttacca atcgcgccac tcaataacaa atacacggat cacgaaatcg 3780
ccacgaaagt ctctctggct taccacttcg ttaccacgac cgttcctagt accatcgtca 3840
cagtgaacta tacggtgaca gcagacggtc agatcaatat cgccacccat tatagcggtc 3900
agtctgattt gccagagcta cccgcatttg gtctgcggtt tatcatgcca actaccgcca 3960
ccggcttcga ctataccggg ttgtccgggt agacttatcc tgaccggctg gctggcgcaa 4020
cacacgggca attccaagtt gacagtctgc cagtcaacac atacttggtc ccacaagaat 4080
gcggcatgca catgcaaact gaacaagtga cagtaacgcg atcaacaaca caaaataacg 4140
ctgaccacga caacacacca tttagtttaa catttagcca aaccgatgca ccattcgctt 4200
tcagctgcct tccctatacc gctgctgaac tagaaaacgc aacacacatg gaagaattac 4260
cattagcagc gcgaacggtc ttatcaatct acggtgccgt tcgtggggtc ggtggcattg 4320
acagttgggg aacggacgta gaagcccat atcatatcct cgctaataca gacattgact 4380
tcagctttta cattcatttc taaaaattaa tttagtttca aaagaatcgc tccggcagtt 4440
atttgccaga gcgttctttt aatatgtttt atggctgagc tttagtcctt tgaagtgggg 4500
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4560
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4620
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4680
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4740
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4800
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4860
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4920
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 4980
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 5040
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 5100
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 5160
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 5220
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 5280
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 5340
gggggggggtc gaccgatgcc cttgagagcc ttcaaccacg tcagctcctt ccggtgggcg 5400
cggggcatga ctatcgtcgc cgcacttatg actgtcttct ttatcatgca actcgtagga 5460
caggtgcggc cagcgtctcg ggtcattttc ggcaggacc gctttcgctg gacgcgcgac 5520
atgatcgccc tgcgtctgc ggtattcgga atcttgacg ccctcgctca agccttcgtc 5580
actggtcccg ccaccaaacg ttctcgcgag aagcaggcca ttatcgccg catggcggcc 5640
gacgcgctgg gctacgtctt gctggcgttc gcgacgcgag gctggatggc cttccccatt 5700
atgattcttc tcgcttcgg cgccatcgg atgcccgcgt tgcaggccat gctgtccagg 5760
caggtagatg acgaccatca gggacagctt caaggatcgc tcgcggctct taccagccta 5820
acttcgatca ctggaccgct gatcgtcacg gcgatttatg ccgcctcggc gagcacatgg 5880
aacgggttg catgattgt aggcgcggcc ctataccttg tctgcctccc cgcgttgctg 5940
cgcggtgcat ggagccgggc cacctcgacc tgaatggaag ccggcggcac ctgcctaacg 6000
gattcaccac tccaagaatt ggagccaatc aattcttgcg gagaactgtg aatgcgcaaa 6060
ccaacccttg gcagaacata tccatcgcgt ccgccatctc cagcagccgc acgcggcgca 6120
tctcgggcag cgttggttcc tggccacggg tgcgcgatg cgtgctcctg tcgttgagga 6180
cccggctagg ctggcggggt tgccttactg gttagcagaa tgaatcaccg atacgcgagc 6240
gaacgtgaag cgactgctgc tgcaaaacgt ctgcgacctg agcaacaaca tgaatggtct 6300
tcggtttccg tgtttcgtaa agtctgaaa cgcggaagtc ccctacgtgc tgctgaagtt 6360
gcccgaaca gagagtggaa ccaaccgtg ataccagat actatgactg agagtcaacg 6420
ccatgagcgg cctcatttct tattctgagt tacaacagtc cgcaccgctg tccggtagct 6480
ccttcgggtg ggcgcggggc atgactatcg tcgccgact tatgactgtc ttctttatca 6540
tgcaactcgt aggacaggtg ccggcagcgc ccaacagtcc cccggccacg gggcctgcca 6600
ccatacccac gccgaaacaa gcgccctgca ccattatgtt ccggatctgc atcgaggat 6660
gctgctggct accctgtgga acacctacat ctgtattaac gaagcgctaa ccgtttttat 6720
caggctctgg gaggcagaat aaatgatcat atcgtcaatt attacctcca cggggagagc 6780
ctgagcaaac tggcctcagg catttgagaa gcacacggtc acactgcttc cggtagtcaa 6840
taaaccggta aaccagcaat agacataaag ggctatttaa cgaccctgcc ctgaaccgac 6900

```

gaccgggtcg	aatttgcttt	cgaatttctg	ccattcatcc	gcttattatc	acttattcag	6960
gcgtagcacc	aggcgtttaa	gggcaccaat	aactgcctta	aaaaaattac	gccccgccct	7020
gccactcatc	gcagtactgt	tgtaattcat	taagcattct	gccgacatgg	aagccatcac	7080
agacggcatg	atgaacctga	atcgccagcg	gcacgagcac	cttgctgcct	tgcgataaat	7140
atttgcccat	ggtgaaaacg	ggggcgaaag	agttgtccat	attggccacg	tttaaatcaa	7200
aactggtgaa	actcacccag	ggattggctg	agacgaaaaa	catattctca	ataaaccttt	7260
tagggaaata	ggccagggtt	tcaccgtaac	acgccacatc	ttgcgaatat	atgtgtagaa	7320
actgccggaa	atcgctcggt	tattcactcc	agagcgatga	aaacgtttca	gtttgctcat	7380
ggaaaacggt	gtaacaagg	tgaacactat	cccatatcac	cagctcaccg	tctttcattg	7440
ccatacgaat	tcagagggtt	gatgactttg	acctttgggt	ttaaattttt	gaaaaaata	7500
aaaaaaggcg	aagcctatta	tatatattatc	atatataatt	taatcttttg	ttcttttgcg	7560
tgaaaaaaa	ggcagtgttt	tcgctagtta	tagaaattaa	acagtcacaa	aaatcgatgt	7620
atacagtcac	aaaaatcgat	ttttgtgact	ctatgcatcg	atgtatacga	gtcacaaaaa	7680
tcgatgtaca	cagtacgact	tttgattttg	tgtactgtat	atagtataat	aaaagcatag	7740
agaaaactca	ctatgaaatg	actttctcta	tgtactact	aaaacacgca	aaggagcgta	7800
tttatactat	gattatacca	gaaaaacaaa	acaaacaaaa	gcagggtgctg	accttgaatg	7860
aattagaaaa	acgcaaagta	gtggagcata	acgctttaat	tcaaagtgtg	gctaaaatgc	7920
aaaaaacagc	tttaaaaatg	tttgagtttag	ccgtgtcttg	tattgatacc	gaagaacccc	7980
caaaaaataa	tactgtttat	cttttaaaat	ctgaactttt	taaatttttt	gaagtatcgt	8040
cagacgtcac	gcatagtcga	ttcaaagaag	cagtttaatta	tatgcaaaag	caagcctttt	8100
tcaatattaa	agctgataag	aaactaggtta	ttgagtatga	aagcattgtg	cctattcctt	8160
atgtaaaaatg	gaacgattat	aacgatgaag	taacaatacg	ttttgaccaa	gccattatgc	8220
cctatctgat	tgattttaag	gcagaattta	ctcaatataa	aatttccgaa	ctacaaaaac	8280
tcaatagcaa	gtattcgatt	attttgtatc	gttggttatc	catgaattat	aaccaatacg	8340
aacattacag	cgtgaaagg	ggacggagag	ttgaacaagt	ggaatcctac	cgaacccct	8400
caataaaagt	gaaagaattg	cgagaaataa	ctgatacaat	aatgaacat	caacattttc	8460
cccattttga	aaactagta	ttaaaaaaag	caattgaaga	aatcaacgct	cacacctctt	8520
ttaatgtaac	ctatgagaaa	aagaaagcag	ggcggagcat	tgactctatt	gtctttcata	8580
ttgagaagaa	acgcatggca	gacgataaca	gttacaagtt	gggagataaa	gactatcaag	8640
acgataaaaa	gcaaaaatca	agaaatgaag	ctgacttact	aaaacaggca	atggaaagta	8700
aatatacaag	acttttatct	gaaaatttct	taattggcat	gaatgatatt	atggatacac	8760
ccacaatggt	aggcttacag	aagaatgttt	atccgcttta	tgacgaactg	aaagaattaa	8820
gagggtctaa	tggtgtcaaa	gaccacttgt	cttatgtttg	cagcaaaaaa	gaagcctatt	8880
ctaaacgtta	tgtagcgaaa	tatcttaaaa	aagccattga	acaatatctg	ccaacggtta	8940
aaaggcagga	cttaaacctat	gagtgaaagc	ttaaaaacga	taaaagagtt	ggcggacgag	9000
ctaggactag	ctaaacagac	cgttcaatac	cattttaaagt	ttttgccgac	cgaaaatcga	9060
caaaaaaaca	atagtgtgtc	gacgtctcta	aaccttgaag	aacaaaagtt	tatcaaaaaa	9120
agactaacia	aaataaaaaa	agttcaaccg	accgaaaaatc	gacaagaaaa	aatgaagaat	9180
taagcgttaa	tagtcttttg	ttaggaattc	cggatgagca	ttcatcaggc	gggcaagaat	9240
gtgaataaag	gccggataaa	acttgtgctt	atttttcttt	acggtcttta	aaaaggccgt	9300
aatatccagc	tgaacggctc	ggttataggt	acattgagca	actgactgaa	atgcctcaaa	9360
atgttcttta	cgatgccatt	gggatataatc	aacggtggta	tatccagtga	tttttttctc	9420
catttttagct	tccttagctc	ctgaaaatct	cgataactca	aaaaatacgc	ccggtagtga	9480
tcttatttca	ttatggtgaa	agttggaacc	tcttacgtgc	cgatcaacgt	ctcattttcg	9540
ccaaaagtgt	gcccagggtc	tcccgggtatc	aacagggaca	ccaggattta	tttattctgc	9600
gaagtgatct	tccgtcacag	gtattttatc	ggcgcaaagt	gcgtcgggtg	atgctgcca	9660
cttactgatt	tagtgtatga	tggtgttttt	gaggtgctcc	agtggcttct	gtttctatca	9720
gctgtccctc	ctgttcagct	actgacgggg	tggtgcgtaa	cggcaaaagc	accgccggac	9780
atcagcgcta	gcggagtcta	tactggctta	ctatgttggc	actgatgagg	gtgtcagtga	9840
agtgtctcat	gtggcaggag	aaaaaaggct	gcaccggtgc	gtcagcagaa	tatgtgatac	9900
aggatataat	ccgcttctc	gctcactgac	tcgctacgct	cggctgcttcg	actgcggcga	9960
gcggaaatgg	cttacgaacg	gggaggagat	ttcctggaag	atgccaggaa	gatacttaac	10020
agggaaagtga	gagggccgcg	gcaaagccgt	ttttccatag	gctccgcccc	cctgacaagc	10080
atcacgaaat	ctgacgctca	aatcagtggt	ggcgaaaccc	gacaggacta	taaagatacc	10140
aggcgtttcc	ccctggcggc	tccctcgtgc	gctctcctgt	tctgccttt	cggtttaccg	10200
tggtcattcc	gctgttatgg	cccggtttgt	ctcatccac	gcctgacact	cagttccggg	10260
taggcagttc	gctccaagct	ggactgtatg	cacgaacccc	ccgttcagtc	cgaccgctgc	10320
gccttatccg	gtaactatcg	tcttgagtcc	aacccggaag	gacatgcaaa	agcaccactg	10380
gcagcagcca	ctggtaattg	atthagagga	gttagtcttg	aagtcatgcg	ccggttaagg	10440
ctaaactgaa	aggacaagtt	ttggtgactg	cgctcctcca	agccagttac	ctcggttcaa	10500
agagtttgta	gctcagagaa	ccttcgaaaa	accgccctgc	aaggcggttt	tttcgttttc	10560

```

agagcaagag attacgcgca gaccaaacg atctcaagaa gatcatctta ttaatcagat 10620
aaaatatattc tagatttcag tgcaatttat ctcttcaa atgtagcacctg aagtcagccc 10680
catacgatat aagttgtaat tctcatgttt gacagcttat catcgataca aattccccgt 10740
aggcgctagg gacctcttta gcttcttgga agctgtcagt agtatatcta ataattttatc 10800
tccattccct ttagtaacgt gtaactttcc aaatttaaaa aagcgactca tagaattatt 10860
tcctcccgtt aaataataga taactattaa aaatagacaa tacttgctca taagtaatgg 10920
tacttaaatt gtttactttg gcgtgtttca ttgcttgatg aaactgattt ttagtaaaca 10980
gttgacgata ttctcgattg acccattttg aaacaaagta cgtatatagc ttccaatatt 11040
tatctggaac atctgtggtg tggcgggtaa gttttattaa gacactgttt acttttggtt 11100
taggatgaaa gcattccgct ggcagcttaa gcaattgctg aatcgagact tgagtgtgca 11160
agagcaaccc tagtgctcgg tgaatatcca aggtacgctt gtagaatcct tcttcaacaa 11220
tcagatagat gtcagacgca cggctttcaa aaaccacttt ttaataaatt tgtgtgctta 11280
aatggtaagg aatattccca acaattttat acctctgttt gttagggaat tgaaactgta 11340
gaatatcttg gtgaattaaa gtgacacgag tattcagttt taatttttct gacgataagt 11400
tgaatagatg actgtcta atcaatagac ttactgtttt acttatttta gccagtttctg 11460
tcgttaaatg ccttttacct gttccaattt cgtaaacggt atcggtttct tttaaattca 11520
attgttttat tatttggttg agtacttttt cactcgttaa aaagttttga gaatatttta 11580
tatttttggt catgtaactt ctctgaagt gattacatct ataaacaaat acagaagtta 11640
aacgatttgt ttgtaatttt agttatctgt ttaaaaagtc ataagattag tcaactggtag 11700
gaattaatct aacgtattta tttatctcgg taactactgt ttttagtctg tttcaaaaca 11760
gtagatgttt tatctacatt acgcatlttg aataccaaca tgacgaatcc ctcttctta 11820
attacaaatt tttagcatct aatttaactt caattcctat tatacaaaat tttagataa 11880
tgcaactatca acacactctt aagtttgctt ctaagtctta tttccataac ttcttttaccg 11940
tttccgccat tctttgctgt ttcgattttt atgatatggt gcaagtcagc acgaacacga 12000
accgtcttat ctccatttat atcttttttt gcaactgattg gtgtatcatt tcgtttttct 12060
ttttgtgcgc ctaaatttcc cacaatcact cacttctttc tatttcttct tattcttatt 12120
ttatcatcaa caatcacaaa tcacttgtga tttgtgataa gtgatttggt ataagtgaat 12180
tgtgattaat atataaaagc cctctttaa gggcttttat gtttattttg agaaagatat 12240
aaaaatcaata tatccctttt cccaattttt tacaacggca ttgtagggt ttttctcttt 12300
gtttttgatt tctctttacc aggttttctt ttccctccag taattctttt acattcggtt 12360
tggttagttt tttctttcta aaatgttcag ctaaagtaaa cttacattca ggataatttg 12420
aacaaccata aaacgatttt tttatataca tattgttgcc acacttagga cattttccta 12480
caatactttt ttctgcttct ttttctttct gttcctggta atcagaaaaa tttagttttt 12540
ctatctcggt aggtacagct tccagtaaat gaacaatgaa ttttttgata ttcgtaataa 12600
agttctcttg attgccttct cttttaccga ttttttttaa atacgtttcc catttagccg 12660
tcatttcagc actcggttaa aggtgctgac tttcaactgc ctggcacaat aattttcctt 12720
tttcagttac aac 12733

```

<210> 48
 <211> 9417
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Vector pEPEF14*

```

<400> 48
accgatgcc cttgagagcc ttcaaccag tcagctcctt ccggtggcg cggggcatga 60
ctatcgtcgc cgcacttatg actgtcttct ttatcatgca actcgtagga caggtgccgg 120
cagcgctctg ggtcattttc ggcgaggacc gctttcgtg gagcgcgacg atgatcgcc 180
tgtcgcttgc ggtattcgga atcttgacg ccctcgctca agccttcgtc actgggtccc 240
ccaccaaacg tttcggcgag aagcaggcca ttatcgccgg catggcggcc gacgcgctgg 300
gctacgtctt gctggcgctt gcgacgcgag gctggatggc cttccccatt atgattcttc 360
tcgcttccgg cggcatcggg atgcccgcgt tgcaggccat gctgtccagg caggtagatg 420
acgaccatca gggacagctt caaggatcgc tcgcggtctt taccagccta acttcgatca 480
ctggaccgct gatcgtcacg gcgatttatg ccgcctcggc gagcacatgg aacgggttg 540
catggattgt aggcgcggcc ctataccttg tctgcctccc cgcgttgctg cgcggtgcat 600
ggagccgggc caccctcgacc tgaatggaag ccggcgccac ctcgctaacg gattcaccac 660
tccaagaatt ggagccaatc aattcttgcg gagaactgtg aatgcgcaaa ccaacccttg 720

```

```

gcagaacata tccatcgct cgcacatctc cagcagccgc acgcggcgca tctcgggcag 780
cgttgggtcc tggccacggg tgcgcatgat cgtgctcctg tcgttgagga cccggctaag 840
ctggcggggt tgccttactg gttagcagaa tgaatcaccg atacgcgagc gaacgtgaag 900
cgactgctgc tgcaaacagt ctgacgactg agcaacaaca tgaatggctc tcgggtttccg 960
tggttcgtaa agtctggaaa cgcggaagt cctacgtgc tgctgaagt gcccgcaaca 1020
gagagtggaa ccaaccggtg ataccacgat actatgactg agagtcaacg ccatgagcgg 1080
cctcatttct tattctgagt tacaacagtc cgcaccgctg tccggtagct ccttcgggtg 1140
ggcgcggggc atgactatcg tcgccgact tatgactgtc ttctttatca tgcaactcgt 1200
aggacaggtg ccggcagcgc ccaacagtcc cccggccacg gggcctgcca ccataccac 1260
gccgaacaa gcgccctgca ccattatgtt ccggatctgc atcgcaggat gctgctggct 1320
acctgtgga acacctacat ctgtattaac gaagcgctaa ccgtttttat caggctctgg 1380
gaggcagaat aaatgatcat atcgtaatt attacctca cggggagagc ctgagcaaac 1440
tggcctcagg gatttgagaa gcacacggtc acactgcttc cggtagtcaa taaaccggtg 1500
aaccagcaat agacataaag ggctatttaa cgaccctgcc ctgaaccgac gaccgggtcg 1560
aatttgcttt cgaatttctg ccattcatcc gcttattatc acttattcag gcgtagcacc 1620
agggctttaa gggcaccaat aactgcctta aaaaaattac gccccgccct gccactcacc 1680
gcagtactgt tgtaattcat taagcattct gccgacatgg aagccatcac agacggcatg 1740
atgaacctga atcgccagcg gcacagcac cttgtgcctc tgcgtataat atttgccct 1800
ggtgaaaacg ggggcaaga agttgtccat attggccacg tttaaatcaa aactggtgaa 1860
actgacctga gatttgctg agacgaaaaa catattctca ataaacctt tagggaaata 1920
ggccagggtt tcaccgtaac acgccacatc ttgcgaatat atgtgtagaa actgccgga 1980
atcgctgtgg tattcactcc agagcgatga aaacgtttca gtttgcctat ggaaaacggt 2040
gtaacaagg tgaacactat cccatatcac cagctcaccg tctttcattg ccatacgaga 2100
ggtttgatga ctttgacctt tggttttaaa ttttgaaaa aaataaaaaa aggcgaagcc 2160
tattatata tttatcata ttttttaac ttttgttctt ttgctgaaa aaaaaggcag 2220
tggtttcgct agttatagaa attaaacagt cacaaaaatc gatgtataca gtcacaaaaa 2280
tcgatttttg tgactctatg catcgatgta tacgagtcac aaaaatcgat gtacacagta 2340
cgacttttgt atttgtgtac tgtatatagt ataataaaag catagagaaa actcactatg 2400
aaatgacttt ctctatgcta ctactaaaac acgcaaagga gcgtatttat actatgatta 2460
taccagaaaa acaaaacaaa caaagcagg tgctgacctt gaatgaatta gaaaaacgca 2520
aagtagtgga gcataacgct ttaattcaaa gtgtggctaa aatgcaaaaa acagctttaa 2580
aaatgttgga gttagccgtg tcttgtattg ataccgaaga acccccaaaa aataatactg 2640
tttatctttt aaaatctgaa ctttttaaat tttttgaagt atcgtctagc agtaagcata 2700
gtcaattcaa agaagcagtt aattatatgc aaaagcaagc ctttttcaat attaaagctg 2760
ataaagaaat aggtattgag tatgaaagca ttgtgcctat tccttatgta aaatggaacg 2820
attataacga tgaagtaaca atacgttttg accaagccat tatgccctat ctgattgatt 2880
taaaggcaga atttactcaa tataaaattt ccgaactaca aaaactcaat agcaagtatt 2940
cgattatttt gtatcggttg ttatccatga attataacca atacgaacat tacagcgtga 3000
aagggggcag gagagttgaa caagtggat cctaccgcaa cccctcaata aaagtgaag 3060
aattgcgaga aataactgat acaataaatg aacatcaaca ttttccccat tttgaaacta 3120
gagtataaaa aaagcaatt gaagaaatca acgttcacac ctcttttaat gtaacctatg 3180
agaaaaagaa agcaggcgag agcattgact atttgtctt tcatattgag aagaaacgca 3240
tgccagacga taacagttac aagttgggag ataaagacta tcaagacgat aaaaagcaaa 3300
aatcaagaaa tgaagctgac ttactaaaac aggcattgga aagtaaatat acaagacttt 3360
tatctgaaaa tttcttaatt ggcattgaat atattatgga tacagccaca atggtaggct 3420
tacagaagaa tgtttatccg ctttatgacg aactgaaaag attaagaggg ctaaatggtg 3480
tcaaagacca cttgtcttat gttgccagca aaaaagaagc ctattctaaa cgtaatgtag 3540
cgaaatatct taaaaagcc attgaacaat atctgccaac ggttaaaagg caggacttaa 3600
accatgatg aagacttaaa aacgataaaa gagttggcgg acgagctagg actagctaaa 3660
cagaccgttc aataccattt aaagtttttg ccgaccgaaa atcgacaaaa aaacaatagt 3720
ggttcgatcg tctaaacct tgaagaacaa aagtttatca aaaaaagact aacaaaaata 3780
aaaaaagttc aaccgaccga aaatcgacaa gaaaaaatga agaattaagc gttaatagtc 3840
ttttgttagg cggatgagca ttcacaggg gggcaagaat gtgaataaag gccggataaa 3900
acttgtgctt atttttctt acggtcttta aaaaggccgt aatatccagc tgaacggtct 3960
ggttataggt acattgagca actgactgaa atgcctcaaa atgttcttta cgatgccatt 4020
gggatatac aacggtggtg tatccagtga ttttttctc cattttagct tcttagctc 4080
ctgaaaatct cgataactca aaaaatacgc ccggtagtga tctattttca ttatgggtga 4140
agttggaacc tcttacgtgc cgatcaacgt ctcatcttcg ccaaaagtgt gccagggt 4200
tcccggtatc aacagggaca ccaggattta tttattctgc gaagtgtatc tccgtcacag 4260
gtatttatc ggcgcaaggt gcgtcgggtg atgctgcaa cttactgatt tagtgtatga 4320
tggtgttttt gaggtgctcc agtggcttct gtttctatca gctgtccctc ctgttcagct 4380

```

```

actgacgggg tgggtgcgtaa cggcaaaagc accgccggac atcagcgcta gcggagtgt 4440
tactggctta ctatgttggc actgatgagg gtgtcagtga agtgcttcat gtggcaggag 4500
aaaaaaggct gcaccgggtgc gtcagcagaa tatgtgatac aggatatatt ccgcttcctc 4560
gctcactgac tcgctacgct cggtcgttcg actgcggcga gcggaatgg cttacgaacg 4620
gggcgagat ttcttggaag atgccaggaa gatacttaac agggagtga gagggccgcg 4680
gcaaagccgt tttccatag gctccgcccc cctgacaagc atcacgaaat ctgacgctca 4740
aatcagtggg ggcgaacccc gacaggacta taaagatacc aggcgtttcc ccctggcggc 4800
tcctcgtgc gctctcctgt tcctgccttt cggtttaccg gtgtcattcc gctgttatgg 4860
ccgcgtttgt tctattccac gcctgacact cagttccggg taggcagttc gctccaagct 4920
ggactgtatg cacgaacccc ccgttcagtc cgaccgctgc gccttatccg gtaactatcg 4980
tcttgagtcc aacccgaaa gacatgcaa agcaccactg gcagcagcca ctggtaattg 5040
atthagagga gttagtcttg aagtcatgcg ccggttaagg ctaaactgaa aggacaagtt 5100
ttggtgactg cgctcctcca agccagttac ctcggttcaa agagtgtgta gctcagagaa 5160
ccttcgaaaa accgccctgc aaggcggttt tttcgttttc agagcaagag attacgcgca 5220
gacaaaaacg atctcaagaa gatcatctta ttaatcagat aaaatatctc tagatttcag 5280
tgcaatttat ctcttcaaat gtacgacctg aagtcagccc catacgatat aagttgtaat 5340
tctcatgttt gacagcttat catcgataca aattccccgt aggcgctagg gacctcttta 5400
gcttcttgga agctgtcagt agtatactta ataatttacc tccattccct ttagtaacgt 5460
gtaactttcc aaatttaaaa aagcgactca tagaattatt tcctcccggt aaataataga 5520
taactattaa aaatagacaa tacttgctca taagtaatgg tacttaaat gtttactttg 5580
gcgtgtttca ttgcttgatg aaactgattt ttagtaaaaa gttgacgata ttctcgattg 5640
acccattttg aaacaaaagta cgtatatagc ttccaatatt tatctggaac atctgtggtg 5700
tggcgggtaa gttttattaa gacactgttt acttttggtt taggatgaaa gcattccgct 5760
ggcagcttaa gcaattgctg aatcgagact tgagtgtgca agagcaaccc tagtggtcgg 5820
tgaatatcca aggtacgctt gtagaatcct tcttcaacaa tcagatagat gtcagacgca 5880
cggctttcaa aaaccacttt ttttaataatt tgtgtgctta aatggtaagg aatattccca 5940
acaattttat acctctgttt gttagggaa tgaaactgta gaatatcttg gtgaattaaa 6000
gtgacacgag tattcagttt taatttttct gacgataagt tgaatagatg actgtctaat 6060
tcaatagacg ttacctgttt acttatttta gccagtttcg tcgttaaatg ccctttacct 6120
gttccaattt cgtaaacggg atcggtttct tttaaattca attgtttat tatttggttg 6180
agtattttt cactcgttaa aaagttttga gaattttta ttttttggt catgtaactc 6240
ctcctgaagt gattacatct ataaacaaat acagaagtta aacgatttgt ttgtaatttt 6300
agttatctgt ttaaaaagtc ataagattag tcactggtag gaattaatct aacgtattta 6360
tttatctgcg taatcactgt ttttagtctg tttcaaaaaca gtagatgttt tatctacatt 6420
acgcatttgg aataccaaca tgacgaatcc ctccctctta attacaaatt tttagcatct 6480
aatttaactt caattcctat tatacaaaat tttagataa tgcactatca acacactctt 6540
aagtttgctt ctaagtctta tttccataac tttctttacg tttccgcat tctttgctgt 6600
ttcgattttt atgatattgg gcaagtcagc acgaacacga accgtcttat ctcccattat 6660
atctttttt gcactgattg gtgtatcatt tcgtttttct ttttgctgcg ctaaatttcc 6720
cacaatcact cacttctttc tatttcttct tattcttatt ttatcatcaa caatcacaaa 6780
tcacttgatg tttgtgataa gtgatttgtg ataagtattg tgtgattaat atataaaagc 6840
cctctttaaa gggcttttat gtttattttg agaaagatat aaaatcaata tatccctttt 6900
ccccattttt tacaacggca ttgtagggtt ttttctcttt gtttttgatt tccttttacc 6960
agggtttctt ttccctccag taattctttt acattcgttt tggttagttt tttctttcta 7020
aaatgttcag ctaaagttaa cttacattca ggataatttg aacaaccata aaacgatttt 7080
tttaatacaa tattgttgcc acacttagga cattttccta caatactttt ttctgcttct 7140
ttttctttct gttcctggta atcagaaaaa tttagttttt ctatatcgtt aggtacagct 7200
tccagtaaat gaacaatgaa ttttttgata ttcgtaataa agttctcttg attgccttct 7260
cttttaccga ttttttttaa atacgtttcc catttagccg tcatctcagc actcgttaaa 7320
aggtgctgac tttcaactgc ctggcacaat aattttcctt tttcagttac aacagcctgc 7380
aggtcgacga taaccacata acagtcataa aaccactcct ttttaacaaa ctttatcaca 7440
agaaatattt tggcatttca cgactataac ttaaaattat attttttact ttataatata 7500
taattgatta tagaataatg ttgctcatat cgtttgccaa catctagtag tcaaattaca 7560
ctatgtttaca cttggtaata ttaaccgaac tttccctgtc caaattagat aagaggtaat 7620
aataaatgga aaataatttt atagtaaatg aaaatgagaa gcgtgtatta aaacaaattt 7680
tcaataacag caatatttca cgaacacaaa tatcgaagaa tttagaactt aataaagcta 7740
ctatttctaa cattctgaac aacttaaaac acaagagttt agttaatgaa gtaggagaag 7800
gtaatagtac taaaagtggg ggacgaaagc ctattttact cgaaattaac caaaaatatg 7860
gtactatat ttctatggat ttaacatatg attccgttga attaatgtac aactactttg 7920
atgtacttat attaaagcaa gattcctacg aattaaatga taaaaatgta agcagtatat 7980
tacaaatttt aaaatctaata ataaacgtct cagaaaaata tgatacgtaa tatgggttac 8040

```

```

ttggtatatac tatatccata cacggtatcg ttgacgatga gcaaaacata atcaatcttc 8100
cttttcataa aaatgagaaa cgcacattta ccgatgaatt aaagtcattc acaaatgttc 8160
ctgtcggtat agaaaatgaa gcaaatttat cagcgctata tgaaaaagt ttatatatta 8220
attcaaacat aaataatttg attactttaa gtattcacia gggatataggc gctggcatca 8280
taataaataa aaaactttat cgtggctcaa atggagaggc tggagagata ggtaagacat 8340
tggttttgga atctataaat aacaatgaca acaaatatta taaaatcgaa gatatatgct 8400
cccaagacgc ttttaatacag aaaataaata ataggttggg cgtcacattg acgtttacag 8460
aactaatcca atattacaac gaaggaaatt caattgttgc tcatgaaatt aaacaattta 8520
ttaataaaaat gacagtctcg attcataaatt tgaatacaca atttaacca gacgctatct 8580
atattaactg tcctttaatt aatgaattac caaatatctt aaatgaaatt aaagagcaat 8640
tctcctgttt ttctcaaggc agtccaattc aattacattt aactactaat gtaaaacaag 8700
ctactttatt tgggtggcact tttagcaataa tgcaaaaaac attaaatata aataacattc 8760
aaatgaatat taaataatta cagcagtctg agttataaaa tagatatctc ggaccgaaaa 8820
atgacagttt attcttgaca gggagagata ggtttgatag aatataatag ttgtctcgag 8880
agttagtgtt tttattaaat taaccaactg gacgaattcc cggggatccg tcgacctgca 8940
gccaaagctt ggcttttcag cctgatacag attaaatcag aacgcagaag cggctcgata 9000
aaacagaatt tgcttgccgg cagtagccgg gtggtccccc ctgaccccat gccgaactca 9060
gaagtgaac gccgtagcgc cgatggtagt gtggggtctc cccatgagag agtagggaac 9120
tgccagtcac caaataaaac gaaaggctca gtcgaaagac tgggcctttc gttttatctg 9180
ttgttttgct gtgaacgctc tcctgagtag gacaaatccg ccgggagcgg atttgaacgt 9240
tgcaagcaa cggcccgag ggtggcgggc aggacgccc ccataaactg ccaggcatca 9300
aattaagcag aaggccatcc tgacggatgg cttttttgag tttctacaaa ctctttttgt 9360
ttatttttct aaatacatcc aaatatgtat ccgctgatcc ccatcctatc gatcccc 9417

```

<210> 49

<211> 9484

<212> DNA

<213> Artificial Sequence

<220>

<223> Vector PlP2/EF14

<400> 49

```

acccgatgcc cttgagagcc ttcaaccag tcagctcctt ccggtgggag cggggcatga 60
ctatcgctgc cgcacttatg actgtcttct ttatcatgca actcgtagga caggtgccgg 120
cagcgctctg ggtcattttc ggagaggacc gctttcgctg gagcgcgagc atgatcgcc 180
tgctcgcttg ggtattcgga atcttgacg ccctcgctca agccttcgtc actggtccc 240
ccaccaaacg ttctggcgag aagcaggcca ttatcgccgg catggcgggc gacgcgctgg 300
gctacgtctt gctggcgctc gcgacgcgag gctggatggc cttcccatc atgattcttc 360
tcgcttcggg cggcctcgag atgcccgcgt tgcaggccat gctgtccagg caggtagatg 420
acgaccatca gggacagctt caaggatcgc tcgcggtctt taccagccta acttcgatca 480
ctggaccgct gatcgtcacg gcgatttatg ccgcctcggc gagcacatgg aacgggttgg 540
catggattgt aggcgcggcc ctataccttg tctgcctccc cgcgttgctg cgcggtgcat 600
ggagccgggc cacctcgacc tgaatggaag ccggcggcac ctcgctaacc gattcaccac 660
tccaagaatt ggagccaatc aattcttgcg gagaactgtg aatgcgcaa ccaacccttg 720
gcagaacata tccatcgctg ccgcatctc cagcagccgc acgcggcgca tctcgggcag 780
cggtgggtcc tggecacggg tgcgcatgat cgtgctcctg tcgttgagga cccggctagg 840
ctggcggggt tgccttactg gttagcagaa tgaatcaccg atacgcgagc gaacgtgaag 900
cgactgctgc tgcaaacgt ctgcgacctg agcaacaaca tgaatggtct tcggtttccg 960
tgtttcgtaa agtctggaag cgcggaagtc ccctacgtgc tgctgaagtt gcccgcaaca 1020
gagagtggaa ccaaccggtg ataccacgat actatgactg agagtcaacg ccatgagcgg 1080
cctcatttct tattctgagt tacaacagtc cgcaccgctg tccggtagct ccttcgggtg 1140
ggcgcggggc atgactatcg tcgcccact tatgactgtc ttctttatca tgcaactcgt 1200
aggacaggtg ccggcagcgc ccaacagtcc cccggccacg gggcctgcca ccataccac 1260
gccgaacaa gcgccctgca ccattatgtt ccggatctgc atcgaggat gctgctggct 1320
accctgtgga acacctacat ctgtattaac gaagcgctaa ccgtttttat caggctctgg 1380
gaggcagaat aaatgatcat atcgtcaatt attacctcca cggggagagc ctgagcaaac 1440
tgccctcagg catttgagaa gcacacggtc acactgcttc cggtagtcaa taaaccggtg 1500
aaccagcaat agacataagc ggctatttaa cgaccctgcc ctgaaccgac gaccgggtcg 1560

```

```

aatttgcttt cgaatttctg ccattcatcc gcttattatc acttattcag gcgtagcacc 1620
aggcggttaa gggcaccaat aactgcctta aaaaaattac gccccgccct gccactcatc 1680
gcagtactgt tgaattcat taagcattct gccgacatgg aagccatcac agacggcatg 1740
atgaacctga atcgccagcg gcatcagcac cttgtcgctt tgcgtataat atttgcccat 1800
ggtgaaaacg gggcggaaga agttgtccat attggccacg tttaaatcaa aactggtgaa 1860
actcacccag ggattggctg agacgaaaaa catattctca ataaaccctt tagggaaata 1920
ggccagggtt tcaccgtaac acgccacatc ttgcgaatat atgtgtagaa actgccggaa 1980
atcgtcgtgg tatttactcc agagcgatga aaacgtttca gtttgctcat ggaaaacggg 2040
gtaacaaggg tgaacactat cccatatcac cagctcaccg tctttcattg ccatacgaat 2100
tcagagggtt gatgactttg acctttggtt ttaaatTTTT gaaaaaata aaaaaaggcg 2160
aagcctatta tatatttatc atatatattt taatcttttg ttcttttgcg tgaaaaaaa 2220
ggcagtggtt tcgctagtta tagaaattaa acagtcacaa aaatcgatgt atacagtcac 2280
aaaaatcgat tttgtgact ctatgcatcg atgtatacga gtcacaaaaa tcgatgtaca 2340
cagtacgact tttgtatttg tgtactgtat atagtataat aaaagcatag agaaaactca 2400
ctatgaaatg actttctcta tgctactact aaaacacgca aaggagcgta tttatactat 2460
gattatacca gaaaaacaaa acaaacaaaa gcagggtgtg acctggaatg aattagaaaa 2520
acgcaaagta gtggagcata acgctttaat tcaaagtgtg gctaaaatgc aaaaaacagc 2580
tttaaaaatg tttgagttag ccgtgtcttg tattgatacc gaagaacccc caaaaaataa 2640
tactgtttat cttttaaaat ctgaactttt taaatttttt gaagtatcgt ctagcagtaa 2700
gcatagtcaa ttcaaagaag cagttaatta tatgcaaaag caagcctttt tcaatattaa 2760
agctgataag aaactaggta ttgagtatga aagcattgtg cctattcctt atgtaaaatg 2820
gaacgattat aacgatgaag taacaatcag ttttgacaa gccattatgc cctatctgat 2880
tgatttaaa gcaaattta ctcaatataa aattttccgaa ctacaaaaac tcaatagcaa 2940
gtattcgatt attttgtatc gttggttatt catgaattat aaccaatacg aacattacag 3000
cgtgaaaggg ggacggagag ttgaacaagt ggaatcctac cgcaacccct caataaaagt 3060
gaaagaattg cgagaataaa ctgatacaat aaatgaacat caacattttc cccattttga 3120
aactagagta ttaaaaaaag caattgaaga aatcaacgct cacacctctt ttaatgtaac 3180
ctatgagaaa aagaaagcag ggcgagcat tgactctatt gtctttcata ttgagaagaa 3240
acgcatggca gacgataaca gttacaagtt gggagataaa gactatcaag acgataaaaa 3300
gcaaaaatca agaaatgaag ctgacttact aaaacaggca atggaagta aatatacaag 3360
acttttatct gaaaatttct taattggcat gaatgatatt atggatacag ccacaatggg 3420
aggcttacag aagaatgttt atccgcttta tgacgaactg aaagaattaa gagggctaaa 3480
tggtgtcaaa gaccacttgt cttatgttgc cagcaaaaaa gaagcctatt ctaaacgtaa 3540
tgtagcgaaa tatcttaaaa aagccattga acaatatctg ccaacggtta aaaggcagga 3600
cttaaaccat gagtgaagac ttaaaaaacga taaaagagtt ggcgagcagag ctaggactag 3660
ctaaacagac cgttcaatac catttaaaagt ttttgccgac cgaaaatcga caaaaaaaca 3720
atagtgttgc gatcgtccta aaccttgaag aacaaaagtt tatcaaaaaa agactaaaca 3780
aaataaaaaa agttcaaccg accgaaaaatc gacaagaaaa aatgaagaat taagcgtaa 3840
tagtcttttg ttaggaattc cggatgagca ttcacaggc gggcaagaat gtgaataaag 3900
gccggataaa acttgtgctt atttttcttt acggtcttta aaaaggccgt aatatccagc 3960
tgaacggtct gggtataggt acattgagca actgactgaa atgcctcaa atgttcttta 4020
cgatgccatt gggatatac aacggtggtat tatccagtga tttttttctc catttttagct 4080
tccttagctc ctgaaaatct cgataactca aaaaatacgc ccggtagtga tcttatttca 4140
ttatggtgaa agttggaacc tcttacgtgc cgatcaacgt ctcatcttcg ccaaaagtgt 4200
gcccagggtc tcccgtatc aacagggaca ccaggattta tttattctgc gaagtgatct 4260
tccgtcacag gtatttatc ggcgcaaagt gcgtcgggtg atgctgcaa cttactgatt 4320
tagtgtatga tggtgtttt gaggtgctcc agtggtctt gtttctatca gctgtccctc 4380
ctgttcagct actgacggg tggtgcgtaa cggcaaaagc accgccggac atcagcgcta 4440
gcggagtgtat tactggctta ctatgttggc actgatgagg gtgtcagtga agtgcttcat 4500
gtggcaggag aaaaaaggct gcaccggtgc gtcagcagaa tatgtgatac aggatataat 4560
ccgtctctc gctcactgac tcgctacgct cggctgttcg actgcggcga gcggaaatgg 4620
cttacgaacg gggcgagat ttcttggaag atgcaggaa gatacttaac agggaagtga 4680
gaggcgccgc gcaaagccgt ttttccatag gctccgccc cctgacaagc atcacgaaat 4740
ctgacgctca aatcagtggt ggcgaaaccc gacaggacta taaagatacc aggcgtttcc 4800
ccctggcggc tccctcgtgc gctctcctgt tctgccttt cggtttacgg gtgtcattcc 4860
gctgttatgg ccgcgtttgt ctattccac gcctgacact cagttccggg taggcagttc 4920
gctccaagct ggactgtatg cacgaacccc ccgttcagtc cgaccgctgc gccttatccg 4980
gtaactatcg tcttgagtcc aaccggaaa gacatgcaaa agcaccactg gcagcagcca 5040
ctgtaattg atttagagga gttagtcttg aagtcatcg cgggttaagg ctaaactgaa 5100
aggacaagtt ttggtgactg cgtcctcca agccagttac ctcggtcaa agagttggta 5160
gctcagagaa ccttcgaaaa accgccctgc aaggcggtt tttcgtttc agagcaagag 5220

```

```

attacgcgca gaccaaaacg atctcaagaa gatcatctta ttaatcagat aaaatatttc 5280
tagatttcag tgcaatttat ctcttcaa atgtagcacctg aagtcagccc catacgatat 5340
aagttgtaat tctcatgttt gacagcttat catcgataca aattccccgt aggcgctagg 5400
gacctcttta gcttcttgga agctgtcagt agtatatcta ataatttatc tccattccct 5460
ttagtaacgt gtaactttcc aaatttaaaa aagcgactca tagaattatt tcctcccgtt 5520
aaataataga taactattaa aaatagacaa tacttgctca taagtaatgg tacttaaat 5580
gtttactttg gcgtgtttca ttgcttgatg aaactgattt ttagtaacaa gttgacgata 5640
ttctcgattg acccattttg aaacaaagta cgtatatagc ttccaatatt tatctggaac 5700
atctgtggta tggcgggtaa gttttattaa gacactgttt acttttggtt taggatgaaa 5760
gcattccgct ggcagcttaa gcaattgctg aatcgagact tgagtgtgca agagcaacc 5820
tagtggttcg tgaatatcca aggtacgctt ttgaatcct tcttcaacaa tcagatagat 5880
gtcagacgca cggctttcaa aaaccacttt ttaataaatt tgtgtgctta aatggtaagg 5940
aatattccca cgaattttat acctctgttt gttagggaat tgaaactgta gaatatcttg 6000
gtgaattaaa gtgacacgag tattcagttt taatttttct gacgataagt tgaatagatg 6060
actgtcta atcaatagacg ttacctgttt acttatttta gccagtttcg tcgttaaagt 6120
ccctttacct gttccaattt cgtaaaacgg atcgggtttct ttaaaattca attgttttat 6180
tatttggttg agtacttttt cactcgttaa aaagttttga gaatatatta tatttttggt 6240
catgtaattc ctctgaagt gattacatct ataaacaaat acagaagtta aacgatttgt 6300
ttgtaatttt agttatctgt ttaaaaagtc ataagattag tcaactggtag gaattaatct 6360
aacgtattta tttatctgct taatcactgt ttttagtctg tttcaaaaca gtagatgttt 6420
tatctacatt acgcatgttg aataccaaca tgacgaatcc ctcttctta attacaaatt 6480
tttagcatct aatttaactt caattcctat tatacaaaat ttaagataa tgcactatca 6540
acacactctt aagtttgctt ctaagtctta tttccataac tttttttacg tttccgcat 6600
tctttgctgt ttcgattttt atgatattgg gcaagtcagc acgaacacga accgtcttat 6660
ctccattat atctttttt gcaactgatt gtgtatcatt tcgtttttct tttgtgctg 6720
ctaaatttcc cacactcact cacttcttct tatttcttct tattcttatt ttatcatcaa 6780
caatcacaaa tcaactgtga tttgtgataa gtgatttggt ataagtgatt tgtgattaat 6840
atataaaagc cctctttaa gggcttttat gtttattttg agaaagatat aaaatcaata 6900
tatccctttt cccaatttt tacaacggca ttgtagggt tttctcttt gtttttgatt 6960
tccttttacc agggtttctt ttccctccag taattctttt acatcggtt tggttagttt 7020
tttctttcta aaatgttcag ctaaagttaa cttacattca ggataatttg aacaaccata 7080
aaacgatttt ttaatacaa tattgttggc acacttagga cattttccta caatactttt 7140
ttctgtctct tttctttct gttcctggta atcagaaaaa tttagttttt ctatctcgtt 7200
agtgacagct tccagttaaa gaacaatgaa ttttttgata ttctgaataa agttctcttg 7260
attgccttct cttttaccga ttttttttaa atacgtttcc catttagccg tcatttcagc 7320
actcgttaaa aggtgctgac tttcaactgc ctggcacaat aattttcctt tttcagttac 7380
aacagcctgc aggtcgacga taaccacata acagtcataa aaccactcct ttttaacaaa 7440
ctttatcaca agaaatattt tggcattcta cgactataac ttaaatattt attttttact 7500
ttataatata taattgatta tagaataatg ttgctcatat cgtttgccaa catctagtac 7560
tcaaattaca ctatgttaca cttggtaata ttaaccgaac tttccctgtc caaattagat 7620
aagaggtaat aataaatgga aaataatttt atagttaatg aaaatgagaa gcgtgtatta 7680
aaacaaaatt tcaataacag caatatttca cgaacacaaa tatcgaagaa tttagaactt 7740
aataaagcta ctatttctaa cattctgaac aacttaaaac acaagagttt agttaatgaa 7800
gtaggagaag gtaatagtag taaaagtggg ggacgaaagc ctattttact cgaaattaac 7860
caaaaatatg gctactatat ttctatggat ttaacatatg attccgttga attaatgtac 7920
aactactttg atgctactat attaaagcaa gattcctacg aattaaatga taaaaatgta 7980
agcagtatat tacaaatttt aaaatcta ataaacgtct cagaaaaata tgatacgta 8040
tatgggttac ttggtatat tatatccata cacggtatcg ttgacgatga gcaaaacata 8100
atcaatcttc cttttcataa aaatgagaaa cgcacattta ccgatgaatt aaagtcattc 8160
acaaatgttc ctgctgttat agaaaatgaa gcaaatattt cagcgtctata tgaaaaagt 8220
ttatatatta attcaaacat aaataatttg attactttaa gtattcaca gggatatagg 8280
gctggcatca taataaataa aaaactttat cgtggctcaa atggagaggc tggagagata 8340
ggtaagacat tggttttgga atctataaat aacaatgaca acaaatatta taaaatcgaa 8400
gatatatgct cccaagacgc ttaataacag aaaataaata ataggttggg cgtcacattg 8460
acgtttacag aactaatcca atattacaac gaaggaaatt caattgttgc tcatgaaatt 8520
aaacaattta ttaataaat gacagttctg attcataatt tgaatacaca atttaacca 8580
gacgctattt atattaaact tcctttaatt aatgaattac caaatatttt aaatgaaatt 8640
aaagagcaat tctcctgttt ttctcaaggc agtccaattc aattacattt aactactaat 8700
gtaaaacaag ctactttatt ggggtggcact ttagcaataa tgcaaaaaac attaaatata 8760
aataacattc aaatgaatat taaataatta cagcagctctg agttataaaa tagatatctc 8820
ggaccggctc agttattctt gacaaaaaat aatattttga tataattaaa tagttgtcgt 8880

```

```

ttgataaacac agttttattct tgacaaaaaa atataaaagt gtataataga aaagtactgt 8940
tctcgaagagt tagtttgttt attaaattaa ccaactggac gaattcccgg ggatccgtcg 9000
acctgcagcc aagcttgggc ttttcagcct gatacagatt aaatcagaac gcagaagcgg 9060
tctgataaaa cagaatttgc ctggcggcag tagcgcggtg gtcccacctg accccatgcc 9120
gaactcagaa gtgaaacgcc gttagcgcca tggtagtggt ggggtctccc atgcgagagt 9180
agggaactgc caggcatcaa ataaaacgaa aggtcagtc gaaagactgg gcctttcgtt 9240
ttatctgttg tttgtcgggt aacgctctcc tgagtaggac aaatccgccg ggagcggatt 9300
tgaacgttgc gaagcaacgg cccggagggt ggcgggcagg acgcccgcga taaactgccca 9360
ggcatcaaat taagcagaag gccatcctga cggatggcct ttttgcggtt ctacaaactc 9420
tttttgttta tttttctaaa tacattcaaa tatgtatccg ctgatcccca tcctatcgat 9480
cccc

```

<210> 50

<211> 9566

<212> DNA

<213> Artificial Sequence

<220>

<223> Vector pEPEF18

<400> 50

```

accagtcag ctcttccgg tgggcgcggg gcatgactat cgtcgcgcga cttatgactg 60
tcttctttat catgcaactc gtaggacagg tgccggcagc gctctgggtc attttcggcg 120
aggaccgctt tcgctggagc gcgacgatga tcggcctgtc gcttgccgta ttcggaatct 180
tgacagccct cgctcaagcc ttctgctact gtcccgccac caaacgtttc ggcgagaagc 240
aggccattat cgcgcgcagc ggcgcgcagc cgtctgggta cgtcttctgt gcgttcgcga 300
cgcgaggttg gatggccttc ccattatga ttcttctcgc ttccggcggc atcgggatgc 360
ccgcgttgca ggccatgctg tccaggcagg tagatgacga ccatcaggga cagcttcaag 420
gatcgctcgc ggctcttacc agcctaactt cgatcactgg accgctgata gtcacggcga 480
tttatgcgcg ctgcgcgagc acatggaacg ggttggcatg gattgtaggc gccgccctat 540
acctgtcttg cctccccgcg ttgcgtcgcg gtgcatggag ccgggccacc tcgacctgaa 600
tggaagcccg cggcacctcg ctaacggatt caccactcca agaattggag ccaatcaatt 660
cttgccggaga actgtgaatg cgcaaaccaa cccttggcag aacatatcca tcgcgtccgc 720
catctccagc agccgcagc gcgcgcatctc ggcgagcgtt gggctcctggc cacgggtgcg 780
catgatcgtg ctctgtcgt tgaggaccog gctaggtctg cggggttgcc ttactggtta 840
gcagaatgaa tcaccgatac gcgagcgaac gtgaagcgac tgctgctgca aaacgtctgc 900
gacctgagca acaaatgaa tggctctcgg ttccctgtt tcgtaaaagtc tggaaacgcg 960
gaagtccctt acgtgctgct gaagttgccc gcaacagaga gtggaaccaa ccggtgatac 1020
cacgatacta tgactgagag tcaacgccat gagcggcctc atttcttatt ctgagttaca 1080
acagtccgca ccgctgtccg gtagctcctt ccggtggcg cgggcatga ctatcgtcgc 1140
cgcacttatg actgtcttct ttatcatgca actcgtagga caggtgcccg cagcgcccaa 1200
cagtcctccg gccacggggc ctgccaccat acccagccg aaacaagcgc cctgcaccat 1260
tatgttccgg atctgcatcg caggatgctg ctggctaccc tgtggaacac ctacatctgt 1320
attaacgaag cgctaaccgt ttttatcagg ctctgggagg cagaataaat gatcatatcg 1380
tcaattatta cctccaaggg gagagcctga gcaaaactgg ctcaggcatt tgagaagcac 1440
acggtcacac tgcttccggt agtcaataaa ccggtaaacc agcaatagac ataagcggct 1500
atttaacgac cctgccctga accgacgacc gggctcgaatt tgctttcgaa tttctgccat 1560
tcacccgctt attatcactt attcaggcgt agcaccaggc gtttaagggc accaataact 1620
gccttaaaaa aattacgccc cgcctgccca ctcatcgag tactgttgta attcattaag 1680
cattctgcgg acatggaagc catcacagac ggcgatgata acctgaatcg ccagcggcat 1740
cagcaccctg tcgccttgcg tataatattt gcccatggtg aaaaacgggg cgaagaagtt 1800
gtccatattg gccacgttta aatcaaaact ggtgaaactc acccagggat tggctgagac 1860
gaaaaacata ttctcaataa accctttagg gaaataggcc aggttttcac cgtaacacgc 1920
cacatcttgc gaatatatgt gtagaaactg ccggaatcg tcgtggtatt cactccagag 1980
cgatgaaaac gtttcagttt gtcctgggaa aacgggtgaa caagggtgaa cactatccca 2040
tatcaccagc tcaccgtctt tcattgccat acgagaggtt tgatgacttt gaccttttgt 2100
tttaaatatt tgaaaaaaat aaaaaaaggc gaagcctatt atatatatt catatatatt 2160
ttaatctttt gttcttttgc gtgaaaaaaa aggcagtgtt ttcgctagtt atagaaatta 2220
aacagtcaca aaaatcgatg tatacagtca caaaaatcga tttttgtgac tctatgcata 2280

```

gatgtatacg	agtcacaaaa	atcgatgtac	acagtacgac	ttttgtat	gtgtactgta	2340
tatagtataa	taaaagcata	gagaaaactc	actatgaaat	gactttctct	atgtactactac	2400
taaaacacgc	aaaggagcgt	atttatacta	tgattatacc	agaaaaacaa	aacaaaacaaa	2460
agcagggtgct	gaccttgaat	gaattagaaa	aacgcaaagt	agtggagcat	aacgctttaa	2520
ttcaaagtgt	ggctaaaatg	caaaaaacag	ctttaaaaat	gtttgagtta	gccgtgtctt	2580
gtattgatac	cgaagaaccc	ccaaaaaata	atactgttta	tcttttaaaa	tctgaacttt	2640
ttaaattttt	tgaagtatcg	tctagcagta	agcatagtca	attcaaagaa	gcagttaatt	2700
atatgcaaaa	gcaagccttt	ttcaatatta	aagctgataa	gaaactaggt	attgagtatg	2760
aaagcattgt	gcctattcct	tatgtaaaat	ggaacgatta	taacgatgaa	gtaacaatac	2820
gttttgacca	agccattatg	ccctatctga	ttgatttaaa	ggcagaattt	actcaatata	2880
aaattttccga	tatgataaca	ctcaatagca	tagtttcgat	tattttgtat	cgttggttat	2940
ccatgaattat	taaccaatac	gaacattaca	gcgtgaaaag	gggacggaga	gttgaacaaag	3000
tggaaatccta	ccgcaacccc	tcaataaaag	tgaagaattt	gcgagaaata	actgatacaa	3060
taaatgaaca	tcaacatttt	ccccattttg	aaactagagt	attaaaaaaa	gcaattgaag	3120
aaatcaacgc	tcacacctct	tttaatgtaa	cctatgagaa	aaagaaagca	gggcgagca	3180
ttgactctat	tgtctttcat	attgagaaga	aacgcattgc	agacgataac	agttacaagt	3240
tgggagataa	agactatcaa	gacgataaaa	agcaaaaatc	aagaaatgaa	gctgacttac	3300
taaaacaggc	aatggaaagt	aaatatacaa	gacttttata	tgaaaatttc	ttaattggca	3360
tgaatgatat	tatggataca	gccacaatgg	taggcttaca	gaagaatgtt	tatccgcttt	3420
atgacgaact	gaaagaatta	agagggctaa	atgggtgtcaa	agaccacttg	tcttatgttg	3480
ccagcaaaaa	agaagcctat	tctaaacgta	atgtagcgaa	atatcttaaa	aaagccattg	3540
aacaatatct	gccaacgggt	aaaaggcagg	acttaaacca	tgagtgaaga	cttaaaaaacg	3600
ataaaaagagt	tgccggacga	gctaggacta	gctaaacaga	ccgttcaata	ccattttaaag	3660
tttttgccga	ccgaaaatcg	acaaaaaaac	aatagtgtt	cgatcgtcct	aaaccttgaa	3720
gaacaaaagt	ttatcaaaaa	aagactaaca	aaaataaaaa	aagttcaacc	gaccgaaaa	3780
cgacaagaaa	aaatgaagaa	ttaagcgtaa	atagcttttt	gttaggcgga	tgagcattca	3840
tcaggcgggc	aagaatgtga	ataaaggccg	gataaaaactt	gtgcttat	ttctttacgg	3900
tcttttaaaa	ggccgtaata	tccagctgaa	cggtctggtt	ataggtacat	tgagcaactg	3960
actgaaatgc	ctcaaaatgt	tctttacgat	gccattggga	tatatcaacg	gtggtatatc	4020
cagtgtat	tttctccatt	ttagcttcct	tagctcctga	aaatctcgat	aactcaaaaa	4080
atacgcccg	tagtgatctt	atttcattat	ggtgaaagt	ggaacctctt	acgtgccgat	4140
caacgtctca	ttttcgccaa	aagttggccc	agggcttccc	ggtatcaaca	gggacaccag	4200
gatttattta	ttctcgcaag	tgatcttccg	tcacaggtat	ttattcgcg	caaagtgcgt	4260
cgggtgatgc	tgccaactta	ctgatttagt	ctgatgtggt	gtttttgagg	tgctccagt	4320
gettctgttt	ctatcagctg	tccctcctgt	tcagctactg	acgggggtgt	gcgtaacggc	4380
aaaagcaccg	ccggacatca	gcgctagcgg	agtgtatact	ggcttactat	gttggcactg	4440
atgagggtgt	cagtgaagt	cttcatgtgg	caggagaaaa	aaggctgcac	cgggtgcgtca	4500
gcagaatatg	tgatacagga	tatatccgcg	ttcctcgctc	actgactcgc	tacgctcgg	4560
cgttcgactg	cggcgagcgg	aaatggctta	cgaacggggc	ggagatttcc	tggaagatgc	4620
caggagaagga	ctaacacagg	aagtgaaggg	gccgcggcaa	agccgttttt	ccataggctc	4680
cgccccctg	ccaagcatca	cgaaatctga	cgctcaaatc	agtgggtggc	aaaccgaca	4740
ggactataaa	gataccaggc	gtttccccct	ggcggtctcc	tcgtgcgctc	tcctgttctc	4800
gcctttcggt	ttaccgggtg	cattccgctg	ttatggccgc	gtttgtctca	ttccacgcct	4860
gacactcagt	tccgggttag	cagttcgctc	caagctggac	tgtatgcacg	aacccccctg	4920
tcagtcggac	cgtgcgcct	tatccggtaa	ctatcgtctt	gagtcacaac	cggaaagaca	4980
tgcaaaagca	ccactggcag	cagccactgg	taattgattt	agaggagtta	gtcttgaaagt	5040
catgcgccgg	ttaaggctaa	actgaaagga	caagttttgg	tgactgcgct	cctccaagcc	5100
agttacctcg	gttcaaaag	ttggtagctc	agagaacctt	cgaaaaaccg	ccctgcaagg	5160
cggttttttc	gttttcagag	caagagatta	cgcgcagacc	aaaacgatct	caagaagatc	5220
atcttattaa	tcagataaaa	tatttctaga	tttcagtgc	atctatctct	tcaaatgtag	5280
cacctgaagt	cagccccata	cgatataagt	tgtaattctc	atgtttgaca	gcttatcatc	5340
gatacaaaat	ccccgtaggc	gctagggacc	tcttttagctt	cttggaagct	gtcagtagta	5400
tatctaataa	tttatctcca	ttcccttttag	taacgtgtaa	ctttccaaat	ttaaaaaagc	5460
gactcataga	attatttctt	cccgttaa	aatagataac	tattaaaaat	agacaatact	5520
tgctcataag	taattggtact	ttaaattgtt	actttggcgt	gtttcattgc	ttgatgaac	5580
tgatttttag	taaacagttg	acgatattct	cgattgacct	atgttgaaac	aaagtacgta	5640
tatagcttcc	aatattttat	tggaaacatct	gtgggtatggc	gggtaagttt	tattaaagaca	5700
ctgtttactt	ttggtttagg	atgaaagcat	tccgctggca	gcttaagcaa	ttgctgaatc	5760
gagacttgag	tgtgcaagag	caaccctagt	gttcgggtgaa	tatccaaggt	acgctttag	5820
aatccttctt	caacaatcag	atagatgtca	gacgcacggc	tttcaaaaac	cactttttta	5880
ataatttgtg	tgcttaaatg	gtaaggaata	ttcccaacaa	ttttatacct	ctgtttgtta	5940

```

gggaattgaa actgtagaat atcttggtga attaaagtga cacgagtatt cagttttaat 6000
ttttctgacg ataagttgaa tagatgactg tctaattcaa tagacgttac ctgtttactt 6060
attttagcca gtttcgtcgt taaatgccct ttacctgttc caatttcgta aacggtatcg 6120
gtttctttta aattcaattg ttttattatt tgggtgagta ctttttcact cgttaaaaaag 6180
ttttgagaat attttatatt tttgttcatt taatctctcc tgaagtgatt acatctataa 6240
acaaatacag aagttaaacg atttgtttgt aattttagtt atctgtttta aaagtcataa 6300
gattagtcac tggtaggaat taatctaacg tatttattta tctgcgtaat cactgttttt 6360
agtctgtttc aaaacagtag atgttttatc tacattacgc atttgggaata ccaacatgac 6420
gaatccctcc ttcttaaaaa caaattttta gcatctaatt taacttcaat tcctattata 6480
caaaatttta agataatgca ctatcaacac actcttaagt ttgcttctaa gtcttatttc 6540
cataacttct tttacgtttc cgccattctt tgctgtttcg atttttatga tatggtgcaa 6600
gtcagcacga acacgaaccg tcttatctcc cattatatct ttttttgac tgattgggtgt 6660
atcatttcgt ttttcttttt gtgcgcctaa atttcccaca atcactcact tctttctatt 6720
tcttcttatt cttattttat catcaacaat cacaaatcac ttgtgatttg tgataagtga 6780
tttgtgataa gtgatttggt attaatatat aaaagccctc tttaaagggc ttttatgttt 6840
attttgagaa agtatataaa tcaatatatc ccttttcccc aatttttaca acggcattgt 6900
agggcttttt ctctttgttt ttgatttcct tttaccaggg tttcttttcc ctccagtaat 6960
tcttttacat tctgttttgt tagttttttc tttctaaaat gttcagctaa agtaaaactta 7020
cattcaggat aatttgaaca accataaaac gattttttta atacaatatt gttgccacac 7080
ttaggacatt ttctacaat acttttttct gcttcttttt ctttctgttc ctggtaatca 7140
gaaaaattta gtttttctat atcgttaggt acagcttcca gtaaatgaac aatgaatttt 7200
ttgatattcg taataaagtt ctcttgattg ctttctcttt taccgatttt ttttaaatac 7260
gtttcccat tagccgtcat ttcagcactc gttaaaaggt gctgactttc aactgcctgg 7320
cacataaatt ttcttttttc agttacaaca gcctgcagggt cgacgataac cacataacag 7380
tcataaaacc actccttttt aacaaacttt atcacaagaa atattttggc attctacgac 7440
tataacttaa atttatattt tttactttat aatatataat tgattataga ataatgttgc 7500
tcatatcgtt tgccaacatc tagtactcaa attacactat gttacacttg gtaattataa 7560
ccgaacttcc cctgtccaaa ttagataaga ggtaataata aatggaaaat aattttatag 7620
taaatgaaaa tgagaagcgt gtattaaaac aaattttcaa taacagcaat atttcacgaa 7680
cacaaatac gaagaattta gaacttaata aagctactat ttctaacatt ctgaacaact 7740
taaaacacaa gagtttagtt aatgaagtag gagaaggtaa tagtactaaa agtgggtggac 7800
gaaagcctat tttactcgaa attaaccaaa aatatggcta ctatatttct atggatttaa 7860
catatgatc cgttgaatta atgtacaact actttgatgc tactatatta aagcaagatt 7920
cctacgaatt aaatgataaa aatgtaagca gtatattaca aattttaaaa tctaataata 7980
acgtctcaga aaaatgatgat acgttatatg ggttacttgg tatatctata tccatacacg 8040
gtatcgttga cgatgagcaa aacataatca atcttctctt tcataaaaaat gagaaacgca 8100
catttaccga tgaattaaag tcattcacaa atgttctgt cgttatagaa aatgaagcaa 8160
atttatcagc gctatagaa aaaagtttat attttaattc aaacataaat aatttgatta 8220
ctttaagtat tcacaagggt ataggcgtg gcatcataat aaataaaaaa ctttatcgtg 8280
gctcaaatgg agaggctgga gagataggta agacattgggt tttggaatct ataaataaca 8340
atgacaacaa atattataaa atcgaagata tatgtctcca agacgcttta atacagaaaa 8400
taaataatag gttgggcgtc acattgacgt ttacagaact aatccaatat tacaacgaag 8460
gaaattcaat tgttgctcat gaaattaaac aattttattaa taaaatgaca gttctgattc 8520
ataatttgaa tacacaattt aaccagacg ctattttatat taactgtcct ttaattaatg 8580
aattaccaaa tattttaaat gaaattaaag agcaattctc ctgtttttct caaggcagtc 8640
caattcaatt acatttaact actaatgtaa aacaagctac tttattgggt ggcactttag 8700
caataatgca aaaaacatta aatataaata acattcaaat gaatattaaa taattacagc 8760
agtctgagtt ataaaataga tatctcggac cgaaacagcc gaataaggct gtttttcttt 8820
tttttatgtt ttagaataag tggctagtt tattcttgac aaaaaataat attttgatat 8880
aattaaatag ttgtcgtttg agacgactga ctttcttatt attcatctaa aatattattt 8940
tgaaaagata acacagttat tcttgacaaa aaaatataaa agtgtataat agaaaagtac 9000
tcgagagtta gtttgtttat taaattaaac aactggacga attcccgggg atccgtcgac 9060
ctgcagccaa gcttgggctt ttcagcctga tacagattaa atcagaacgc agaagcggtc 9120
tgataaaaca gaatttgctt ggcggcagta gcgcggtggt cccacctgac cccatgccga 9180
actcagaagt gaaacgccgt agcgccgatg gtagtgtggg gtctcccat gcgagagtag 9240
ggaactgcc aagcatcaaat aaaacgaaag gctcagtcga aagactgggc ctttcgtttt 9300
atctgttgtt tgtcggtgaa cgctctcctg agtaggacaa atccgcccgg agcggatttg 9360
aacgttgca agcaacggcc cggagggtgg cgggcaggac gcccgccata aactgccagg 9420
catcaaatta agcagaaggc catcctgacg gatggccttt ttgcgtttct acaaactctt 9480
tttgtttatt tttctaaata cattcaaata tgtatccgct gatcccatc ctatcgatcc 9540

```

ccacccgatg cccttgagag ccttca

9566

<210> 51
 <211> 9412
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> pEPEF20

<400> 51
 acccgatgcc cttgagagcc ttcaaccag tcaagctcctt ccggtgggag cggggcatga 60
 ctatcgctgc cgcacttatg actgtcttct ttatcatgca actcgttaga caggtgccgg 120
 cagcgctctg ggtcattttc ggcgaggacc gctttcgctg gagcgcgacg atgatcggcc 180
 tgtcgcttgc ggtattcgga atcttgacg ccctcgctca agccttcgctc actgggtccc 240
 ccaccaaagc tttcgcgagc aagcaggcca ttatcgccgg catggcgggc gacgcgctgg 300
 gctacgtctt gctggcgctc gcgacgcgag gctggatggc cttccccatt atgattcttc 360
 tcgcttccgg cggcatcggg atgcccgctg tgcaggccat gctgtccagg caggtagatg 420
 acgaccatca gggacagctt caaggatcgc tcgcgctctt taccagccta acttcgatca 480
 ctggaccgct gatcgtaacg gcgatttatg ccgcctcggc gagcacatgg aacgggttgg 540
 catggattgt aggcgcggcc ctataccttg tctgcctccc cgcgttgctg cgcggtgcat 600
 ggagccgggc cactcgacc tgaatggaag ccgcgggcac ctcgctaagc gattcaccac 660
 tccaagaatt ggagccaatc aattcttgcg gagaactgtg aatgcgcaaa ccaacccttg 720
 gcagaacata tccatcgctg ccgcatctc cagcagccgc acgcggcgca tctcgggcag 780
 cgttgggtcc tggccacggg tgcgatgat cgtgctcctg tcgttgagga ccgggctagg 840
 ctggcggggt tgccttactg gttagcagaa tgaatcacg atacgcgagc gaacgtgaag 900
 cgactgctgc tgcaaaacgt ctgcgacctg agcaacaaca tgaatggtct tcggtttccg 960
 tgtttcgtaa agtctgaaa cgcggaagtc ccctacgtgc tgctgaagtt gcccgcaaca 1020
 gagagtggaa ccaaccggtg ataccacgat actatgactg agagtcaacg ccatgagcgg 1080
 cctcatttct tattctgagt tacaacagtc cgcaccgctg tccggtagct ctttcgggtg 1140
 ggcgcggggc atgactatcg tcgcccact tatgactgtc ttctttatca tgcaactcgt 1200
 aggcagagtg ccggcagcgc ccaacagtc ccggccacg gggcctgcca ccataccac 1260
 gccgaacaa gcgcccgtga ccattatgtt ccggtctgac atcgaggat gctgctggct 1320
 accctgtgga acacctacat ctgtattaac gaagcgctaa ccgtttttat caggctctgg 1380
 gaggcagaat aaatgatcat atcgtaatt attacctca cggggagagc ctgagcaaac 1440
 tggcctcagg catttgagaa gcacacggtc acactgcttc cggtagtcaa taaaccggta 1500
 aaccagcaat agacataagc ggctatttaa cgaccctgcc ctgaaccgac gaccgggtcg 1560
 aatttgcttt cgaatttctg ccattcatcc gcttattatc acttattcag gcgtagcacc 1620
 agcggtttat gggcaccaat aactgcctta aaaaaattac gccccgccct gccactcatc 1680
 gcagtactgt tgtaattcat taagcattct gccgacatgg aagccatcac agacggcatg 1740
 atgaacctga atcgccagcg gcacacgac cttgtcgctt tgcgtataat atttgcccat 1800
 ggtgaaaacg ggggcgaaga agttgtccat attggccacg tttaaatcaa aactggtgaa 1860
 actacccag ggattggctg agacgaaaaa catattctca ataaaccctt tagggaaata 1920
 ggccagggtt tcaccgtaac acgccacatc ttgcgaatat atgtgtagaa actgccgga 1980
 atcgtcgtgg tattcactcc agagcgatga aaacgtttca gtttgctcat ggaaaacggt 2040
 gtaacaaggg tgaacactat cccatatcac cagctcaccg tctttcattg ccatacgaga 2100
 ggtttgatga ctttgacctt tggttttaaa tttttgaaaa aaataaaaaa aggcgaagcc 2160
 tattatata tttatcata ttttttaac ttttgttctt ttgctgaaa aaaaaggcag 2220
 tgttttcgct agttatagaa attaaacagt cacaaaaatc gatgtataca gtcacaaaaa 2280
 tcgatttttg tgactctatg catcgatgta tacgagtcac aaaaatcgat gtacacagta 2340
 cgacttttgt atttgtgtac tgtatatagt ataataaaag catagagaaa actcactatg 2400
 aaatgacttt ctctatgcta ctactaaaac acgcaaagga gcgtatttat actatgatta 2460
 taccagaaaa acaaaacaaa caaaagcagg tgctgacctt gaatgaatta gaaaaacgca 2520
 aagtagtgga gataacgct ttaattcaaa gtgtggctaa aatgcaaaaa acagctttta 2580
 aaatgtttga cttagccgtg tcttgtattg ataccgaaga acccccaaaa aataatactg 2640
 tttatctttt aaaatctgaa ctttttaaat tttttgaagt atcgtctagc agtaagcata 2700
 gtcaattcaa agaagcagtt aattatatgc aaaagcaagc ctttttcaat attaaagctg 2760
 ataagaaact aggtattgag tatgaaagca ttgtgcctat tccttatgta aaatggaacg 2820
 attataacga tgaagtaaca atacgttttg accaagccat tatgccctat ctgattgatt 2880

taaaggcaga	atttactcaa	tataaaat	ccgaactaca	aaaactcaat	agcaagtatt	2940
cgattat	gtatcgttg	ttatccatga	attataacca	atacgaacat	tacagcgtga	3000
aagggggacg	gagagttgaa	caagtggaa	cctaccgcaa	ccctcaata	aaagtgaag	3060
aattgcgaga	aataactgat	acaataaatg	aacatcaaca	ttttcccat	tttgaaacta	3120
gagtattaaa	aaaagcaatt	gaagaaatca	acgctcacac	ctctttta	gtaacctatg	3180
agaaaaagaa	agcagggcgg	agcattgact	ctattgtctt	tcatattgag	aagaaacgca	3240
tggcagacga	taacagttac	aagttgggag	ataaagacta	tcaagacgat	aaaaagcaaa	3300
aatcaagaaa	tgaagctgac	ttactaaaac	aggcaatgga	aagtaaata	acaagacttt	3360
tatctgaaaa	tttcttaatt	ggcatgaatg	atattatgga	tacagccaca	atggtaggct	3420
tacagaagaa	tgtttatccg	ctttatgacg	aactgaaaga	attaagaggg	ctaaatgggtg	3480
tcaaagacca	cttgctttat	gttgccagca	aaaaagaagc	ctattctaaa	cgtaatgtag	3540
cgaatatct	taaaaaagcc	attgaacaat	atctgccaac	ggttaaaagg	caggacttaa	3600
accatgagtg	aagacttaaa	aacgataaaa	gagttggcgg	acgagctagg	actagctaaa	3660
cagaccgttc	aataccattt	aaagtttttg	ccgaccgaaa	atcgacaaa	aaacaatagt	3720
ggttcgatcg	tctaaccct	tgaagaacaa	aggtttatca	aaaaaagact	aacaaaaata	3780
aaaaaagttc	aaccgaccga	aaatcgacaa	gaaaaaatga	agaattaagc	gttaatagtc	3840
ttttgttagg	cggtatgagca	ttcatcaggc	gggcaagaat	gtgaataaag	gccggataaa	3900
acttgtgctt	atttttcttt	acggtcttta	aaaaggccgt	aatatccagc	tgaacggctc	3960
ggttataggf	acattgagca	actgactgaa	atgcctcaaa	atgttcttta	cgatgccatt	4020
gggatataatc	aacggtggta	tatccagtga	ttttttctc	catttttagct	tccttagctc	4080
ctgaaaatct	cgataactca	aaaaatacgc	ccggtagtga	tcttatttca	ttatggtgaa	4140
agttggaacc	tcttactgtc	cgatcaacgt	ctcattttcg	ccaaaagtgg	gcccagggtc	4200
tcccggtatc	aacagggaca	ccaggattta	tttattctgc	gaagtatct	tcggtcacag	4260
gtatttatc	ggcgcaaaagt	gcgtcgggtg	atgctgcaa	cttactgatt	tagtgtatga	4320
tggtgtttt	gaggtgctcc	agtggtctct	gtttctatca	gctgtccctc	ctgttcagct	4380
actgacgggg	tggtgcgtaa	cggcaaaagc	accgccggac	atcagcgcta	gcggagtgtg	4440
tactggctta	ctatgttggc	actgatgagg	gtgtcagtga	agtgttcat	gtggcaggag	4500
aaaaaaggct	gcaccgggtg	gtcagcagaa	tatgtgatac	aggatatatt	ccgcttctc	4560
gctcactgac	tgcctacgct	cggtcgttcg	actgcggcga	gcggaatgg	cttacgaacg	4620
gggcggagat	ttcctggaag	atgccaggaa	gatacttaac	agggaagtga	gagggccgct	4680
gcaaagccgt	ttttccatag	gtccgcgcc	cctgacaagc	atcacgaaat	ctgacgctca	4740
aatcagtggt	ggcgaaaccc	gacaggacta	taaagatacc	aggcgtttcc	ccctggcggc	4800
tcctcgtgc	gctctcctgt	tcctgccttt	cggtttaccg	gtgtcatctc	gctgttatgg	4860
ccgcgtttgt	ctcattccac	gcctgacact	cagttccggg	taggcagttc	gctccaagct	4920
ggactgtatg	cacgaacccc	ccgttcagtc	cgaccgctgc	gccttatccg	gtaactatcg	4980
tcttgagtc	aacccgaaa	gacatgcaaa	agcaccactg	gcagcagcca	ctggtaattg	5040
atttagagga	gttagtcttg	aagtcatgcg	ccggttaagg	ctaaactgaa	aggacaagtt	5100
ttggtgactg	cgctccctcca	agccagttac	cctggttcaa	agagttggtg	gctcagagaa	5160
ccttcgaaaa	accgccctgc	aaggcggttt	tttcgttttc	agagcaagag	attacgcgca	5220
gaccaaaacg	atctcaagaa	gatcatctta	ttaatcagat	aaaatatctc	tagatttcag	5280
tgcaatttat	ctcttcaaat	gtagcacctg	aagtcagccc	catacgatat	aagttgtaat	5340
tctcatgttt	gacagcttat	catcgataca	aattccccgt	aggcgctagg	gacctcttta	5400
gcttcttgga	agctgtcagt	agtatatcta	ataatttatc	tccattccct	ttagtaacgt	5460
gtaactttcc	aaatttaaaa	aagcgactca	tagaattatt	tcctcccggt	aaataataga	5520
taactattaa	aaatagacaa	tacttgctca	taagtaatgg	tacttaaat	gtttactttg	5580
gcgtgtttca	ttgcttgatg	aaactgattt	ttagtaaaaca	gttgacgata	ttctcgattg	5640
acccattttg	aaacaaagta	cgtatatagc	ttccaatatt	tatctggaac	atctgtggta	5700
tggcgggtaa	gttttatata	gacactgttt	acttttggtt	taggatgaaa	gcattccgct	5760
ggcagcttaa	gcaattgctg	aatcgagact	tgagtgtgca	agagcaaccc	tagtgttcgg	5820
tgaatatcca	aggtacgctt	gtagaatcct	tcttcaacaa	tcagatagat	gtcagacgca	5880
cggctttcaa	aaaccacttt	tttaataaatt	tgtgtgctta	aatggtaagg	aatattccca	5940
acaattttat	acctctgttt	gttagggaat	tgaactgtga	gaatatcttg	gtgaattaaa	6000
gtgacacgag	tttcaagttt	taatttttct	gacgataagt	tgaatagatg	actgtctaat	6060
tcaatagacg	ttacctgttt	acttatttta	gccagtttctg	tcgttaaatg	ccctttacct	6120
gttccaattt	cgtaaacggg	atcggtttct	tttaaatcca	attgttttat	tatttggttg	6180
agtacttttt	cactcgttaa	aaagttttga	gaatatttta	tatttttggt	catgtaatct	6240
ctcctgaagt	gattacatct	ataaacaaat	acagaagtta	aacgatttgt	ttgtaatttt	6300
agttatctgt	ttaaaaagtc	ataagattag	tcactggtag	gaattaatct	aacgtattta	6360
tttatctgcg	taatcactgt	ttttagtctg	tttcaaaaca	gtagatgttt	tatctacatt	6420
acgcatttgg	aataccaaca	tgacgaatcc	ctccttctta	attacaaatt	tttagcatct	6480
aatttaactt	caattcctat	tatacaaaat	tttaagataa	tgcactatca	acacactctt	6540

```

aagtttgctt ctaagtctta tttccataac ttcttttacg tttccgcat tctttgctgt 6600
ttcgattttt atgatatggg gcaagtcagc acgaacacga accgtcttat ctcccattat 6660
atcttttttt gcaactgattg gtgtatcatt tcgtttttct tttgtgctgc cttaaatttcc 6720
cacaatcact cacttctttc tatttcttct tatttcttatt ttatcatcaa caatcacaaa 6780
tcactttgtga tttgtgataa gtgattttgtg ataatgtgatt tgtgattaat atataaaagc 6840
cctcttttaa gggtttttat gtttattttg agaaagatat aaaatcaata tatccctttt 6900
ccccaatttt tacaacggca ttgtagggtt ttttctcttt gtttttgatt tcctttttacc 6960
agggttttct ttcctccag taattctttt acattcggtt tggtagttt tttctttcta 7020
aaatgttcag ctaaagtaaa cttacattca ggataatttg aacaaccata aaacgatttt 7080
tttaatacaa tattgttgcc acacttagga cattttccta caatactttt ttctgcttct 7140
tttcttttct gttcctggta atcagaaaaa tttagttttt ctatatcggt aggtacagct 7200
tccagtaaat gaacaatgaa ttttttgata ttcgtaataa agttctcttg attgccttct 7260
cttttaccga ttttttttaa atacgttttc catttagccg tcatttcagc actcggttaa 7320
aggtgctgac tttcaactgc ctggcacaaat aattttcctt tttcagttac aacagcctgc 7380
aggtcgacga taaccacata acagtcataa aaccactcct ttttaacaaa ctttatcaca 7440
agaaatattt tggcattcta cgactataac ttaaatttat attttttact ttataatata 7500
taattgatta tagaataatg ttgctcatat cgtttgccaa catctagtag tcaaattaca 7560
ctatgttaca cttggtaata ttaaccgaac ttcccctgtc caaattagat aagaggtaat 7620
aataaatgga aataaatttt atagttaagt aaaatgagaa gcgtgtatta aaacaaattt 7680
tcaataacag caatatattca cgaacacaaa tatcgaagaa tttagaactt aataaagcta 7740
ctatttctaa cattctgaac aacttaaaac acaagagttt agttaatgaa gtaggagaag 7800
gtaaatgtac taaaagtggg ggacgaaagc ctattttact cgaaattaac caaaaatatg 7860
gctactatat ttctatggat ttaacatatg attccgttga attaatgtac aactactttg 7920
atgctactat attaaagcaa gattcctacg aattaaatga taaaaatgta agcagtatat 7980
tacaattttt aaaatctaata ataaacgtct cagaaaaata tgatacgtta tatgggttac 8040
ttggtatata tatatccata cacggtatcg ttgacgatga gcaaacata atcaatcttc 8100
cttttcataa aaatgagaaa cgcacattta ccgatgaatt aaagtcattc acaaatgttc 8160
ctgtcgttat agaaaaatgaa gcaaatttat cagcgtctata tgaaaaaagt ttatatatta 8220
attcaaacat aaataatttg attactttaa gtattcaciaa gggatataggc gctggcatca 8280
taataaataa aaaactttat cgtggctcaa atggagaggc tggagagata ggtaagacat 8340
tgggttttga atctataaat aacaatgaca acaaatatta taaaatcgaa gatatatgct 8400
cccaagacgc tttataacag aaaataaata ataggttggg cgtcacattg acgtttacag 8460
aactaatcca atattacaac gaaggaaatt caattgttgc tcatgaaatt aaacaattta 8520
ttaataaaat gacagttctg attcataatt tgaatacaca atttaacca gacgtatttt 8580
atattaactg tcttttaatt aatgaattac caaatatttt aaatgaaatt aaagagcaat 8640
tctctgtttt ttctcaaggc agtccaattc aattacattt aactactaat gtaaaacaag 8700
ctactttatt ggggtggcact ttagcaataa tgcaaaaaac attaaatata aataacattc 8760
aaatgaatat taaataatta cagcagctcg agttataaaa tagatatctc ggaccgtttc 8820
gtgaactttt ttgttgacaa agataaaaaac acatgatata cttaaatcac tcgagagtta 8880
gtttgtttat taaattaacc aactggacga attcccgggg atccgtcgac ctgcagccaa 8940
gcttgggctt ttacgcctga tacagattaa atcagaacgc agaagcggtc tgataaaaca 9000
gaatttgcct ggcggcagta gcgcgtggg cccactgac cccatgccga actcagaagt 9060
gaaacgccgt agcgcgcatg gtatgtgtgg gtctcccat gcgagagtag ggaactgcca 9120
ggcatcaaat aaaacgaaag gctcagtcga aagactgggc ctttcgtttt atctgttgtt 9180
tgtcggtgaa cgctctctcg agtaggacaa atccgcggg agcggatttg aacgttgcga 9240
agcaacggcc cggagggttg cgggcaggac gcccgccata aactgccagg catcaaatta 9300
agcagaaggc catcctgacg gatggccttt ttgcgtttct acaaactctt tttgtttatt 9360
tttctaaata cattcaataa tgtatccgct gatcccatc ctatcgatcc cc 9412

```

<210> 52

<211> 9412

<212> DNA

<213> Artificial Sequence

<220>

<223> pEPEF22

<400> 52

accgatgcc cttgagagcc ttcaaccagg tcagctcctt ccggtgggag cggggcatga 60

ctatcgctgc	cgcacttatg	actgtcttct	ttatcatgca	actcgtagga	caggtgccgg	120
cagcgctctg	ggcatttttc	ggcgaggacc	gctttcgctg	gagcgcgacg	atgatcggcc	180
tgctgcttgc	ggtattcgga	atcttgacag	ccctcgctca	agccttcgtc	actggtcccc	240
ccaccaaacg	tttcggcgag	aagcaggcca	ttatcgccgg	catggcgggc	gacgcgctgg	300
gctacgtctt	gctggcggtt	gcgacgcgag	gctggatggc	cttccccatt	atgattcttc	360
tcgcttcggg	cggcatcggg	atgcccgcgt	tgcaggccat	gctgtccagg	caggtagatg	420
acgaccatca	gggacagctt	caaggatcgc	tcgcggctct	taccagccta	acttcgatca	480
ctggaccgct	gatcgtcacg	gcgatttatg	ccgcctcggc	gagcacatgg	aacgggttgg	540
catggattgt	agggcccgcc	ctataccttg	tctgcctccc	cgcgttgctg	cgcggtgcat	600
ggagccgggc	cacctcgacc	tgaatggaag	ccggcgccac	ctcgctaacg	gattcaccac	660
tccaagaatt	ggagccaatc	aattcttgcg	gagaactgtg	aatgcgcaaa	ccaacccttg	720
gcagaacata	tccatcgctg	ccgcctatct	cagcagccgc	acgcggcgca	tctcgggcag	780
cgttggggtc	tggccacggg	tgcgcgatg	cgtgctcctg	tcgttgagga	cccggctagg	840
ctggcggggt	tgcttactg	gttagcagaa	tgaatcaccg	atacgcgagc	gaacgtgaag	900
cgactgctgc	tgcaaaacgt	ctgcgacctg	agcaacaaca	tgaatggctc	tcggtttccg	960
tgtttcgtaa	agtcctgaaa	cgcggaagtc	ccctacgtgc	tgctgaagtt	gcccgcacaa	1020
gagagtggaa	ccaaccgggtg	ataccacgat	actatgactg	agagtcaacg	ccatgagcgg	1080
cctcatttct	tattctgagt	tacaacagtc	cgcaccgctg	tccggtagct	ccttcgggtg	1140
ggcgcggggc	atgactatcg	tcgcccact	tatgactgtc	ttctttatca	tgcaactcgt	1200
aggacagggtg	ccggcagcgc	ccaacagtc	cccgccacg	gggctgcca	ccatacccac	1260
gccgaacaa	gcgccttgc	ccattatgtt	ccggatctgc	atcgcaggat	gctgctggct	1320
accctgtgga	acacctacat	ctgtattaac	gaagcgctaa	ccgtttttat	caggctctgg	1380
gaggcagaat	aaatcgatcat	atcgctcaatt	attgacctca	cggggagagc	ctgagcaaac	1440
tggcctcagg	catttgagaa	gcacacggtc	acactgcttc	cggtagtcaa	taaaccggta	1500
aaccagcaat	agacataagc	ggctatttaa	cgacctgcc	ctgaaccgac	gaccgggtcg	1560
aatttgcttt	cgaatttctg	ccattcatcc	gcttattatc	acttatcag	gcgtagcacc	1620
aggcgtttaa	gggcaccaat	aactgcctta	aaaaaattac	gccccccct	gccactcatc	1680
gcagtactgt	tgtaattcat	taagcattct	gccgacatgg	aagccatcac	agacggcatg	1740
atgaacctga	atcgccagcg	gcacagcac	cttgtcgctt	tgctataaat	atttgcccat	1800
ggtgaaaacg	ggggcgaaag	agttgtccat	attggccacg	tttaaatcaa	aactggtgaa	1860
actcaccag	tgattggctg	agacgaaaaa	catatttctc	ataaacctt	tagggaaata	1920
ggccagggtt	tcaccgtaac	acgccacatc	ttgcgaatat	atgtgtagaa	actgccggaa	1980
atcgctgctg	tattcactcc	agagcgatga	aaacgtttca	gtttgctcat	ggaaaacggt	2040
gtaacaaggg	tgaacactat	cccatacac	cagctcaccg	tctttcattg	ccatacgaga	2100
ggtttgatga	ctttgacctt	tggttttaaa	tttttgaaaa	aaataaaaaa	aggcgaaagg	2160
tattatata	ttatcatata	tattttaatc	ttttgttctt	ttgcgtgaaa	aaaaaggcag	2220
tgtttctgct	agttatagaa	attaaacagt	cacaaaaatc	gatgtataca	gtcacaaaaa	2280
tcgatttttg	tgactctatg	catcgatgta	tacgagtcac	aaaaatcgat	gtacacagta	2340
cgacttttgt	atttgtgtac	tgtatatagt	ataataaaa	catagagaaa	actcactatg	2400
aaatgacttt	ctctatgcta	ctactaaaac	acgcaaagga	gcgtatttat	actatgatta	2460
taccagaaaa	acaaaaacaa	caaaagcagg	tgctgacctt	gaatgaatta	gaaaaacgca	2520
aagtagtgga	gcataacgct	ttaattcaaa	gtgtggctaa	aatgcaaaaa	acagctttaa	2580
aaatgtttga	gttagccgtg	tcttgtattg	ataccgaaga	acccccaaaa	aataatactg	2640
tttatctttt	aaaatctgaa	ctttttaaat	tttttgaagt	atcgtctagc	agtaagcata	2700
gtcaattcaa	agaagcagtt	aatttatatg	aaaagcaagc	ctttttcaat	attaaagctg	2760
ataagaaact	aggtattgag	tatgaaaagca	ttgtgcctat	tccttatgta	aaatggaaag	2820
attataacga	tgaagtaaca	atacgttttg	accaagccat	tatgccctat	ctgattgatt	2880
taaaggcaga	atttactcaa	tataaaat	ccgaactaca	aaaactcaat	agcaagtatt	2940
cgattatttt	gtatcgttgg	ttatccatga	attataacca	atacgaaat	tacagcgtga	3000
aagggggacg	gagagtggaa	caagtggaa	cctaccgcaa	cccccaata	aaagtgaag	3060
aattgcgaga	aataactgat	acaataaatg	aacatcaaca	ttttcccat	tttgaaacta	3120
gagtatttaa	aaaagcaatt	gaagaaatca	acgtcacac	ctctttta	gtaacctatg	3180
agaaaaagaa	agcaggcg	agcattgact	ctattgtctt	tcataatgag	aagaaacgca	3240
tggcagacga	taacagttac	aagttgggag	ataaagacta	tcaagacgat	aaaaagcaaa	3300
aatcaagaaa	tgaagctgac	ttactaaaac	aggcaatgga	aagtaaatat	acaagacttt	3360
tatctgaaaa	tttcttaatt	ggcatgaatg	atattatgga	tacagccaca	atggtaggct	3420
tacagaagaa	tgtttatccg	ctttatgacg	aactgaaaga	attaagaggg	ctaaatggtg	3480
tcaaagacca	cttgccttat	gttgccagca	aaaaagaagc	ctattctaaa	cgtaatgtag	3540
cgaaatatct	taaaaaagcc	attgaacaat	atctgccaac	ggttaaaaag	caggacttaa	3600
accatgagtg	aagacttaaa	aacgataaaa	gagttggcgg	acgagctagg	actagctaaa	3660
cagaccgttc	aataccattt	aaagtttttg	ccgaccgaaa	atcgacaaaa	aaacaatagt	3720

```

ggttcgatcg tcctaaacct tgaagaacaa aagtttatca aaaaaagact aacaaaaata 3780
aaaaaagttc aaccgaccga aaatcgacaa gaaaaaatga agaattaagc gttaatagtc 3840
ttttgttagg cggtatgagca ttcatcaggc gggcaagaat gtgaataaag gccggataaa 3900
acttgtgctt atttttcttt acggtcttta aaaaggccgt aatatccagc tgaacggctc 3960
ggttataggt acattgagca actgactgaa atgcctcaaa atgttcttta cgatgccatt 4020
gggatataac aacggtggta tatccagtga ttttttctc catttttagc tccttagctc 4080
ctgaaaatct cgataactca aaaaatacgc ccggtagtga tcttatttca ttatggtgaa 4140
agttggaacc tcttacgtgc cgatcaacgt ctcatcttcg ccaaaagttg gccagggct 4200
tcccgtatc aacagggaca ccaggattta tttattctgc gaagtatct tccgtcacag 4260
gtatttattc ggcgcaaaagt gcgtcgggtg atgctgccaa ctactgatt tagtgtatga 4320
tgggtgtttt gaggtgctcc agtggcttct gtttctatca gctgtccctc ctgttcagct 4380
actgacgggg tgggtgcgtaa cggcaaaagc accgccggac atcagcgcta gcggagtga 4440
tactggctta ctatgttgc actgatgagg gtgtcagtga agtgcttcac gtggcaggag 4500
aaaaaaggct gcaccggtgc gtcagcagaa tatgtgatac aggatatatt ccgcttctc 4560
gctcactgac tcgctacgct cggctgttcg actgcggcga gcggaatgg cttacgaacg 4620
gggcggagat ttcttggaag atgccaggaa gatacttaac agggaaagtga gagggccgcg 4680
gcaaagccgt ttttccatag gtcgccccc cctgacaagc atcacgaaat ctgacgctca 4740
aatcagtggt ggcgaaaccc gacaggacta taaagatacc aggcgtttcc ccctggcggc 4800
tccctcgctc gctctctgt tctgccttt cggtttaccg gtgtcattcc gctgttatgg 4860
ccgcgtttgt ctcatccac gcctgacact cagttccggg taggcagttc gctccaagct 4920
ggcatgtatg cacgaacccc ccgttcagtc cgaccgctgc gccttatccg gtaactatcg 4980
tcttgagtc aaccgggaaa gacatgcaaa agcaccactg gcagcagcca ctggtaattg 5040
atthagagga gttagtcttg aagtcatgcg ccggttaagg ctaaaactgaa aggacaagtt 5100
ttgggtgactg cgctcctcca agccagttac ctcggttcaa agagttggtg gctcagagaa 5160
ccttcgaaaa accgccctgc aaggcggttt tttcgtttc agagcaagag attacgcgca 5220
gaccaaaacg atctcaagaa gatcatctta ttaatcagat aaaatatctc tagatttcag 5280
tgcaatttat ctcttcaaat gtagcacctg aagtcagccc catacgatac aagttgtaat 5340
tctcatgttt gacagcttat catcgataca aattcccgt aggcgctagg gacctctta 5400
gcttcttggg agctgtcagt agtatactta ataatttatc tccattccct ttagtaacgt 5460
gtaactttcc aaatttaaaa aagcgactca tagaattatt tctcccgtt aaataataga 5520
taactattaa aaatagacaa tacttgctca taagtaatgg tacttaaat gtttactttg 5580
gcgtgtttca ttgcttgatg aaactgattt ttagtaaaaca gttgacgata ttctcgattg 5640
acccattttg aaacaaagta cgtatatagc ttccaatatt tatctggaac atctgtggtg 5700
tggcgggtaa gttttattaa gacactgttt acttttggtt taggatgaaa gcattccgct 5760
ggcagcttaa gcaactgtg aatcgagact tgagtgtgca agagcaaccc tagtgttcgg 5820
tgaatatcca aggtacgctt gtagaatcct tcttcaacaa tcagatagat gtcagacgca 5880
cggctttcaa aaaccacttt ttttaataatt tgtgtgctta aatggtaagg aatattccca 5940
acaattttat acctctgttt gttagggaat tgaaactgta gaatatcttg gtgaattaaa 6000
gtgacacgag tattcagttt taatttttct gacgataagt tgaatagatg actgtctaatt 6060
tcaatagacg ttacctgttt acttattttta gccagtttcg tcgttaaatg ccctttacct 6120
gttccaattt cgtaaacgggt atcggtttct tttaaattca attgttttat tatttggttg 6180
agtactttt cactcgttaa aaagttttga gaattttta tatttttgtt catgtaattc 6240
ctcctgaagt gattacatct ataacaacaa acagaagtta aacgatttgt ttgtaatttt 6300
agttatctgt ttaaaaagtc ataagattag tcaactgtag gaattaatct aacgtattta 6360
tttatctgcg taatcactgt ttttagtctg tttcaaaaca gtagatgttt tatctacatt 6420
acgcatttgg aataccaaca tgacgaatcc ctcttctta attacaaatt tttagcatct 6480
aatttaactt caattcctat tatacaaaat tttagataa tgcactatca acacactctt 6540
aagtttgctt ctaagtctta tttccataac tttttttacg tttccgccat tctttgctgt 6600
ttcgattttt atgatatggt gcaagtcagc acgaacacga accgtcttat ctccattat 6660
atctttttt gcactgattg gtgtatcatt tegttttct tttgtgctc cttaaatttc 6720
cacaatcact cacttcttct tatttcttct tattcttatt ttatcatcaa caatcacaaa 6780
tcaactgtga tttgtgataa gtgatttgtg ataagtgatt tgtgattaat atataaaagc 6840
cctctttaaa gggcttttat gtttattttg agaaagatat aaaatcaata tatccctttt 6900
ccccaatttt tacaacggca ttgtagggtt ttttctctt gtttttgatt tctttttacc 6960
agggtttctt ttccctccag taattctttt acattcgttt tggttagttt tttctttcta 7020
aaatgttcag ctaaagtaaa cttacattca ggataatttg aacaaccata aaacgatttt 7080
tttaatacaa tattgttgcc acatttagga cattttccta caatactttt ttctgcttct 7140
tttctttct gttcctggta atcagaaaaa ttttagtttt ctatatcgtt aggtacagct 7200
tccagtaaat gaacaatgaa ttttttgata ttcgtaataa agttctcttg attgccttct 7260
cttttaccga ttttttttaa atacgtttcc catttagccg tcaatttcagc actcgttaaa 7320
aggtgctgac tttcaactgc ctggcacaat aattttcctt tttcagttac aacagcctgc 7380

```

```

aggtcgacga taaccacata acagtcataa aaccactcct ttttaacaaa ctttatcaca 7440
agaaatattt tggcattcta cgactataac ttaaatttat attttttact ttataatata 7500
taattgatta tagaataatg ttgctcatat cgtttgccaa catctagtagt tcaaattaca 7560
ctatgttaca cttggtaata ttaaccgaac ttcccctgtc caaattagat aagaggtaat 7620
aataaatgga aaataatttt atagtaaatg aaaatgagaa gcgtgtatta aaacaaattt 7680
tcaataacag caatatttca cgaacacaaa tatcgaagaa tttagaactt aataaaagcta 7740
ctatttctaa cattctgaac aacttaaaac acaagagttt agttaatgaa gtaggagaag 7800
gtaaatgtac taaaagtggg ggacgaaagc ctattttact cgaaattaac caaaaatatg 7860
gctactatat ttctatggat ttaacatattg attccgttga attaatgtac aactactttg 7920
atgctactat attaaagcaa gattcctacg aattaaatga taaaaatgta agcagtatat 7980
tacaaatttt aaaatctaata ataaacgtct cagaaaaata tgatacgta tatgggttac 8040
ttggtatata tatatccata cacggtatcg ttgacgatga gcaaaacata atcaatcttc 8100
cttttcataa aaatgagaaa cgcacattta ccgatgaatt aaagtcattc acaaattgtt 8160
ctgtcggtat agaaaaatgaa gcaaatttat cagcgctata tgaaaaaagt ttatatatta 8220
attcaaacat aaataatttg attactttaa gtattcaciaa gggatataggc gctggcatca 8280
taataaataa aaaactttat cgtggctcaa atggagaggc tggagagata ggtaagacat 8340
tgggttttga atctataaat aacaatgaca acaaatatta taaaatcgaa gatatatgct 8400
cccaagacgc tttaatacag aaaataaata ataggttggg cgtcacattg acgtttacag 8460
aactaatcca atattacaac gaaggaaatt caattgttgc tcatgaaatt aaacaattta 8520
ttaataaaat gacagttctg attcataatt tgaatacaca atttaacca gacgctattt 8580
atattaactg tcctttaatt aatgaattac caaatatttt aaatgaaatt aaagagcaat 8640
tctcctgttt ttctcaaggc agtccaattc aattacattt aactactaat gtaaaacaag 8700
ctactttatt ggggtggcact tttagcaataa tgcaaaaaac attaaatata aataacattc 8760
aaatgaatat taaataatta cagcagctcg agttataaaa tagatatctc ggaccgtttc 8820
gtgaactttt ttgttgacaa agataaaaac acatgatata attaaatcac tcgagagtta 8880
gtttgtttat taaattaacc aactggacga attcccgggg atccgtcgac ctgcagccaa 8940
gcttggtgctt ttcagcctga tacagattaa atcagaacgc agaagcggtc tgataaaaaca 9000
gaatttgctt ggcggcagta gcgcggtggt cccacctgac cccatgccga actcagaagt 9060
gaaacgcctg agcgccgatg gtagtgtggg gtctcccat gcgagagtag ggaactgcca 9120
ggcatcaaat aaaacgaaag gctcagtcga aagactgggc ctttcgtttt atctgttgtt 9180
tgtcgttgaa cgctctcctg agtaggacaa atccgccggg agcggatttg aacgttgcca 9240
agcaacggcc cggagggttg cgggcaggac gcccgccata aactgccagg catcaaatta 9300
agcagaaggc catcctgacg gatggccttt ttgcgtttct acaaaactct tttgtttatt 9360
tttctaaata cattcaaata tgtatccgct gatcccatc ctatcgatcc cc 9412

```

<210> 53

<211> 6852

<212> DNA

<213> Artificial Sequence

<220>

<223> pXyl-T5-DD p15a

<400> 53

```

tggcattcta cgactataac ttaaatttat attttttact ttataatata taattgatta 60
tagaataatg ttgctcatat cgtttgccaa catctagtagt tcaaattaca ctatgttaca 120
cttggtaata ttaaccgaac ttcccctgtc caaattagat aagaggtaat aataaatgga 180
aaataatttt atagtaaatg aaaatgagaa gcgtgtatta aaacaaattt tcaataacag 240
caatatttca cgaacacaaa tatcgaagaa tttagaactt aataaaagcta ctatttctaa 300
cattctgaac aacttaaaac acaagagttt agttaatgaa gtaggagaag gtaatagtagt 360
taaaagtggg ggacgaaagc ctattttact cgaaattaac caaaaatatg gctactatat 420
ttctatggat ttaacatattg attccgttga attaatgtac aactactttg atgctactat 480
attaaagcaa gattcctacg aattaaatga taaaaatgta agcagtatat taaaaattt 540
aaaatctaata ataaacgtct cagaaaaata tgatacgta tatgggttac ttggtatata 600
tatatccata cacggtatcg ttgacgatga gcaaaacata atcaatcttc cttttcataa 660
aaatgagaaa cgcacattta ccgatgaatt aaagtcattc acaaattgtt ctgtcggtat 720
agaaaaatgaa gcaaatttat cagcgctata tgaaaaaagt ttatatatta attcaaacat 780
aaataatttg attactttaa gtattcaciaa gggatataggc gctggcatca taataaataa 840
aaaactttat cgtggctcaa atggagaggc tggagagata ggtaagacat tgggttttga 900

```

```

atctataaat aacaatgaca acaaatatta taaaatcgaa gatatatgct cccaagacgc 960
ttaataacag aaaataaata ataggttggg cgtcacattg acgtttacag aactaatcca 1020
atattacaac gaaggaaatt caattgttgc tcatgaaatt aaacaattta ttaataaaat 1080
gacagttctg attcataatt tgaatacaca atttaacca gacgtatatt atattaactg 1140
tcctttaatt aatgaattac caaatatatt aaatgaaatt aaagagcaat tctcctgttt 1200
ttctcaaggc agtccaatfc aattacattt aactactaat gtaaaacaag ctactttatt 1260
gggtggcact ttagcaataa tgcaaaaaac attaaatata aataacattc aaatgaatat 1320
taaataatta cagcagtcct agttataaaa tagatatctc ggaccgtcat aaaaaattta 1380
ttttacatca ggaaaaattt tctgtatatatt agattcaagt tagtttgttt attaaattaa 1440
ccaactaaaa tgtagaattc gagctcggta cccggggatc ctctagagtc gacctgcagc 1500
caagcttggg cttttcagcc tgatacagat taaatcagaa cgcagaagcg gtctgataaa 1560
acagaatttg cctggcgcca gtagcgcgtt ggtccacct gaccccatgc cgaactcaga 1620
agtgaacgc cgtagcgccg atggtagtgt ggggtctccc catgcgagag tagggaaactg 1680
ccaggcatca aataaaacga aaggctcagt cgaaagactg ggcttttcgt tttatctgtt 1740
gtttgtcggg gaacgctctc ctgagtagga caaatccgcc gggagcggat ttgaacgttg 1800
cgaagcaacg gcccgagggg tggcgggcag gacgcccgcc ataaactgcc aggcatacaa 1860
ttaagcagaa ggccatcctg acggatggcc tttttgcgtt tctacaaaact ctttttgttt 1920
atttttctaa atacattcaa atatgtatcc gctgatcccc atcctatcga tgataagctg 1980
tcaaacatga gaattaaatc aatctaaagt atatatgagt aaacttggtc tgacagttaac 2040
caatgcttaa tcagtggagg acctatctca cgcgtctgtc tatttcgttc atccatagtt 2100
gcctgactcc ccgtcgtgta gataactacg atacgggagg gcttaccatc tggccccagt 2160
gctgcaatga taccgcgaga cccacgctca ccggctccag atttatcagc aataaaccag 2220
ccagccgga gggccgagcg cagaagtggg cctgcaactt tatccgcctc catccagtct 2280
attaattgtt gccgggaagc tagagtaagt agttcgccag ttaatagttt gcgcaacgtt 2340
gttgccattg ctacaggcat cgtggtgtca cgctcgtcgt ttggtatggc ttcatcagc 2400
tcgggttccc aacgatcaag gcgagttaca tgatcccca tgttggtgaa aaaagcgggt 2460
agtcctctcg gtcctccgat cgttgtcaga agtaaagttg ccgcagtgtt atcactcatg 2520
gttatggcag cactgcataa ttctcttact gtcatgccat ccgtaagatg cttttctgtg 2580
actggtgagt actcaaccaa gtcattctga gaatagtgtg tgccggcgacc gagttgctct 2640
tgcccggcgt caacacggga taataccgcg ccacatagca gaactttaaa agtgctcatc 2700
attgaaaaac gctcttcggg gcgaaaactc tcaaggatct taccgctgtt gagatccagt 2760
tcgatgtaac ccactcgtgc acccaactga tcttcagcat cttttacttt caccagcgtt 2820
tctgggtgag caaaaacagg aaggcaaaat gccgaaaaa agggaataag ggcgacacgg 2880
aaatgttgaa tactcatact cttccttttt gaagcattta tcagggttat 2940
tgtctcatga gcgatatact atttgaatgt atttgaaaa ataaacaaat aggggttccg 3000
cgcacatttc cccgaaaagt gccacctggc ggccgcataa gatgatcttc ttgagatcgt 3060
tttggctcgc gcgtaatctc ttgctctgaa aacgaaaaaa ccgccttgca gggcggtttt 3120
tcgaaggttc tctgagctac caactctttg aaccgaggta actggcttgg aggagcgag 3180
tcaccaaaac ttgtccttcc agtttagcct taaccggcgc atgacttcaa gactaaactc 3240
tctaaatcaa ttaccagtgg ctgctgccag tgggtccttt gcatgtcttt ccgggttggg 3300
ctcaagacga tagttaccgg ataaggcgca gcggtcggg tgaacggggg gttcgtgcat 3360
acagtccagc ttggagcgaa ctgcctaccc ggaactgagt gtcaggcgtg gaatgagaca 3420
aacgcggcca taacagcgga atgacaccgg taaaccgaaa ggcaggaaac ggagagcgca 3480
cgaggggagc gccaggggaa acgcctggtg tctttatagt cctgtcgggt ttcgccacca 3540
ctgatttgag cgtcagattt cgtgatgctt gtcagggggg cggagcctat ggaaaaacgg 3600
ctttgcgcgc gccctctcac ttccctgtta agtatcttcc tggcatcttc caggaaatct 3660
ccgccccgtt cgtaagccat ttccgctcgc cgcagtcgaa cgaccgagcg tagcgagtca 3720
gtgagcgagg aagcggaata tatcctgtat cacatatctt gctgacgcac cgggtgcagc 3780
ttttttctcc tgccacatga agcacttcac tgacaccctc atcagtgcc aatagtaag 3840
ccagtgcggc cgcttcttcc ctgcgttatc ccctgattct gtggataacc gtattaccgc 3900
ctttgagtga gctgataccg ctgcgcgcag ccgaacgacc gagecgagcg agtcagtga 3960
cgagggaagc gaagagcgcc tgatgcggtg ttttctcctt acgcatctgt gcggtatttc 4020
acaccgata agatcccctc gacctgcagg catgcaagct tctgtaggtt tttaggcata 4080
aaactatatg atttaccctt aaatctttaa aatgccctt aaaattcaaa ataaaggcat 4140
ttaaaattta aatatttctt gtgataaagt ttgttaaaaa ggagtgggtt tatgactgtt 4200
atgtggttat cgattatagg tatgtggtt ttgattggaa tggcattttt tgctatcaag 4260
gttattaaaa ataaaaattg gaccacgcat tttatgccgag aaaatttatt gtgcgttgag 4320
aagaaccctt aactaaactt gcagacgaat gtcggcatag cgtgagctat taagccgacc 4380
attcgacaag ttttgggatt gttaagggtt ccgaggctca acgtcaataa agcaattgga 4440
ataaagaagc gaaaaaggag aagtcggttc agaaaaagaa ggatatggat ctggagctgt 4500
aatataaaaa ctttcttcaa ctaacggggc aggttagtga cattagaaaa ccgactgtaa 4560

```

```

aaagtacagt cggcattatc tcatattata aaagccagtc attaggccta tctgacaatt 4620
cctgaataga gttcataaac aatcctgcat gataaccatc acaaacagaa tgatgtacct 4680
gtaaagatag cggtaaatat attgaattac ctttattaat gaattttcct gctgtaataa 4740
tgggtagaag gtaattacta ttattattga tatttaagtt aaaccagta aatgaagtcc 4800
atggaataat agaaagagaa aaagcatttt cagggtatag tgttttggga aacaatttcc 4860
ccgaaccatt atatttctct acatcagaaa ggtataaatc ataaaactct ttgaagtcac 4920
tctttacagg agtccaaata ccagagaatg ttttagatac accatcaaaa attgtataaa 4980
gtggctctaa cttatcccaa taacctaaact ctccgtcgct attgtaacca gttctaaaaa 5040
ctgtatttga gtttatcacc cttgtcacta agaaaataaa tgcagggtaa aatttatatc 5100
cttcttgttt tatgtttcgg tataaaacac taatatcaat ttctgtggtt atactaaaaa 5160
tcgtttgttg gttcaaataa tgattaaata tctcttttct cttccaattg tctaaatcaa 5220
ttttattaaa gttcatttga tatgcctcct aaatttttat cttaaagtga tttaggaggc 5280
ttacttgtct gctttcttca ttagaatcaa tcctttttta aaagtcaata ttactgtaac 5340
ataaatatat attttaaaaa tatcccactt tatccaattt tcgtttgttg aactaatggg 5400
tgctttagtt gaagaataaa gaccacatta aaaaatgtgg tcttttgtgt ttttttaaag 5460
gatttgagcg tagcgaaaaa tccttttctt tcttatcttg ataataaggg taactattgc 5520
cggcgaggct agttaccctt aagttatttg tatgactggg ttttaagcga aaaaaagttg 5580
ctttttcgta cctattaatg tatcgtttta aatgactagt aaaaaacata catagaaagg 5640
ggaaaaagca acttttttta ttgtcatagt ttgtgaaaac taagttgttt ttatgtgtta 5700
taacatggaa aagtatactg agaaaaaaca aagaaatcaa gtatttcaga aatttattaa 5760
acgtcatatt ggagagaatc aaatggattt agttgaagat tgcaatacat ttctgtcttt 5820
tgtagctgat aaaactttag aaaaacagaa attatataaa gctaattctt gtaaaaatcg 5880
attttgcct gtctgtgctt ggagaaaagc tagaaaagat gcattgggtt tatctttgat 5940
gatgcaatat attaaagcgc aagagaaaaa ggagtttata tttttaactt tgactacacc 6000
taatgtaatg agtgatgaat tagaaaatga aataaaacgt tataataatt cttttagaaa 6060
acttataaag agaaaaaaag taggtagtgt tataaaggga tatgttcgta agttagagat 6120
tacatataat aaaaaaagag atgattataa tcctcatttt catgtgttaa ttgcagttaa 6180
taaatcgtat ttcacagata aaagatatta tattagccaa caagaatggg tagattttatg 6240
gcgtgatgta acgggcattt cagaaataac acaagttcaa gttcaaaaaa taagacaaaa 6300
taataataaa gaattatatg aaatggctaa gtattctggt aaagatagtg attatttaat 6360
aaataaatca aaaagtcttt gatgcatttt ataaatcact taaaggtaaa caggattatg 6420
tttattcagg attattttaa gaggctaaaa agaaattaaa aaatggggat ttagattact 6480
taaaagaaat tgatccaacc gtaatatatc tatcaaattt tttatatttg gaaacaaaaa 6540
gagtatttag ctagtgaact ttatgactta acagaacaag aaaaaagaga aattaatcac 6600
aaaatgatag acgaaatcga ggaagaacaa taacaaaata taagtgtctaa cagtcgtctg 6660
caagtttagt taagggttct tctcaacgca caataaattt tctcggcata aatgcgtggg 6720
ctaattttta tttttaataa ccttgatagc aaaaaatgcc attccaatac aaaaccacat 6780
acctataatc gataaccaca taacagtcac aaaaaccact ctttttaaca aactttatca 6840
caagaaatat tt                                     6852

```

<210> 54

<211> 16

<212> DNA

<213> Artificial Sequence

<220>

<223> OR1 operator sequence

<400> 54

tatcaccgca gaggta

16

<210> 55

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> OR2 operator sequence

<400> 55
taacaccgtg cgtgttg 17

<210> 56
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> OR3 operator sequence

<400> 56
tatcaccgca agggata 17

<210> 57
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> OL1 operator sequence

<400> 57
tatcaccgcc agtggtta 17

<210> 58
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> OL2 operator sequence

<400> 58
tatctctggc ggtgttg 17

<210> 59
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> OL3 operator sequence

<400> 59
tatcaccgca gatggtt 17

<210> 60
<211> 15
<212> DNA
<213> Artificial Sequence

<220>
<223> MalM operator sequence

<400> 60
cgcaaacggt tgcgt 15

<210> 61
<211> 14
<212> DNA

<213> Artificial Sequence

<220>

<223> MalX operator sequence

<400> 61

cgcaaacggtt ttgg

14

<210> 62

<211> 38

<212> DNA

<213> Escherichia coli

<400> 62

actttatcac tgataaaca acttatcagt gataaaga

38

<210> 63

<211> 38

<212> DNA

<213> Escherichia coli

<400> 63

actctatcat tgatagagtt ccctatcagt gatagaga

38

<210> 64

<211> 38

<212> DNA

<213> Plasmid pSC101

<400> 64

agcttatcat cgataagcta gtttatcaca gttaaatt

38

<210> 65

<211> 39

<212> DNA

<213> Salmonella

<400> 65

actctatcat tgataaggga actctatcaa tgataggga

39

<210> 66

<211> 38

<212> DNA

<213> Transposon Tn10

<400> 66

aatctatcac tgatagagta ccctatcatc gatagaga

38

<210> 67

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Trp O operator sequence

<400> 67

cgaactagtt aactagtacg

20

<210> 68

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Xyl O operator sequence

<400> 68

agttagtttg ttattaaat taaccaact

29

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.



THIS PAGE BLANK (USPTO)